A strangulated hernia as a result of ERCP

An 83-year-old woman attended for endoscopic retrograde cholangiopancreatography (ERCP) following an episode of acute jaundice. A computed tomography (CT) scan confirmed a dilated common bile duct with a distal filling defect that was later confirmed on magnetic resonance imaging (MRI) to be a biliary stone. She had undergone a right inguinal hernia repair 5 years previously and the CT scan suggested a possible recurrence (● Fig. 1).

Sphincterotomy was performed at ERCP and a 10-mm stone was extracted. On the first postoperative day, she developed a temperature of 38.5°C and produced a feculent vomit. A plain abdominal radiograph showed distended small-bowel loops. Her arterial blood gases were normal but her white cell count was elevated at $16.2 \times 10^9/L$, as was her C-reactive protein (CRP) at 268 mg/L. A repeat CT scan revealed an incarcerated inguinal hernia causing small-bowel obstruction (● Fig. 2).

She underwent an emergency laparotomy, which revealed the strangulated hernia (● Fig. 3a) with a segment of necrotic small bowel 270 cm from the ligament of Treitz (● Fig. 3b). This segment was resected using a linear stapler to create a stapled side-to-side anastomosis. The patient recovered well and was fit for discharge 4 days later.

During any endoscopic procedure visualization of the lumen is achieved by the insufflation of gas. However, because ambient “room air” is not well absorbed by the gastrointestinal tract, it must either be suctioned at the end of the procedure or be passed through the gastrointestinal tract as flatus [1]. ERCP can lead to prolonged examination time and overinsufflation [2]. More pertinently it may cause bowel hyperextension limiting the flexibility of the endoscope. It has previously been reported that this may cause barotrauma resulting in perforation or abdominal compartment syndrome [1]. Carbon dioxide is better absorbed by the bowel

Fig. 1 Image from a computed tomography (CT) scan in an 83-year-old woman who had presented with an acute episode of jaundice showing an incidental finding of a possible recurrent inguinal hernia in the right inguinal region (arrow).

Fig. 2 Image from a computed tomography (CT) scan performed following development of fever and vomiting 1 day after endoscopic retrograde cholangiopancreatography (ERCP) showing an incarcerated inguinal hernia (arrow) with distension of the proximal small bowel and collapse of the distal small bowel.

and has been shown to reduce pain both during and after the procedure, as well as sedation requirements and procedure and recovery times [3].

We suggest that the presence of any abdominal hernias is documented when carrying out a prolonged endoscopic procedure such as an ERCP. Also, in such cases judicious use of room air or carbon dioxide insufflation should be considered to prevent the conversion of an incarcerated to a strangulated hernia as in this case.

Endoscopy_UCTN_Code_CPL_1AK_2AJ

Competing interests: None

J. Horsnell¹, V. Pegna¹, A. Bell², A. Mickevicius¹

¹ Department of Surgery, Weston General Hospital, Weston-Super-Mare, Somerset, UK

² Department of Gastroenterology, Weston General Hospital, Weston-Super-Mare, Somerset, UK

Fig. 3 Images during laparotomy showing: **a** necrotic small bowel at the medial aspect of the previous inguinal hernia repair; **b** a knuckle of necrotic small bowel following reduction of the strangulated inguinal hernia.

References

- 1 Manser CN, Bauerfeind P, Gubler C. Iatrogenic complications in five patients with upper GI bleeding due to ambient air: Case series and literature review. *Case Rep Gastroenterol* 2012; 6: 197–204
- 2 Muraki T, Arakura N, Kodama R et al. Comparison of carbon dioxide and air insufflation use by non-expert endoscopists during ERCP. *Dig Endosc* 2012, DOI: 10.1111/j.1443-1661.2012.01344.x
- 3 Dellon E, Hawk J, Grimm I et al. The use of carbon dioxide for insufflation during GI endoscopy: a systematic review. *Gastrointest Endosc* 2009; 69: 843–849

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326452>
Endoscopy 2013; 45: E299–E300
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

V. Pegna, MBBS
 Department of General Surgery
 Weston General Hospital
 Grange Road
 Weston-Super-Mare
 Somerset
 BS23 4TQ
 UK
 Fax: +44-1934-636366
victoriapegna@hotmail.co.uk