Multidisciplinary laparo-endoscopic management of a subcardial gastric diverticulum

Fig. 1 Accidental finding of a suprenal mass (arrow) on a magnetic resonance scan of the spine of a 46-year-old obese woman with acute exacerbation of abdominal pain, recurring nausea and vomiting, and a history of chronic pain in the lumbar spine.

Fig. 2 Computed tomography scan showing gastric diverticulum (arrow).

Fig. 3 Preoperative oral contrast study showing gastric diverticulum (arrow).

A 46-year-old obese female patient presented in the emergency room with acute exacerbation of abdominal pain, recurring nausea and vomiting, and a history of chronic pain in the lumbar spine. An external report of recently taken magnetic resonance images described a suspicious left adrenal mass (● Fig. 1). The patient also had reflux symptoms and fetor ex ore. A history of rapid weight loss of 16 kg over the past 3 months and increasing fatigue raised suspicion for malignant disease. Assessment for pheochromocytoma was negative. A computed tomography (CT) scan excluded any malignancy and

showed subcardial pouching of the dorsal gastric wall (● Fig. 2). Oral contrast swallow (● Fig. 3) and endoscopy confirmed the diagnosis of a gastric diverticulum with an aperture of 35 mm, filled with food remnants. Due to the clinical relevance of this benign finding, the patient was referred to general surgery. Using an endoscopic rendezvous technique, laparoscopic resection was carried out with an EchelonFlex60 Endopath Stapler (Ethicon Endo-Surgery, Cincinnati, Ohio, USA) under visual control from both sides of the diverticulum (● Fig. 4). Histologic examination showed a full-thickness stomach specimen with slight chronic inflammation of otherwise normal gastric mucosa, indicative of a congenital diverticulum. Both the fetor and reflux symptoms ceased immediately after the surgery. A second contrast swallow on postoperative day 5 showed a patent suture and the patient was discharged thereafter.

Gastric diverticula are a rare entity of diverticula of the gastrointestinal tract. Due to their rarity, they are often mistaken for pathologies such as, most commonly, left adrenal masses [1,2]. Usually asymptomatic, they sometimes present with diffuse abdominal symptoms and gastric bleeding [3,4]. The preferred treatment is laparoscopic resection of the diverticulum, however, open approaches have also been reported [5]. A rendezvous technique as carried out in our department has not yet been described, but constitutes a safe treatment method.

Endoscopy_UCTN_Code_TTT_1AT_2AD

Competing interests: None

B. Globke¹, P. Fikatas¹, A. Beck², A. Adler³, S. C. Schmidt¹

¹ Department of General, Visceral and Transplantation Surgery, Campus Virchow-Klinikum, Charité – Universitätsmedizin Berlin, Berlin, Germany

² Department of Radiology, Campus Virchow-Klinikum, Charité – Universitätsmedizin Berlin, Berlin, Germany

³ Department of Interdisciplinary Endoscopy, Division of Gastroenterology and Hepatology, Campus Virchow-Klinikum, Charité – Universitätsmedizin Berlin, Berlin, Germany

Fig. 4 **a** Intraoperative endoscopic visualization of the diverticulum aperture (arrow). **b** Laparoscopic view of diverticulum locked in GIA stapler. **c** Intraoperative endoscopic visualization after removal of the diverticulum (arrow).

References

- 1 Noguera JJ, Benito A, Hernandez-Sastre C et al. Gastric diverticulum mimicking cystic lesion in left adrenal gland. *Urology* 2009; 73 (Suppl. 05): 997–998
- 2 Kadera R, Otsuka F, Inagaki K et al. Gastric diverticulum simulating left adrenal incidentaloma in a hypertensive patient. *Endocr J* 2007; 54 (Suppl. 06): 969–974
- 3 Perbet S, Constantin JM, Poincloux L et al. Gastric diverticulum: a rare cause of hemorrhagic shock. *Intensive Care Med* 2008; 34 (Suppl. 07): 1353–1354
- 4 Lajoie A, Strum WB. Gastric diverticulum presenting as acute hemorrhage. *Gastrointest Endosc* 2008; 67 (Suppl. 01): 175–176
- 5 Elliott S, Sandler AD, Meehan JJ et al. Surgical treatment of a gastric diverticulum in an adolescent. *J Pediatr Surg* 2006; 41 (Suppl. 08): 1467–1469

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326455>
Endoscopy 2013; 45: E138–E139
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

B. Globke

Charité Universitätsmedizin Berlin
 Klinik für Allgemein-, Viszeral-
 und Transplantationschirurgie
 Campus Virchow-Klinikum
 Augustenburger Platz 1
 13353 Berlin
 Germany
 Brigitta.Globke@charite.de