Suppression of Melanin Synthesis by the Phenolic Constituents of Sappanwood (*Caesalpinia sappan*)

Authors

Kaoru Mitani¹, Fumihide Takano^{1,2}, Tetsuro Kawabata¹, Ahmed E. Allam¹, Mayumi Ota¹, Tomoya Takahashi¹, Nobuo Yahagi¹, Chikai Sakurada², Shinji Fushiya², Tomihisa Ohta¹

Affiliations

¹ Pharmacognosy and Chemistry of Natural Products, School of Pharmacy and Pharmaceutical Sciences, Kanazawa University, Kanazawa, Japan

² Nihon Pharmaceutical University, Saitama, Japan

Key words

- *Caesalpinia sappan* Linn.
- Fabaceae
- antimelanogenesis
- brazilin
- mRNAs expression

Abstract

Sappanwood (*Caesalpinia sappan* Linn.) is used as an herbal medicine. It is sometimes used to treat skin damage or as a facial cleanser. In the present study, the methanol (MeOH) extract of sappanwood was found to inhibit melanin synthesis in cultured human melanoma HMV-II cells stimulated with forskolin, and six active compounds (**1–5** and **7**) were isolated from the extract along with a non-active compound (**6**). Compounds **2–7** were identified as sappanchalcone (**2**), 3'-deoxy-4-O-methylsappanol (**3**), brazilein, (**4**), brazilin (**5**), sappanol (**6**), and 4-O-methylsappanol (**7**). Compound **1** was a new compound, and its structure was determined to be (6aS,11bR)-7,11b-dihydro-6H-indeno[2,1-c]chromene-3,6a,10,11-tetrol by spectroscopic analyses. Among the six active

compounds, brazilin (**5**) (EC₅₀: 3.0 ± 0.5 μM) and 4-O-methylsappanol (**7**) (EC₅₀: 4.6 ± 0.7 μM) strongly suppressed melanin synthesis in HMV-II cells. Bioactive compounds showed moderate cytotoxicities against HMV-II cells with IC₅₀ values of 83.1 ± 4.0 μM (for **2**), 72.0 μM ± 2.4 (for **3**), 33.8 ± 1.1 μM (for **4**), 18.4 ± 0.8 μM (for **5**), and 20.2 ± 0.8 (for **7**), respectively. Brazilin (**5**) selectively suppressed the expression of mRNAs for tyrosinase-related protein (TYRP) 2 and tyrosinase but did not influence the expression of TYRP1. These results suggest that brazilin (**5**) is a new class of melanin inhibitor and that sappanwood could be used as a cosmetic material.

Supporting information available online at <http://www.thieme-connect.de/ejournals/toc/plantamedica>

received May 16, 2012
revised Sept. 29, 2012
accepted October 9, 2012

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1327897>
Published online November 15, 2012
Planta Med 2013; 79: 37–44
© Georg Thieme Verlag KG
Stuttgart · New York ·
ISSN 0032-0943

Correspondence

Associate Professor
Dr. Fumihide Takano
Nihon Pharmaceutical
University
10281 Komuro
Kita-Adachi-Gun,
362-0806 Saitama
Japan
Phone: + 81 487 21 11 55
(ext. 6040)
Fax: + 81 487 21 62 41
takano@nichiyaku.ac.jp

Introduction

Melanin is a biopigment synthesized in melanosomes. Melanosomes enclosing melanin polymers are transferred from melanocytes to surrounding keratinocytes in the human epidermis. The color of human skin is determined primarily by the quantity, type, and distribution of melanin in keratinocytes [1, 2]. Melanin has a critical role in the protection of the skin from solar ultraviolet (UV) radiation [3], but excess synthesis of melanin leads to hyperpigmentation disorders such as melasma, lentigo, and age spots [4]. Hence, the inhibitors of melanin biosynthesis have received considerable attention in clinical and cosmetic research, and melanin inhibitors such as hydroquinone [5], kojic acid [6, 7], arbutin [8, 9], and linoleic acid [10, 11] have been discovered.

“Sappan Lignum” is the heartwood of sappanwood (*Caesalpinia sappan* Linn.; Fabaceae) and is used both as a dyestuff and as an herbal medicine

[12]. It has been used to treat wounds, skin diseases, leprosy, dysentery, menorrhagia, menoxenia, leukorrhea, and diabetic complications [13]. Several pharmacological studies have been undertaken focusing on the extracts and constituents of this plant as well as on the various activities found (e.g., immunomodulatory, [14], anti-inflammatory [15], antioxidant [16], hypoglycemic [17], vasorelaxant [12], spasmolytic [18], and anti-arthritis [19] effects). However, the effect on melanin synthesis of the extracts and constituents of this plant has not been studied in detail. In the present study, we investigated the inhibitory effects of the methanol (MeOH) extract of sappanwood on melanin synthesis in human melanoma HMV-II cells stimulated with forskolin. The MeOH extract showed inhibitory activity, and six compounds (one of which is novel) were isolated as active principles from the extract. Furthermore, the mechanism of action for the most potent inhibitory compound, brazilin **5**, was investigated.

Materials and Methods

General

Values for optical rotation were determined with a Horiba SEPA-3000 high-sensitivity polarimeter (Horiba). UV data were obtained with a Shimadzu UV-1600 UV-visible spectrometer (Shimadzu). IR data were obtained with a Shimadzu IR-460 IR spectrophotometer. Circular dichroism (CD) data were obtained using a JASCO J-820 spectrometer (JASCO).

NMR spectra were recorded with a JEOL GSX-500 spectrometer (JEOL) in acetone- d_6 and dimethyl sulfoxide (DMSO)- d_6 . Chemical shifts were referenced to the residual solvent peaks acetone- d_6 (δ_H 2.04 and δ_C 29.8) and DMSO- d_6 (δ_H 2.49 and δ_C 39.5). Mass spectra were measured on a JEOL SX-102 mass spectrometer. Reversed-phase high-performance liquid chromatography (HPLC) was performed on an ODS column (particle size: 5 μ m, TOSO, 18 mm $\times$ 250 mm). Silica gel (63–210 μ m; Kanto Kagaku) and ODS (63–212 μ m; Wako Pure Chemical Industries) were used for open-column chromatography. Thin-layer chromatography (TLC) was done on silica gel 60 F₂₅₄ (Merck) and RP-18 F_{254S} (Merck).

Plant material

C. sappan for medicinal use (lot number: 313116) was purchased from Uchida Wakanyaku Co. Ltd. and identified by the authors. A voucher specimen of the plant material (T-232) was deposited in the Laboratory of the Pharmacognosy and Chemistry of Natural Products, School of Pharmacy and Pharmaceutical Sciences, Kanazawa University (Kanazawa, Japan).

Extraction and isolation

The air-dried heartwood of *C. sappan* (5.0 kg) was extracted thrice with MeOH (3 L) at room temperature. The MeOH solution (9 L) was passed through filter paper (Advantec MFS, Inc.), and the solvent removed under reduced pressure at 35 °C to yield the MeOH extract (408 g). The MeOH extract (390 g) was suspended in H₂O and partitioned successively with *n*-hexane (1 L $\times$ 3), ethylacetate (EtOAc) (1 L $\times$ 5), and *n*-butanol (*n*-BuOH) (1 L $\times$ 3), respectively, to give *n*-hexane (9.46 g), EtOAc (345 g), *n*-BuOH (17.4 g), and H₂O (22.7 g) fractions.

The EtOAc fraction was further separated by column chromatography (80 mm $\times$ 200 mm) on SiO₂, eluted with a gradient of *n*-hexane and EtOAc (*n*-hexane:EtOAc = 5:5 $\rightarrow$ 4:6 $\rightarrow$ 2:8 $\rightarrow$ EtOAc) and MeOH to afford six fractions. The third fraction (54.0 g, eluted by *n*-hexane:EtOAc = 2:8) was subjected to additional column chromatography on SiO₂ and eluted with a gradient of chloroform and methanol (CHCl₃:MeOH = 14:1 $\rightarrow$ 9:1 $\rightarrow$ 7:1 $\rightarrow$ 5:1 $\rightarrow$ MeOH) to give nine fractions. A part of the fraction eluted by CHCl₃:MeOH (14:1) (20 mg) was purified by HPLC on a SiO₂ column (TOSO; 18 mm $\times$ 250 mm; particle size, 5 μ m; flow rate, 5 mL/min) with the eluting solvent (CHCl₃:MeOH = 97:3) to give compound **2** (6.04 mg; purity > 91%). A further portion of the same fraction (80 mg) was also separated by HPLC on a SiO₂ column with the eluting solvent (CHCl₃:MeOH = 14:1) to give compound **3** (2.21 mg; purity > 90%). A further portion of the same fraction (850 mg) was subjected to ODS column (80 mm $\times$ 200 mm) chromatography with elution with a gradient of aqueous acetonitrile solution (10% $\rightarrow$ 25% $\rightarrow$ 40% $\rightarrow$ 70% $\rightarrow$ 100%) to give seven fractions. The fraction eluted by 25% aqueous acetonitrile was purified by HPLC on an ODS column by elution with the same aqueous acetonitrile solvent to afford compound **4** (1.79 mg; purity > 90%). The fraction eluted by CHCl₃:MeOH (14:1) (3.7 g) was separated further by HPLC on a preparative

ODS column with the eluting aqueous acetonitrile solvent system (10% $\rightarrow$ 25% $\rightarrow$ 40% $\rightarrow$ 100%). Compound **5** (2.99 g; purity > 90%) was purified from the fraction eluted by 10% aqueous acetonitrile. Compounds **6** (3.97 mg; purity $\approx$ 92%) and **7** (16.0 mg; purity > 92%) were separated from the fraction eluted by 25% aqueous acetonitrile. A new compound **1** (15 mg; purity > 93%) was obtained from the fraction eluted by 40% aqueous acetonitrile. Compounds **2–7** (Fig. 1) were isolated and identified as sappanchalcone (**2**), 3'-deoxy-4-O-methylsappanol (**3**), brazilein (**4**), brazilin (**5**), sappanol (**6**), and 4-O-methylsappanol (**7**) by comparison of their spectral data with those reported in the literature [20–24].

Compound 1: orange needles, $[\alpha]_D^{25} + 65.0$ (c 0.75, DMSO); UV (MeOH) λ_{max} (log ϵ) 282 (3.46), 203 (4.31) nm; FABMS (negative) m/z 285 [$M - H$][−]; HRFABMS (negative) m/z 285.0789 [$M - H$][−] (calcd. for C₁₆H₁₅O₅, 285.0763). CD ($c = 2.0 \times 10^{-5}$ M, MeOH) $\Delta\epsilon$ (nm): −1.41 (282.0). ¹H NMR and ¹³C NMR data for **1** in acetone- d_6 are shown in Table 1.

Measurement of melanin content in cultured human melanoma HMV-II cells

Intracellular and extracellular melanin contents in cultured HMV-II human melanoma cells (Dainippon Pharmaceutical Co.) were determined according to methods described previously with minor modification [25–28]. Cells were passed once a week and used between passages 5 and 8. Briefly, cultured HMV-II cells were trypsinized [0.25% trypsin and 0.1% ethylenediamine tetraacetic acid (EDTA) at 37 °C for 5–10 min]. Cells [2.5×10^4 cells/well in 980 μ L Dulbecco's modified Eagle's medium (DMEM)] were inoculated with a pipette into 24-well plates and incubated for 24 h at 37 °C in a CO₂ incubator. After 24-h incubation, 20 μ L of sample solution was added to each well, and the 24-well plate incubated for 2 days at 37 °C in a CO₂ incubator. Each sample was tested in triplicate. Test samples were dissolved in dimethyl sulfoxide/phosphate-buffered saline (DMSO/PBS) and then diluted with DMEM to achieve the appropriate concentrations. The final concentration of DMSO was 0.03%. In the control group, DMSO/PBS (1:1, v/v) solution diluted with DMEM to 0.03% of the final DMSO concentration was used instead of the sample solution. We used α -arbutin (purity $\geq$ 98%; Sigma-Aldrich) or kojic acid (purity = 98.0%; Sigma-Aldrich) as the reference compounds for the inhibition of intracellular and extracellular melanogenesis. HMV-II cells were incubated with tested samples, reference compounds, or vehicle in the absence or presence of phorbol-12-myristate-13-acetate (PMA; an activator of protein kinase C) or forskolin (an activator of adenylate cyclase) [29] for 120 h. At the indicated time points, the culture medium was removed with a pipette and assayed for extracellular melanin, as described below. The remaining melanoma cells were trypsinized (0.25% trypsin and 0.1% EDTA at 37 °C for 5–10 min) and washed with 100 μ L PBS. Cells were digested by the addition of 400 μ L 1 N NaOH and left to stand for 16 h at room temperature. The optical density (OD) of the resulting solution was measured at 475 nm, and the amount of intracellular melanin calculated. The culture medium was centrifuged (600 $\times g$ for 10 min at 4 °C). An aliquot of the supernatant (1 mL) was added to 1 mL of a mixture of 0.4 M 2-[4-(2-hydroxyethyl)-1-piperazinyl] ethanesulfonic acid (HEPES) buffer (pH 6.8) and ethanol (9:1, v/v). The OD of the resulting solution was measured at 475 nm. The amount of melanin retained in the cells (intracellular melanin) and that secreted into cultured medium (extracellular melanin) were determined separately as described by Kawabata et al. [26].

Fig. 1 Chemical structures of isolated compounds from sappanwood.

Position	δ_C		δ_H (J in Hz)	HMBC
1	132.8	CH	7.72 d (8.2)	2, 4, 4a
2	110.2	CH	6.35 dd (2.0, 8.9)	3, 4
3	157.4	qC		
4	103.6	CH	6.25 d (2.0)	2, 3, 4a, 11c
4a	155.0	qC		
6	70.4	CH ₂	4.08 d (10.3), 3.85 d (11.0)	4a, 6a, 7, 11b
6a	79.4	qC	–OH 5.05 ^b	
7	41.7	CH ₂	3.21 d (15.8), 2.68 d (15.8)	6, 6a, 7a, 8, 9, 10, 11, 11b
7a	134.6	qC		
8	116.5	CH	6.47 d (8.2)	7, 10, 11, 11a, 11b
9	114.8	CH	6.62 d (8.5)	7a, 10, 11
10	144.2	qC		
11	143.3	qC		
11a	132.5	qC		
11b	51.1	CH	4.29 s	1, 4a, 6a, 7, 11, 11a, 11c
11c	117.8	qC		

Table 1 1H and ^{13}C NMR data for **1**^a.

^a 1H and ^{13}C NMR for **1** were measured in acetone- d_6 , ^b For detection of the hydroxyl signal of position 6a; 1H NMR for **1** was measured in DMSO- d_6 .

Cytotoxicity assay

To determine the cytotoxicity of the tested samples, HMV-II cells (180 μ L) were seeded on 96-well plates at 1.0×10^5 cells per well with tested samples (purity > 93%; 20 μ L in DMSO/PBS) at various concentrations. After 48-h cultivation, supernatants were removed, and adherent (B16F1) or non-adherent (THP-1) cells were incubated with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide (MTT; 10 μ L, 5 mg/mL in PBS) for 4 h and then solubilized with 10% (w/v) sodium dodecyl sulfate [SDS; in

60% (v/v) dimethyl formamide] solution (100 μ L) for 10 h. Absorbance was measured at 570 nm using a microplate reader (Bio-Rad Laboratories, Model 550). Cell viability was calculated by comparing the absorbance with that of the non-treated control culture. A cell growth curve was plotted. Values for the half-maximal inhibitory concentration (IC_{50}) were calculated using simple linear regression.

Fig. 2 Time course and dose-dependent melanin synthesis in cultured human melanoma HMV-II cells triggered by forskolin or PMA. HMV-II cells were treated with various concentrations of PMA or forskolin. Culture supernatants were removed at the indicated times. Intracellular melanin contents (panels A, C, and D) were measured by the methods described in Materials and Methods. Cell viability (panel B) was also measured after administration of forskolin or PMA. Data are the mean $\pm$ SD of quadruple cultures. * $p < 0.05$; significantly different from individual control culture.

Reverse transcription-polymerase chain reaction (RT-PCR)

Total ribonucleic acid (RNA) was prepared directly from HMV-II cells administered with samples for 8 h using a commercially available kit (RNeasy Mini Kit; Qiagen GmbH). Isolated RNA was quantified by photometry at 260 nm/280 nm. The quality of RNA was determined by measuring the ratio of 18S/28S ribosomal band intensity in an ethidium bromide containing 1.5% agarose gel after electrophoresis. A PrimerScript One-step RT-PCR kit ver2 (Takara Bio, Inc.) was used for preparing complementary deoxyribonucleic acid (cDNA). First-strand cDNAs were synthesized by one cycle of reverse transcription at 50 °C for 30 min and one cycle of reverse transcriptase incubated at 94 °C for 2 min. Second-strand cDNAs synthesis and PCR amplification for tyrosinase as well as tyrosinase-related proteins 1 and 2 (TYRP1 and TYRP2, respectively) were carried out at 94 °C for 30 s, 53–57 °C for 30 s, and 72 °C for 1 min. For each combination of primers, the kinetics of the PCR amplification was determined beforehand and semi-quantitative PCR undertaken in the exponential range. The β -actin (ACT) gene was used as an endogenous internal standard and amplified with specific primers simultaneously. PCR products were subjected to electrophoresis in 1% agarose gel and visualized by ethidium bromide staining under UV irradiation. The primer sequences used were: TYR forward, 5'-AGGCAGAGGTTCTGTCTAGA-3' and TYR reverse, 5'-ATTGTGTCTCGTCTTTGAG-3'; TYRP1 forward, 5'-CATGATGGCGAGAGATGACG-3' and TYRP1 reverse, 5'-TCTGTGAAGGTGTGCAGGAG-3'; TYRP2 forward, 5'-AGTGATTCGGCACGAACATCC-3' and TYRP2 reverse, 5'-CTGGAGGGAAGAAAGGAACC-3'; ACTB forward, 5'-AGAGCTACGAGCTGCCTGAC-3' and ACT reverse, 5'-AGCACTGTGTTGGCGTACAG-3'.

Data analyses

Mean melanin production ($\mu\text{g}/\text{mL}$) and the percentage of control of melanogenesis are reported as the mean $\pm$ SD of two or three independent experiments (triplicate/one experiment). Statistical significance was determined by Dunnett's multiple test after one-way analysis of variance (ANOVA) with comparison to a control group. $p < 0.05$ was considered statistically significant.

Supporting information

Spectroscopic data for compound **1** are available as Supporting Information.

Results

In the present study, we used human melanoma HMV-II cells instead of the previously used murine melanoma B16F1 cells [26] to assess the inhibitory effect on melanin synthesis. The potential of human melanoma HMV-II cells to synthesize melanin was therefore investigated. Melanin contents in cultured HMV-II cells increased depending upon the concentration of the stimulants forskolin, an activator of adenylate cyclase [30], and PMA, an activator of protein kinase C (● Fig. 2A). When stimulated by 500 ng/mL of forskolin, melanin contents increased markedly. Forskolin did not show toxicity at concentrations up to 500 ng/mL, whereas PMA decreased cell viability at > 50 ng/mL (● Fig. 2B). On the basis of these data, subsequent experiments were carried out using 500 ng/mL of forskolin as a stimulant.

The time course of intracellular melanin production in HMV-II cells is shown in ● Fig. 2D. Intracellular production of melanin

Fig. 3 Effects of kojic acid or arbutin on melanin synthesis in cultured HMV-II cells stimulated with forskolin. The positive control agents kojic acid or arbutin were added to the HMV-II cell culture system with or without forskolin (500 ng/mL), and the intracellular melanin contents measured at the indicated times (panel A and B). Dose–response relationships for the positive control agents were also determined 120 min after stimulation with forskolin (panel C). Panel D shows microscopic photographs of cells administered with kojic acid or arbutin for 96 h (1: no treatment, 2: forskolin only, 3: 100 µM arbutin + forskolin, and 4: 100 µM kojic acid + forskolin). Data are the means ± SD of quadruple cultures. * $p < 0.05$; significantly different from individual control culture.

increased appreciably 48 h after the challenge with 500 ng/mL forskolin, reached a peak at 96 h and decreased to the level seen in controls within 120 h. Thus, melanin contents were determined 96 h after forskolin stimulation in subsequent experiments.

Next, melanin contents in various cell concentrations were measured. Maximal synthesis of melanin was observed at a density of 2×10^5 cells/mL (stimulated by 500 ng/mL forskolin) (● Fig. 2C). At a density of 2×10^5 cells/mL, cell viability and melanin content decreased (data not shown). Hence, subsequent experiments were carried out using a density of 2×10^5 cells/mL.

The effects of arbutin and kojic acid [31] as positive control agents to inhibit melanin synthesis were assessed. Arbutin (● Fig. 3A) and kojic acid (● Fig. 3B) appreciably reduced the melanin synthesis induced by forskolin at the indicated time points. These inhibitions were concentration-dependent (● Fig. 3C). The whitening effect of arbutin and kojic acid could be visualized under a microscope (● Fig. 3D).

Thus, investigation of the inhibitory effect of *C. sappan* was carried out using the *in vitro* assay system under the conditions mentioned above. The MeOH extract inhibited melanin synthesis by 84.5% at a 10 µg/mL concentration (● Table 2). Furthermore, most of the activity was transferred into EtOAc and *n*-BuOH fractions. The H₂O fraction also inhibited melanin synthesis but showed marked cytotoxicity.

To identify the active constituent(s), the EtOAc fraction was further separated using SiO₂ or ODS column chromatography and HPLC. A new brazilin derivative (1) was isolated along with six known compounds (2–7) identified as sappanchalcone (2), 3'-deoxy-4-O-methylsappanol (3), brazilein (4), brazilin (5), sappanol (6), and 4-O-methylsappanol (7) by analyses of their NMR data and comparison of their spectroscopic data with those in the literature [20–24]. All six known compounds (2–7) were reportedly isolated from sappanwood [20–24].

Compound 1 showed a quasimolecular ion peak at m/z 285 [$M - H$][−]. The molecular formula was determined to be C₁₆H₁₃O₅ on the basis of HRFABMS and was identical to that of brazilin (5). The NMR data of 1 (● Table 1) were similar with those of brazilin (5) [22], except for two aromatic proton signals. Thus, aromatic proton signals due to ring D at δ 6.81 (1H, s, H-8) and δ 6.92 (1H, s, H-11) in brazilin (5) were observed as two doublets at δ 6.47 (1H, d, $J = 8.2$ Hz, H-8) and δ 6.62 (1H, d, $J = 8.5$ Hz, H-9) in 1, which indicated that the location of the OH group at C-9 in brazilin (5) was at C-11 in 1. This was supported by analyses of the heteronuclear multiple bond correlation (HMBC) (● Table 1 and Fig. 4A) and also by the correlated spectroscopy (COSY) spectrum (correlation from H-8: δ 6.47 to H-9: δ 6.62 was observed). A significant nuclear Overhauser effect (NOE) correlation (● Fig. 4B) was also observed between H-11b and the OH group (C-6a), indicating a *cis* relationship. Furthermore, the CD spectrum of 1 showed a negative Cotton effect at UV 282 nm (● Fig. 4C), similar to that of brazilin (5) (data not shown). Thus, compound 1 was determined to be (6*a*S,11*b*R)-7,11*b*-dihydro-6*H*-indeno[2,1-*c*]chromene-3,6*a*,10,11-tetrol.

After establishing the structures, the inhibitory activities of these compounds on melanin synthesis were investigated. Among the seven compounds, brazilin (5) and *O*-methylsappanol (7) strongly suppressed melanin synthesis in cultured HMV-II cells with concentration values that gave a half-maximal response (EC₅₀) of 3.0 ± 0.5 and 4.6 ± 0.7 µM, respectively. The five compounds (2, 3, 4, 5, and 7) showed cytotoxic effects, but selectivity factors (IC₅₀/EC₅₀) ranging from 1.43 (for 3), 1.82 (for 4), 1.95 (for 2), 4.39 (for 7), to 6.13 (for 5) were observed (● Table 3). At a lower concentration, 1 µM of brazilin (5) inhibited melanin synthesis by 53%, and the cell viability was 94%.

Interestingly, the inhibitory activity as judged by the half-maximal effective concentration (EC₅₀; 61.0 ± 3.1 µM) of the new compound (1), an isomer of brazilin (5), was >20-times weaker than

Table 2 Effects of MeOH extract and separated fractions on melanin synthesis in cultured HMV-II cells costimulated with forskolin.

Treatment	Yield (%)	Concentration ($\mu\text{g/mL}$)	Inhibition (%)	Cell viability (%)	n
MeOH extract	8.20 ^b	1	33.2 $\pm$ 1.9	94.3 $\pm$ 0.7	4
		10	84.5 $\pm$ 13.5 ^a	51.8 $\pm$ 3.1 ^a	3
<i>n</i> -hexane fr.	2.43 ^c	1	-2.0 $\pm$ 2.4	101 $\pm$ 2	4
		10	0.3 $\pm$ 1.2	99.1 $\pm$ 5.0	4
EtOAc fr.	88.5 ^c	1	42.7 $\pm$ 2.8 ^a	87.0 $\pm$ 4.2	4
		10	79.4 $\pm$ 9.0 ^a	44.4 $\pm$ 3.0 ^a	4
<i>n</i> -BuOH fr.	4.46 ^c	1	30.8 $\pm$ 2.6	89.0 $\pm$ 2.0	4
		10	78.3 $\pm$ 20.4 ^a	85.2 $\pm$ 4.8	4
H ₂ O fr.	5.82 ^c	1	50.4 $\pm$ 12.0 ^a	48.0 $\pm$ 1.2 ^a	4
		10	ND	10.0 $\pm$ 4.2 ^a	4
Arbutin	–	100 μM	58.7 $\pm$ 7.5 ^a	92.6 $\pm$ 2.9	3
Kojic acid	–	100 μM	76.1 $\pm$ 18.0 ^a	47.2 $\pm$ 6.6 ^a	4

HMV-II cells were treated with two concentrations of test samples for 96 h and costimulated with forskolin (500 ng/mL). After incubation, melanin contents and cell viabilities were measured. Data are the means percent of control $\pm$ SD of 3 or 4 independent experiments. ^a $p < 0.05$; significantly different from non-treated control culture. ^b Yield of the extract is indicated as percent (w/w) of dried plant. ^c Yield of each fraction was indicated as percent (w/w) of the MeOH extract. ND: not determined

Fig. 4 Key HMBC and NOE correlations and CD spectrum for **1**.

that of braziliin (**5**). Positive control agents arbutin and kojic acid showed moderate inhibitory activities with EC_{50} values of 123.3 ± 3.9 and $70.6 \pm 3.0 \mu\text{M}$, respectively (Table 3). The viability of HMV-II cells treated with kojic acid was decreased by 47% at 100 μM (Table 2).

To explore the mechanism of inhibition by the MeOH extract and active constituents on melanin synthesis, we first planned to investigate the inhibitory effect on tyrosinase in a cell-free condition. However, tyrosinase activity could not be analyzed because of the reddish color of the samples, which interfered with measurement of the OD of the chromophore formed by tyrosinase-catalyzed melanin. Therefore, the effects of the samples on the amounts of messenger ribonucleic acids (mRNA) molecules for tyrosinase and two related genes [TYRP1 and TYRP2/dopachrome tautomerase (DCT)] in HMV-II cells were examined using semi-quantitative RT-PCR. β -actin served as the housekeeping gene

through the experiment. The expression for three mRNAs encoding tyrosinase, TYRP1, and TYRP2/DCT was upregulated by adding 500 ng/mL forskolin for 8 h compared with that of the non-treated control group (Fig. 5A).

Arbutin and kojic acid (the positive control agents) reduced the gene expression for tyrosinase and TYRP1 induced by 50 μM forskolin but did not reduce the gene expression for TYRP2/DCT (Fig. 5). The MeOH extract at 10 $\mu\text{g/mL}$ concentration reduced the gene expression for tyrosinase but did not influence the mRNAs for TYRP1 and TYRP2/DCT (Fig. 5). Braziliin (**5**) reduced the expression of mRNAs for tyrosinase and TYRP2/DCT at the 10 μM concentration (Fig. 5A, B, and D) and reduced the latter more effectively (Fig. 5B and D). The MeOH extract and braziliin (**5**) did not reduce the expression of TYRP1 mRNA.

Discussion

The human melanoma HMV-II cells synthesized intracellular melanin within 96 h in response to forskolin without causing cytotoxicity. PMA also triggered melanin synthesis [32], whereas melanogenesis was lower than that seen for forskolin, and its action was accompanied by cytotoxicity. This finding suggested that forskolin was a more suitable stimulator than PMA for triggering melanogenesis in this culture system. The well-known whitening agents arbutin and kojic acid [31] appreciably suppressed melanin synthesis in HMV-II cells, suggesting that this assay system can be used to investigate inhibitors of melanin synthesis.

By using this assay system, the MeOH extract of sappanwood was found to inhibit melanin synthesis, and six active compounds were isolated. The five tested compounds (**2**, **3**, **4**, **5**, and **7**) and positive control agent (kojic acid) showed cytotoxicity against HMV-II cells. However, significant anti-melanogenic activities (EC_{50}) were observed when cells were treated with these samples at two- or four-times lower concentrations. Braziliin (**5**) exhibited strong suppression of melanogenesis ($\text{EC}_{50} = 3.0 \pm 0.5 \mu\text{M}$), and cell viability was $\leq 95\%$ at this concentration. Furthermore, 3'-deoxy-4-O-methylsappanol (**7**) also suppressed melanogenesis with an IC_{50} of $4.6 \pm 0.7 \mu\text{M}$, whereas nonsignificant cytotoxicity (cell viability $< 92\%$) was shown at this concentration. Therefore, the anti-melanogenic activity of the tested compounds was prob-

Compound	EC ₅₀ value for anti-melanogenic activity (μM)	IC ₅₀ value for cytotoxicities (μM)	n
1	61.0 ± 3.1	ND	8
2	42.6 ± 1.8	83.1 ± 4.0	8
3	50.4 ± 2.0	72.0 ± 2.4	8
4	18.6 ± 1.1	33.8 ± 1.1	8
5	3.0 ± 0.5	18.4 ± 0.8	8
6	> 100	ND	8
7	4.6 ± 0.7	20.2 ± 0.8	8
Arbutin	123.3 ± 3.9	ND	8
Kojic acid	70.6 ± 3.0	99.7 ± 2.1	8

Values are mean values from two independent experiments (n = 4/experiment). Cell viabilities were measured using the MTT method described in Materials and Methods, and IC₅₀ values for individual compounds against HMV-II cells were calculated. ND: not determined

Table 3 Inhibitory activities of isolated compounds on melanin synthesis in cultured human melanoma HMV-II cells costimulated with forskolin.

Fig. 5 Effects of the MeOH extract, braziliin (5), arbutin, and kojic acid on the expression of mRNAs related to melanin synthesis in cultured HMV-II cells. Cells were treated with test samples [kojic acid, 100 μM; arbutin, 100 μM; braziliin (5), 5 μM; MeOH extract, 10 μg/mL]. mRNAs were extracted 8 h after incubation, and the expression of tyrosinase (TYR), TYRP1, or TYRP2/DCT was measured by semiquantitative RT-PCR using specific primers (panel A). β-Actin gene expression (ACT) was used as an endogenous internal standard. Mean expression values for non-treated control cultures are normalized to the expression of ACT. Expression levels of TYR (panel B), TYRP1 (panel C), and TYRP2/DCT (panel D) are indicated as the means ± SD of three independent experiments.

ably not due to their cytotoxicities. We are currently investigating the cytotoxic activity of isolated compounds on cultured human skin melanocytes, keratinocytes, and fibroblast cells to clarify this issue.

In addition, compound 1, which is an analog of 5, moderately inhibited melanogenesis, suggesting that the position of the benzoyl hydroxyl group is important for inhibition.

To clarify the mechanism of inhibition of braziliin (5), semiquantitative RT-PCR analyses of tyrosinase-related mRNAs in HMV-II cells were carried out. Arbutin and kojic acid suppressed the expression of tyrosinase and TYRP1 mRNAs at 50 μM. Braziliin (5) (5 μM) reduced the expression of tyrosinase and TYRP2 mRNAs, but did not act upon TYRP1 mRNA. Lee et al. [33] reported that the flavanone butin inhibited melanin synthesis in human epidermal melanocytes and suppressed the expression of tyrosi-

nase, TYRP1, and TYRP2/DCT mRNAs. Therefore, the mechanism of inhibition of braziliin (5) is different from those of arbutin, kojic acid, and butin.

In the present study, we revealed the suppressive effect of the MeOH extract of sappanwood upon melanin synthesis in cultured human melanoma HMV-II cells and isolated six active compounds, one of which, compound 7, was a new braziliin analog. We also demonstrated that 5, which showed the most potent inhibition, selectively suppressed the expression of mRNAs for tyrosinase and TYRP2. These results support the traditional use of sappanwood for treating skin damage or as a facial cleanser. These compounds are candidates for new cosmetic agents.

Conflict of Interest

The authors declare no conflict of interest.

References

- 1 Parvez S, Kang M, Chung HS, Cho C, Hong MC, Shin MK, Bae H. Survey and mechanism of skin depigmenting and lightening agents. *Phytother Res* 2006; 20: 921–934
- 2 Wakamatsu K, Kavanagh R, Kadekaro AL, Terzieva S, Sturm RA, Leachman S, Abdel-Malek Z, Ito S. Diversity of pigmentation in cultured human melanocytes is due to differences in the type as well as quantity of melanin. *Pigment Cell Res* 2006; 19: 154–162
- 3 Solano F, Briganti S, Picardo M, Ghanem G. Hypopigmenting agents: an updated review on biological, chemical and clinical aspects. *Pigment Cell Res* 2006; 19: 550–571
- 4 Briganti S, Camera E, Picardo M. Chemical and instrumental approaches to treat hyperpigmentation. *Pigment Cell Res* 2003; 16: 101–110
- 5 Jimbow K, Obata H, Pathak MA, Fitzpatrick TB. Mechanism of depigmentation by hydroquinone. *J Invest Dermatol* 1974; 62: 436–449
- 6 Mishima Y, Ohyama Y, Shibata T, Seto H, Hatae S. Inhibitory action of Kojic acid on melanogenesis and its therapeutic effect for various human hyper-pigmentation disorders. *Skin Res* 1994; 36: 134–150
- 7 Mishima Y, Hatta S, Ohyama Y, Inazu M. Induction of melanogenesis suppression: cellular pharmacology and mode of differential action. *Pigment Cell Res* 1988; 1: 367–374
- 8 Maeda K, Fukuda M. Arbutin: mechanism of its depigmenting action in human melanocyte culture. *J Pharmacol Exp Ther* 1996; 276: 765–769
- 9 Sugai T. Clinical effects of arbutin in patients with chloasma. *Skin Res* 1992; 34: 522–529
- 10 Ando H, Ryu A, Hashimoto A, Oka M, Ichihashi M. Linoleic acid and alpha-linolenic acid lightens ultraviolet-induced hyperpigmentation of the skin. *Arch Dermatol Res* 1998; 290: 375–381
- 11 Ando H, Watabe H, Valencia JC, Yasumoto K, Furumura M, Funasaka Y, Oka M, Ichihashi M, Hearing VJ. Fatty acids regulate pigmentation via proteasomal degradation of tyrosinase: a new aspect of ubiquitin-proteasome function. *J Biol Chem* 2004; 279: 15427–15433
- 12 Sasaki Y, Suzuki M, Matsumoto T, Hosokawa T, Kobayashi T, Kamata K, Nagumo S. Vasorelaxant activity of sappan lignum constituents and extracts on rat aorta and mesenteric artery. *Biol Pharm Bull* 2010; 33: 1555–1560
- 13 Wu J, Hou B, Zhang MM, Zhou YP, Yu B. Protosappanin A, an immunosuppressive constituent from a Chinese herb, prolongs graft survival and attenuates acute rejection in rat heart allografts. *Transplant Proc* 2008; 40: 3719–3722
- 14 Ye M, Xie WD, Lei F, Meng Z, Zhao YN, Su H, Du LJ. Brazilin, an important immunosuppressive component from *Caesalpinia sappan* L. *Int Immunopharmacol* 2006; 6: 426–432
- 15 Hu CM, Liu YH, Cheah KP, Li JS, Lam CSK, Yu WY, Choy CS. Heme oxygenase-1 mediates the inhibitory actions of brazilin in RAW264.7 macrophages stimulated with lipopolysaccharide. *J Ethnopharmacol* 2009; 121: 79–85
- 16 Badami S, Moorkoth S, Rai SR, Kannan E, Bhojraj S. Antioxidant activity of *Caesalpinia sappan* heartwood. *Biol Pharm Bull* 2003; 26: 1534–1537
- 17 Parameshwar S, Srinivasan KK, Rao CM. Oral antidiabetic activities of different extracts of *Caesalpinia bonducella* seed kernels. *Pharm Biol* 2002; 40: 590–595
- 18 Rodriguez-Lopez V, Salazar L, Estrada S. Spasmolytic activity of several extracts obtained from some Mexican medicinal plants. *Fitoterapia* 2003; 74: 725–728
- 19 Wang YZ, Sun SQ, Zhou YB. Extract of the dried heartwood of *Caesalpinia sappan* L. attenuates collagen-induced arthritis. *J Ethnopharmacol* 2011; 136: 271–278
- 20 Namikoshi M, Nakata H, Saitoh T. Homoisoflavonoids from *Caesalpinia sappan*. *Phytochemistry* 1987; 26: 1831–1833
- 21 Namikoshi M, Nakata H, Saitoh T. Homoisoflavonoids and related compounds. V. A novel dibenzoxocin derivative from *Caesalpinia sappan* L. *Chem Pharm Bull* 1987; 35: 3615–3619
- 22 Kim DS, Baek NI, Oh SR, Jung KY, Lee IS, Lee HK. NMR assignment of brazilin. *Phytochemistry* 1997; 46: 177–178
- 23 Hulme AN, McNab H, Pegg DA, Quye A. Negative ion electrospray mass spectrometry of neoflavonoids. *Phytochemistry* 2005; 66: 2766–2770
- 24 Fu LC, Huang XA, Lai ZY, Hu YJ, Liu HJ, Cai XL. A new 3-benzylchroman derivative from sappan lignum (*Caesalpinia sappan*). *Molecules* 2008; 13: 1923–1930
- 25 Sugimoto K, Nishimura T, Nomura K, Sugimoto K, Kuriki T. Inhibitory effects of α -arbutin on melanin synthesis in cultured human melanoma cells and a three-dimensional human skin model. *Biol Pharm Bull* 2004; 27: 510–514
- 26 Kawabata T, Cui MY, Hasegawa T, Takano F, Ohta T. Anti-inflammatory and anti-melanogenic steroidal saponin glycosides from *Fenugreek* (*Trigonella foenum-graecum* L.) seeds. *Planta Med* 2011; 77: 705–710
- 27 Bhatnagar V, Srirangam A, Abburi R. *In vitro* modulation of proliferation and melanization of melanoma cells by citrate. *Mol Cell Biochem* 1998; 187: 57–65
- 28 Matsuda H, Kawaguchi Y, Yamazaki M, Hirata N, Naruto S, Asanuma Y, Kaihatsu T, Kubo M. Melanogenesis stimulation in murine B16 melanoma cells by *Piper nigrum* leaf extract and its lignan constituents. *Biol Pharm Bull* 2004; 27: 1611–1616
- 29 D'Orazio JA, Nobuhisa T, Cui R, Arya M, Spry M, Wakamatsu K, Igras V, Kunisada T, Granter SR, Nishimura EK, Ito S, Fisher DE. Topical drug rescue strategy and skin protection based on the role of Mc1r in UV-induced tanning. *Nature* 2006; 443: 340–344
- 30 Busca R, Ballotti R. Cyclic AMP: a key messenger in the regulation of skin pigmentation. *Pigment Cell Res* 2000; 13: 60–69
- 31 Callender VD, St Surin-Lord S, Davis EC, Maclin M. Postinflammatory hyperpigmentation: etiologic and therapeutic considerations. *Am J Clin Dermatol* 2011; 12: 87–99
- 32 Tatsuno T, Jinno M, Arima Y, Kawabata T, Hasegawa T, Yahagi N, Takano F, Ohta T. Anti-inflammatory and anti-melanogenic proanthocyanidin oligomers from peanut skin. *Biol Pharm Bull* 2012; 35: 909–916
- 33 Lee MH, Lin YP, Hsu FL, Zhan GR, Yen KY. Bioactive constituents of *Spatholobus suberectus* in regulating tyrosinase-related proteins and mRNA in HEMn cells. *Phytochemistry* 2006; 67: 1262–1270