Chemical Constituents from the Fungus *Amauroderma amoiensis* and Their *In Vitro* Acetylcholinesterase Inhibitory Activities

Shuang Shuang Zhang^{1,2}, Qing Yun Ma², Xi Sheng Zou³, Hao Fu Dai², Sheng Zhuo Huang², Ying Luo², Zhi Fang Yu¹, Huai Rong Luo³, You Xing Zhao²

¹ Food Science and Technology College, Nanjing Agricultural University, Nanjing, China

² Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou, China

³ State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China

Abstract

One new compound named amauroamoienin (**1**), together with thirteen known compounds (**2–14**), was isolated from the EtOAc extract of *Amauroderma amoiensis*. The structures of these compounds were elucidated by the analysis of 1D and 2D spectroscopic data and the MS technique. The bioassays of inhibitory activities of these isolates against acetylcholinesterase were evaluated, and compounds **1**, **3**, and **5** exhibited acetylcholinesterase inhibitory activities.

Key words

Amauroderma amoiensis · Ganodermataceae · amauroamoienin · acetylcholinesterase inhibitory activity

Supporting information available online at <http://www.thieme-connect.de/ejournals/toc/plantamedica>

The family Ganodermataceae comprising more than 200 fungal species is mainly distributed in the tropical and subtropical areas of Asia, Australia, Africa, and America [1]. More than 100 species of this family grow in China, of which 78 wild species were found in Hainan Province [2]. *Ganoderma* and *Amauroderma* are two genera of the Ganodermataceae family. A great deal of work has been carried out on the genus *Ganoderma* [3]. Two of its species, recorded in the Chinese Pharmacopoeia, *G. lucidum* and *G. sinense*, have been used for the treatment of migraine, hypertension, arthritis, bronchitis, asthma, gastritis, diabetes, nephritis, and hepatitis problems for centuries [4,5]. Recent research on the chemical constituents of *Ganoderma* species showed the presence of natural products including triterpenes, steroid, alkaloids, flavonoids, polysaccharides, and fatty acids [6–8]. Previous screening of acetylcholinesterase (AChE) inhibitors from fungal extracts showed that many fungi, including the genera of the Ganodermataceae family, exhibited inhibitory activities against AChE [9]. At present, chemical constituents from many *Amauroderma* species, including *Amauroderma amoiensis* (Zhao) found in Fujian and Hainan Province of China [10], and their biological activities are still not reported.

In order to study the bioactive constituents of *A. amoiensis*, a chemical investigation was carried out and led to the isolation of one new compound, named amauroamoienin (**1**), along with

thirteen known compounds (**2–14**) (● Fig. 1), 4-hydroxy-17-methylincisterol (**2**) [11], (17R)-17-methylincisterol (**3**) [12], 1,5-dihydroxy-6',6'-dimethylpyrano[2',3':3,2] xanthone (**4**) [13], jacareubin (**5**) [13], ergosterol peroxide (**6**) [14], ergosta-7,22-dien-3 β -ol (**7**) [15,16], (22E,24R)-ergosta-8,22E-diene-3 β ,5 α ,6 β ,7 α -tetraol (**8**) [17,18], 22E-7 α -methoxy-5 α ,6 α -epoxyergosta-8 (14),22-dien-3 β -ol (**9**) [19], 3 β ,5 α ,9 α -trihydroxyergosta-7,22-dien-6-one (**10**) [20,21], 1H-indole-3-carboxylic acid (**11**) [22], *p*-hydroxybenzoic acid (**12**) [23], methyl 3,4-dihydroxybenzoate (**13**) [24], and 7,8-dimethylalloxazine (**14**) [25]. Herein, we describe the isolation and structural elucidation of the new compound (**1**), as well as the AChE inhibitory activities of all isolates. Compound **1** was obtained as a yellow amorphous powder, and its molecular formula was assigned to be C₄₆H₅₄O₇ from its HREIMS (*m/z* 718.3871 [M]⁺, calcd. for C₄₆H₅₄O₇, 718.3870) and NMR data (● Table 1), indicating twenty degrees of unsaturation. The IR spectrum displayed the presence of hydroxyl (3353 cm⁻¹), carbonyl (1726 cm⁻¹), and double bond (1621 cm⁻¹) absorptions. Analysis of ¹³C NMR and DEPT spectra (● Table 1) showed 46 carbon resonances including eight for methyls, six for methylenes, sixteen for methines (two oxygenated), and sixteen for quaternary carbons (including one carbonyl). The ¹³C NMR spectra showed resonances characteristic of the structure consisting of an ergostane moiety and a xanthone moiety. Comparison of 18 carbon signals of compound **1** with those of jacareubin (**5**) [14] hinted that **1** had a xanthone moiety with the same structure of that of **5**, which was also confirmed by olefinic and aromatic protons signals [δ_H 6.71 (1H, d, *J* = 10.1 Hz), 5.57 (1H, d, *J* = 10.1 Hz), δ_H 6.39 (1H, s), 6.88 (1H, d, *J* = 8.9 Hz), and 7.67 (1H, d, *J* = 8.9 Hz)], and two resonance signals [δ_H 1.45 and 1.46 (each 1H, s)] for methyl groups similar to those of jacareubin. The ¹H-¹H COSY correlations of H-7'/H-8' and H-13'/H-14' and HMBC correlations (● Fig. 2) further deduced the presence of a jacareubin moiety in **1**. The remaining 28 carbon signals in compound **1** were reasonably characteristic of an ergostane steroid skeleton which was the basic natural product isolated from this genus. The steroid moiety possessed six olefinic carbons [δ_C 115.9 (d), 124.0 (d), 132.5 (d), 135.6 (d), 140.0 (s), 141.0 (s)], two oxygenated methines [δ_C 66.8, 73.6], and one oxygenated quaternary carbon [δ_C 77.3], among which an epoxy group was formed based on analysis of its molecular formula. Comparison of these ¹³C NMR data with those of (22E,24R)-ergosta-7,9(11),22-triene-3 β ,5 α ,6 β -triol [26] suggested that they might have identical skeletons except for the difference of three oxygenated carbons. The key HMBC correlations from H-6 (δ_H 4.73, s) to C-6', C-7, and C-8 indicated that the steroid and xanthone moieties were connected via an oxygen at C-6 and C-6', which also confirmed the epoxy group formed at two other oxygenated carbons. The HMBC correlation from H-19 (δ_H 1.24, s) to C-5 (δ_C 77.3) showed that the epoxy group was at C-4 and C-5. The stereochemistry of the chiral centers (C-10, C-13, C-14, C-17) and the side chain at C-20 and C-24 of the steroid moiety in **1** were proposed to be the same as those of (22E,24R)-ergosta-7,9(11),22-triene-3 β ,5 α ,6 β -triol by comparison of their ¹³C NMR data, with the possible β -orientations of CH₃-18 and CH₃-19 and α -orientations of H-14 and H-17 [26]. The β -orientation of H-6 was determined by the key NOE of H-6/H-19 (● Fig. 2). The weak correlations of H-6 and H-4 (δ_H 3.93, m) with the same proton H-2b (δ_H 1.61, m) proposed the β -orientation of H-4, which accordingly hinted the α -orientation of the epoxy group in **1**. NOE of H-18/H-20 further confirmed

Fig. 1 Structures of compounds 1–14.

the β -orientation of CH₃-18 and the α -orientation of H-17. Thus, the structure of compound **1** was assigned as shown in **Fig. 1**, and this compound was named amauroamoienin.

The AChE inhibitory activities for compounds **1–11** were tested. The purities of all compounds were more than 98% by HPLC analyses. Compound **3** showed a certain inhibitory activity (inhibition percentage was 46.33%) at the concentration of 100 μ M, and compounds **1** and **5** showed weak inhibitory activity (inhibition percentages were 14.63% and 25.49%, respectively) (**Table 2**). Meanwhile, the other compounds were inactive with inhibition ratios less than 10%.

Materials and Methods

The fruiting bodies of *A. amoensis* were collected in Lingshui County, Hainan Province, People's Republic of China in June 2011 and identified by Prof. Xing-Liang Wu of the Hainan University. A voucher specimen (No. 2011JZ01) was deposited at the Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences.

Extraction and isolation: The dried and powdered fruiting bodies of *A. amoensis* (6.0 kg) were extracted at room temperature with 95% EtOH (3 $\times$ 20 L, 7 d each). The extract was concentrated and suspended in water followed by successive partition with EtOAc and *n*-BuOH, respectively. The EtOAc extract (70.8 g) was separated by silica gel column chromatography (CC) (10 $\times$ 40 cm, 560.0 g) under vacuum using a gradient eluent mixture of petroleum ether/EtOAc (20:1–3:1, 3 L each) to afford nine fractions (Fr 1–Fr 9). Fraction 4 (3.2 g) was subjected to silica gel CC (4 $\times$ 30 cm, 30.0 g) under vacuum eluted with petroleum ether/EtOAc (10:1–3:1, 1 L each) to give 6 subfractions 4a–4f. Subfraction 4d (350.0 mg) was subjected to CC with Sephadex LH-20 (2 $\times$ 100 cm, CHCl₃/MeOH 1:1, 1 L) to yield compound **4** (5.0 mg). Fraction 5 (7.3 g) was subjected to silica gel CC (5.5 $\times$ 30 cm, 70.0 g) eluted with petroleum ether/EtOAc (4:1–1:1, 3 L each) to give 5 subfractions 5a–5e. Subfraction 5a (447.0 mg) was repeatedly purified by silica gel CC (2 $\times$ 45 cm, 24.0 g) eluted with petroleum ether/EtOAc (7:1, 3 L) and Sephadex LH-20 (2 $\times$ 100 cm, CHCl₃/MeOH 1:1, 500 mL) to yield compound **1** (11.9 mg). Subfraction 5b (2.4 g) was chromatographed on silica gel CC (3 $\times$ 30 cm, 20.0 g) eluted with petroleum ether/EtOAc

Table 1 ^1H NMR (500 MHz) and ^{13}C NMR (125 MHz) data of compounds **1** and **5** in CDCl_3 (δ_{H} in ppm, J in Hz).

1			5			
No	δ_{H}, J (Hz)	δ_{C}	No	δ_{H}	δ_{C}	δ_{C}
1	1.69 (m), 1.90 (m)	30.0	1'		160.5	159.5
2	1.28 (m), 1.61 (m)	23.2	2'		104.8	103.7
3	1.59 (m), 2.16 (m)	35.4	3'		157.9	156.9
4	3.93 (m)	66.8	4'	6.39 (s)	95.6	94.6
5		77.3	4a'		157.2	156.4
6	4.73 (s)	73.6	5'		130.2	132.5
7	5.05 (s)	115.9	6'		146.7	152.1
8		141.0	7'	6.88 (d, 8.9)	114.1	113.2
9		140.0	8'	7.67 (d, 8.9)	117.2	115.9
10		40.6	8a'		115.5	112.9
11	5.69 (d, 6.4)	124.0	9'		180.7	179.9
12	2.22 (m), 2.35 (m)	42.3	9a'		103.6	102.2
13		42.8	10a'		147.1	146.0
14	2.19 (m)	51.1	12'		78.3	78.1
15	1.71 (m), 2.24 (m)	28.9	13'	5.57 (d, 10.1)	127.6	128.1
16	1.69 (m), 2.05 (m)	31.0	14'	6.71 (d, 10.1)	115.8	114.5
17	1.27 (m)	56.2	15'	1.45 (s)	28.7	27.9
18	0.56 (s)	12.0	16'	1.46 (s)	28.6	27.9
19	1.24 (s)	24.5				
20	2.01 (m)	40.6				
21	1.00 (d, 6.6)	21.0				
22	5.11 (dd, 8.3, 15.3)	135.6				
23	5.20 (dd, 7.7, 15.3)	132.5				
24	1.92 (m)	43.1				
25	2.06 (m)	33.3				
26	0.79 (d, 6.8)	19.9				
27	0.81 (d, 6.8)	20.2				
28	0.89 (d, 6.8)	17.9

Fig. 2 Key HMBC and NOESY correlations of compound **1**.

(4:1, 4L) to yield compound **6** (19.6 mg). Subfraction 5c (169.2 mg) was separated by silica gel CC (1.2 × 34 cm, 9.0 g) eluted with petroleum ether/EtOAc (5:1, 3 L) to yield compound **7** (3.8 mg). Fraction 6 (3.7 g) was separated by silica gel CC (4 × 30 cm, 32 g) using as eluent petroleum ether/EtOAc (4:1–1:1, 2 L each) to afford subfractions 6a–6e. Subfraction 6e (671.7 mg) was gel-filtrated on Sephadex LH-20 (2 × 100 cm,

$\text{CHCl}_3/\text{MeOH}$ 1:1, 1.2 L) to yield compound **2** (10.9 mg). Subfraction 6d (332.0 mg) was gel-filtrated on Sephadex LH-20 (2 × 100 cm, $\text{CHCl}_3/\text{MeOH}$ 1:1, 500 mL) and then separated by silica gel column (1.2 × 34 cm, 8.0 g) using as eluent petroleum ether/EtOAc (15:1, 1 L) to yield compound **3** (8.9 mg). Fraction 7 (4.7 g) was separated by vacuum liquid column (4 × 30 cm, 50.0 g) using as eluent petroleum ether/EtOAc (3:2–1:2, 2 L) to afford subfractions 7a–7g. Subfraction 7c (520 mg) was subjected to Sephadex LH-20 (2 × 100 cm, $\text{CHCl}_3/\text{MeOH}$ 1:1, 800 mL) to yield compound **11** (5.2 mg). Subfractions 7a and 7b were separated by silica gel column to yield compounds **12** (8.1 mg), using as eluent $\text{CHCl}_3/\text{MeOH}$ (50:1, 800 mL), and **13** (8.1 mg) (2.5 × 40 cm, $\text{CHCl}_3/\text{MeOH}$ 22:1, 600 mL) from 7a (830 mg), as well as compounds **8** (21.8 mg), **9** (12.9 mg), and **10** (24.3 mg) (2.5 × 40 cm, $\text{CHCl}_3/\text{EtOAc}$ 50:1–14:1, 600 mL each) from 7b (790 mg). Fraction 8 (7.7 g) was separated by vacuum liquid column (5.5 × 30 cm, 90 g) using as eluent $\text{CHCl}_3/\text{MeOH}$ (35:1–10:1, 1.5 L each) to afford subfractions 8a–8g. Subfractions 8d (55.3 mg) and 8e (880.3 mg) were gel-filtrated on Sephadex LH-20 (2 × 100 cm, $\text{CHCl}_3/\text{MeOH}$ 1:1, 400 mL and 700 mL) to yield compounds **5** (13.0 mg) and **14** (5.3 mg), respectively.

Isolate: Amauroamoienin (1): Yellow amorphous powder; $[\alpha]_{\text{D}}^{27} + 12.5$ (c 3.2, MeOH); UV (MeOH) λ_{max} (log ϵ) 365 (1.35), 315 (1.56), 257 (3.02), 241 (2.77); IR (KBr) ν_{max} $\cdot\text{cm}^{-1}$ 3353 cm^{-1} , 2885, 2270, 1726, 1621, 1590, 1490, 1407, 1296, 1253, 1195, 1093, 884, 888, 670; ^1H and ^{13}C NMR data, see **Table 1**; positive ESIMS m/z $[\text{M} + \text{H}]^+$ 719 (100); HREIMS m/z $[\text{M}]^+$ 718.3871 (calcd. for $\text{C}_{46}\text{H}_{54}\text{O}_7$, 718.3870).

Bioassay of AChE inhibitory activity: Acetylcholinesterase inhibitory activity of all compounds was assayed by the spectrophoto-

Compound	Percentage of inhibition	Compound	Percentage of inhibition
1	14.6	7	< 10
2	< 10	8	< 10
3	46.3	9	< 10
4	< 10	10	< 10
5	25.5	11	< 10
6	< 10	Tacrine*	53.0

* Positive control (0.333 μ M)

Table 2 AChE inhibitory activity of compounds **1–11** (%) at 50 μ M.

metric method developed by Ellman et al. [27] with slight modifications. *S*-Acetylthiocholine iodide, 5,5'-dithio-bis-(2-nitrobenzoic) acid (DTNB, Ellman's reagent), and acetylcholinesterase derived from human erythrocytes were purchased from Sigma Chemical. Compounds were dissolved in DMSO. The reaction mixture (totally 200 μ L), containing phosphate buffer (pH 8.0), test compound (50 μ M), and acetylcholinesterase (0.02 U/mL), was incubated for 20 min (30 °C). The reaction was initiated by the addition of 20 μ L of DTNB (0.625 mM) and 20 μ L acetylthiocholine iodide (0.625 mM) for the AChE inhibitory activity assay. The hydrolysis of acetylthiocholine was monitored at 405 nm after 30 min. Tacrine (Sigma-Aldrich 99%) was used as a positive control with final concentration of 0.333 μ M, and DMSO was used as negative control with final concentration of 0.1%. All the reactions were performed in triplicate. The percentage of inhibition was calculated as follows: % inhibition = $(E-S)/E \times 100$ (*E* is the activity of the enzyme without test compound and *S* is the activity of enzyme with test compounds).

Supporting information

The original spectra of NMR and HREIMS data for the new compound (**1**) and general experimental procedures are available as Supporting Information.

Acknowledgements

This work was financially supported by the Natural Science Foundation of Hainan (211020) and the National Nonprofit Institute Research Grant of CATAS (ITBB110301; 1630052012014). The authors are grateful to the members of the analytical group of the Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, for the spectral measurements.

Conflict of Interest

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, and there is no professional or other personal interest of any nature or kind in any product, service, and/or company that could be construed as influencing the position presented in, or the review of the manuscript entitled.

References

- Wu XL, Dai YC, Lin LH. Study on the Ganodermataceae of China I. Guizhou Sci 2004; 22: 27–34
- Wu XL, Guo JR, Liao QZ, Xie SH, Xiao M. The resources and ecological distribution of Ganodermataceae in Hainan island. Mycosystema 1998; 17: 122–129
- Paterson RRM. *Ganoderma* – a therapeutic fungal biofactory. Phytochemistry 2006; 67: 1985–2001
- Sliva D. *Ganoderma lucidum* in cancer research. Leuk Res 2006; 30: 767–768
- Stanley G, Harvey K, Slivova V, Jiang J, Sliva D. *Ganoderma lucidum* suppresses angiogenesis through the inhibition of secretion of VEGF and TGF- β 1 from prostate cancer cells. Biochem Biophys Res Commun 2005; 330: 46–52
- Aryantha INP, Adinda A, Kusmaningati S. Occurrence of triterpenoids and polysaccharides on *Ganoderma tropicum* with *Ganoderma lucidum* as reference. Aust Mycol 2002; 20: 123–129
- Liu JQ, Wang CF, Li Y, Luo HR, Qiu MH. Isolation and bioactivity evaluation of terpenoids from the medicinal fungus *Ganoderma sinense*. Planta Med 2012; 78: 368–376
- Lee IS, Kim HJ, Youn UJ, Kim JP, Min BS, Jung HJ, Na MK, Hattori M, Bae KH. Effect of lanostane triterpenes from the fruiting bodies of *Ganoderma lucidum* on adipocyte differentiation in 3T3-L1 cells. Planta Med 2010; 76: 1558–1563
- Oinonen P, Mettälä A, Vuorela P, Hatakka A. Screening of acetylcholinesterase inhibitors from fungal extracts. Planta Med 2006; 72: P-041 DOI: 10.1055/s-2006-949841
- Zhao JD, Xu LW, Zhang XQ. Taxonomic studies on the family Ganodermataceae of China II. Mycosystema 1983; 2: 159–167
- Togashi H, Mizushima Y, Takemura M. 4-Hydroxy-17-methylincisterol, an inhibitor of DNA polymerase- α activity and the growth of human cancer cells *in vitro*. Biochem Pharmacol 1998; 56: 583–590
- Ciminiello P, Fattorusso E, Magno S, Mangon A, Pansini M. Incisterols, a new class of highly degraded sterols from the marine sponge *Dictyonella incisa*. J Am Chem Soc 1990; 112: 3505–3509
- Rukachaisirikul V, Ritthiwigrom T, Pinsa A, Sawangchote P, Taylor WC. Xanthenes from the stem bark of *Garcinia nigrolineata*. Phytochemistry 2003; 24: 1149–1156
- Westerman PW, Gunasekera SP, Sultanbawa S, Uvais M, Kazlauskas R. Carbon-13 n.m.r. study of naturally occurring xanthenes. Org Magn Reson 1977; 17: 631–636
- Wan H, Sun RQ, Wu DJ. Three sterols from *Gyroporus castaneus*. Nat Prod Res Dev 1999; 11: 18–20
- Keller AC, Maillard MP, Hostettmann K. Antimicrobial sterols from the fungus *Fomitopsis pinicola*. Phytochemistry 1996; 41: 1041–1046
- Ishikzuka T, Yaoita Y, Kikuchi M. Sterol constituents from the fruit bodies of *Grifola frondosa* (Fr.) S.F.Gray. Chem Pharm Bull 1997; 45: 1756–1760
- Sun Y, Tian L, Huang J, Li W, Pei YH. Cytotoxic sterols from marine-derived fungus *Pennicillium* sp. Nat Prod Res 2006; 20: 381–384
- Gao H, Hong K, Chen GD, Wang CX, Tang JS, Yu Y, Jiang MM, Li MM, Wang NL, Yao XS. New oxidized sterols from *Aspergillus awamori* and the endo-boat conformation adopted by the cyclohexene oxide system. Magn Reson Chem 2010; 48: 38–43
- Kawagishi H, Katsumi R, Sazawa T, Mizuno T, Hagiwara T, Nakamura T. Cytotoxic sterols from the mushroom *Agaricus blazei*. Phytochemistry 1988; 27: 2777–2779
- Yue JM, Chen SN, Lin ZW, Sun HD. Sterols from the fungus *Lactarium vollemus*. Phytochemistry 2001; 56: 801–806
- Lai GF, Zhu XD, Luo SD, Wang YF. Chemical constituents from *Elsholtzia rugulosa*. Chin Tradit Herb Drugs 2008; 39: 661–664

- 23 Liu JJ, Liu XK. Chemical constituents from edible part of *Pistacia chinensis*. Chin Tradit Herb Drugs 2009; 40: 186–189
- 24 Liu ZY, Luo DQ. Chemical constituents from *Trollius chinensis*. Chin Tradit Herb Drugs 2010; 41: 370–373
- 25 Kwon HC, Kim KR, Zee SD, Cho SY, Lee KR. A new indolinepeptide from *Peecilomyces* sp.J300. Arch Pharm Res 2004; 27: 604–609
- 26 Ishizuka T, Yaoita Y, Kikuchi M. Sterol constituents from the fruit bodies of *Grifola frondosa* (Fr.) S. F. GRAY. Chem Pharm Bull 1997; 45: 1756–1760
- 27 Ellman GL, Courtney KD, Andres VJ, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. Biochem Pharmacol 1961; 7: 88–95

received September 12, 2012
revised October 14, 2012
accepted October 21, 2012

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1327951>

Published online November 23, 2012

Planta Med 2013; 79: 87–91

© Georg Thieme Verlag KG Stuttgart · New York ·
 ISSN 0032-0943

Correspondence

Prof. Dr. You Xing Zhao

Institute of Tropical Bioscience and Biotechnology
 Chinese Academy of Tropical Agricultural Sciences
 Xueyuan road 4#
 Haikou 571101
 China
 Phone: + 86 8 98 66 98 90 95
 Fax: + 86 8 98 66 98 90 95
zhaoyouxing@itbb.org.cn

Prof. Dr. Huai Rong Luo

State Key Laboratory of Phytochemistry and Plant Resources in West China
 Kunming Institute of Botany
 Chinese Academy of Sciences
 Lanhei road 132#
 Kunming 650204
 China
 Phone: + 86 87 15 22 32 25
 Fax: + 86 87 15 22 32 25
luohuairong@mail.kib.ac.cn