Characterisation of Nox4 Inhibitors from Edible Plants

Authors

Philipp A. Kofler¹, Haymo Pircher¹, Susanne von Grafenstein², Thomas Diener¹, Monika Höll^{1,3}, Klaus R. Liedl³, Karsten Siems⁴, Pidder Jansen-Dürr^{1,3}

Affiliations

The affiliations are listed at the end of the article

Key words

- NADPH oxidase 4
- Nox4
- ROS
- inhibitors
- dehydrogenase domain
- broken cell assay

Abstract

NADPH oxidases transport electrons from cytosolic NADPH through biological membranes to generate reactive oxygen species. NADPH oxidase 4, broadly expressed in humans, is an interesting pharmacological target, since its activity is deregulated in several diseases, including pulmonary fibrosis, diabetic nephropathy, and cardiac hypertrophy. Whereas several candidate NADPH oxidase 4 inhibitors were recently described, most of these compounds are either unspecific or toxic. Here we set out to identify new NADPH oxidase 4 inhibitors from edible plants, in an attempt to decrease the number of hits with toxic side effects. We screened a compound library prepared from edible plants for new bioactives with the ability to inhibit the activity of NADPH oxidase 4. Using both cell-based and cell-free assays, we identified several compounds with significant inhibitory activity towards NADPH oxidase 4. For selected compounds, the activity profile towards NADPH oxidase 2 and NADPH oxidase 5 was established, and controls were carried out to exclude general reactive oxygen species scavengers. A number of promising NADPH oxidase 4 inhibitors from edible plants was identified and characterised. Several new chemical entities are disclosed which act as NADPH oxidase 4 inhibitors, and the efficacies of our best hits, in particular

several diarylheptanoids and lignans, are comparable to the best available pharmacological NADPH oxidase 4 inhibitors. These findings will provide valuable tools to study mechanisms of NADPH oxidase inhibition.

Abbreviations

AR:	Amplex Red
AUC:	area under the curve
CL:	chemiluminescence
DHE:	dihydroethidium
DPI:	diphenyleneiodonium chloride
EC ₅₀ :	effective concentration at 50% activity
ETC:	electron transport chain
HEK 293:	human embryonic kidney 293 cells
MBP:	maltose binding protein
NBT:	nitroterazolium blue chloride
Nox:	NADPH oxidase
Nox4DH:	Nox4 dehydrogenase domain
PBMC:	peripheral blood mononuclear cells
PMA:	phorbol 12-myristate 13-acetate
ROCS:	rapid overlay of chemical structures
ROS:	reactive oxygen species
SF9:	Spodoptera frugiperda insect cells

Supporting information available online at <http://www.thieme-connect.de/ejournals/toc/plantamedica>

received July 4, 2012
revised October 26, 2012
accepted Dec. 6, 2012

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1328129>
Published online January 23, 2013
Planta Med 2013; 79: 244–252
© Georg Thieme Verlag KG
Stuttgart · New York ·
ISSN 0032-0943

Correspondence

Dr. Pidder Jansen-Dürr
Institute for Biomedical Aging
Research
University of Innsbruck
Rennweg 10
6020 Innsbruck
Austria
Phone: + 43 5 12 58 39 19 44
pidder.jansen-duerr@uibk.ac.at

Introduction

ROS induce damage to biological systems and were shown to contribute to ageing in cells. Aberrant ROS levels are linked to a variety of cancers [1] and many other diseases, e.g., vascular disease [2–6]. Whereas ROS are obligatory byproducts of mitochondrial oxidative metabolism, Nox2, the founding member of the Nox family [7,8], was found responsible for the oxidative burst of phag-

ocytes, where ROS are deliberately produced by Nox2 to enforce pathogen elimination [7,8]. In humans, the Nox family consists of seven enzymes, i.e., five NADPH oxidases (Nox1 through Nox5; for review, see [7,9]) and two Nox homologues (Duox1 and Duox2) [10], along with several subunits and regulatory proteins. All Nox proteins are transmembrane proteins with six (Nox1–Nox5) or seven (Duox1,2) transmembrane domains. In each case, four conserved histidines

bind two heme groups that are essential for the electron transport from cytosolic NADPH through membranes to generate ROS. An integral part is also covered by the FAD and NADPH binding domains that are similarly conserved [2, 7]. Both Nox2, a multisubunit NADPH oxidase best known for its role in the host defense against pathogens, and Nox5 require activation by upstream stimuli, such as PMA (Nox2) or calcium agonists (Nox5) for full activity; in contrast, Nox4, in complex with its obligatory subunit p22^{Phox}, is constitutively active [7]. Nox4 was identified in kidney [11] and was later found to be expressed in a variety of different cell types [12–19], and regulation occurs primarily at the mRNA level [20]. Whereas altered Nox4 activity has been found in a large variety of diseases, including pulmonary fibrosis, diabetic nephropathy (for review, see [4]), and benign prostatic hyperplasia [21], experiments with Nox4^{-/-} mice revealed a protective role of Nox4 in the vascular system [22], and Nox4 deletion was shown to predispose to diet-induced obesity [23]. Given the pleiotropic effects of Nox4 on a variety of physiological and pathophysiological processes, isotype-specific inhibitors of Nox4 would be potentially very useful [2, 3, 24, 25]. Whereas extracts of *Piper sarmentosum* regulate expression of Nox4 [26], inhibition of Nox4 catalytic activity was not described for such extracts. Various efforts to search for new Nox4 inhibitors have been published in recent years, but so far there are no Nox4 inhibitors clinically available [21, 27], although preclinical studies seem to support potential benefits in certain settings [28]. Also most known inhibitors are either unspecific or toxic [29, 30]. However, there are attempts to chemically synthesise inhibitors from promising lead compounds, but the claimed effects so far remain uncertified [24, 25, 31–34]. This study aimed to identify formerly unknown Nox4 inhibitors from edible plants, in an attempt to satisfy ADMET criteria and avoid toxicity, which is becoming a high priority in this field [35].

Materials and Methods

Chemicals

All reagents were purchased from Sigma-Aldrich, unless stated otherwise.

Test compounds

The majority of the compounds used for the screening were provided as purified compounds in solid state by AnalytiCon Discovery. For details of sample preparation, see Supplementary Methods as Supporting Information. As control compounds, we used DPI (Sigma), VAS2870 (Vasopharm), as well as honokiol and magnolol (both kindly provided by Hermann Stuppner, Innsbruck University). All control compounds were provided at >95% purity.

Cell culture

Peripheral blood mononuclear cells were isolated fresh every day from volunteers and suspended in RPMI medium. Osteosarcoma cells (U2-OS) were acquired from ATCC, and human embryonic kidney (HEK 293 FT) cells were bought from Invitrogen. Both were grown in DMEM (Sigma) and provided with fresh medium every 2 and 3 days, respectively. DMEM was supplemented with 10% FCS (Biochrome AG, heat inactivated at 56°C for 40 min), 4 mM L-glutamine (Gibco), and 1/100 penicillin/streptomycin (Gibco). SF9 cells, a gift from Alexandra Lusser (Division for Molecular Biology, Medical University Innsbruck), were incubated in

SF9 II medium (Gibco) with 10% FCS, 1.4 mM glutamine, and pen/strep at 27°C.

Nox dependent chemiluminescence

Chemiluminescence was performed in white 96-well plates, tissue culture-treated, optical flat bottom (Matrix, Nunc, or Perkin Elmer), essentially as described before [36]. For details, see Supporting Information.

ROS-scavenging assessment by DHE staining

U2-OS cells were treated with rotenone and stained with dihydroethidium to assess ROS levels, as described [37]. For details, see Supporting Information.

Preparation of cell membranes and Nox4 *in vitro* assay

Membrane fractionation and Nox4 activity measurement was performed similar to a method previously described [24, 27, 34, 38]; for details, see Supporting Information.

Shape-based alignment

In a retrospective analysis of the hits in the Nox-dependent chemiluminescence assay, we performed a shape-based analysis applying ROCS (version 3.1.1. OpenEye Scientific Software. <http://www.eyesopen.com>), as described [39, 40]. For details, see Supporting Information.

Results and Discussion

To assess the activity of Nox4 in intact cells, HEK cells expressing high levels of p22^{Phox} were transfected with an expression vector for Nox4. Stable transfection of HEK cells led to a significant increase of the Nox4 protein and activity, determined by chemiluminescence, which was sensitive to inhibition by the general Nox inhibitor DPI (**Fig. 1S**). This system was applied to determine the inhibitory activity of plant-derived compounds, selected from a compound library containing 3557 purified compounds, as outlined in **Fig. 2S** and briefly described below. For the first screening, a structural diverse subset of compounds derived from edible plants was selected, with a focus on compounds inside Lipinsky's Rule of Five. Results obtained with the diversity set were used to select additional compounds for screening, based on chemical similarity. The third screening was done with compounds mostly not structurally related to the hit compounds.

Selected experimental data are shown in **Fig. 1** for three selected plant compounds, representative of the activity spectrum (weak, intermediate, strong) obtained for the 14 best hits. DMSO vehicle had no effect on chemiluminescence activity, which was however significantly reduced by DPI when added at increasing concentrations (**Fig. 1A**). Under these conditions, compound ACD005 strongly suppressed Nox4 activity when assessed at a concentration as low as 0.5 µM ("strong" inhibitory activity) with a clear dose dependency observed (**Fig. 1B**). Under the same conditions, ACD018 induced a clear-cut inhibition of Nox4 activity when added at a concentration of 7.5 µM ("weak" inhibitory activity) (**Fig. 1C, Table 1**), whereas ACD212 displayed significant inhibitory activity when assayed at a concentration of 2.5 µM ("intermediate" inhibitory activity), and residual inhibitory activity was still visible at a concentration of 0.5 µM (**Fig. 1D**). The results of all measurements are summarised in **Fig. 1E** for 14 selected plant-derived compounds and the pharmacological Nox4 inhibitor VAS2870 [41, 42], and approximative

Fig. 1 Identification of Nox4 inhibitors. **A–D** Nox4-dependent chemiluminescence measurements of transfected HEK 293 cells using controls (**A** DMSO, DPI) and substances with strong (**B** ACD005), medium (**C** ACD212), and weak (**D** ACD018) inhibitory effect. **E** The signal of selected inhibitors at different concentrations 10 min after the start of the experiment. All values were derived from 3 independent experiments.

EC₅₀ values were calculated (● **Table 1**). Apocynin, gomisin C, magnolol, and honokiol, described as potential Nox4 inhibitors in the literature [25,43,44], were included in our screening program; however, these compounds were at best moderately active in our assays. Thus, EC₅₀ values were 60 μM for apocynin, 25 μM for gomisin C, 13 μM for honokiol, and 150 μM for magnolol. Because of the relatively low inhibitory activity in comparison to our best hits, these compounds were not considered further. Appropriate controls were included to identify inhibitors specific for Nox4 activity. Thus, we set up a Nox2 activity screen based on normal PBMC isolated from healthy donors, in which Nox2 activity was stimulated by addition of PMA (● **Fig. 2A**), and compounds were applied at a concentration suitable for Nox4 inhibition (see above, ● **Table 1**). Some of the Nox4 inhibitors, such as ACD042, did not inhibit Nox2 to any visible degree; in contrast, compound ACD018 displayed Nox2 inhibitory activity close to that of DPI, whereas VAS2780 displayed intermediate inhibitory activity (● **Fig. 2B**). The relative Nox2 inhibition by selected compounds is displayed in ● **Fig. 2C**. To further address Nox specificity we also established a HEK-derived cell line constitutively expressing Nox5. Nox5 expression

Table 1 EC₅₀ of selected Nox4 inhibitors. This table summarises the calculated EC₅₀ values for several inhibitors, as obtained in experiments with HEK-Nox4 cells. All values were derived from 3 independent experiments with 3 data points each, except for * × 2 independent experiments with 3 data points each and ** 1 experiment with 3 data points.

Substance	Mean [μM]	± S
ACD 005	0.54	0.10
ACD 018*	7.42	6.59
ACD 042*	2.06	0.76
ACD 047	0.87	0.42
ACD 053	0.92	0.28
ACD 055**	1.88	0.26
ACD 056	1.62	1.12
ACD 063*	2.55	2.19
ACD 066	0.83	0.44
ACD 067	1.12	0.43
ACD 069	2.37	0.69
ACD 084	3.08	2.77
ACD 176**	1.56	0.16
ACD 212	1.59	0.52
VAS 2870	2.65	1.13

Fig. 2 Assessment of inhibitory activity towards Nox2. **A** Nox2-dependent chemiluminescence of PBMCs was measured after the addition of 2 µg/mL PMA in the presence of compounds at the indicated concentration. DMSO 0.1% was the vehicle control, DPI 10 µM was the positive control. **B** A selection of typical Nox2-CL curves over time. 10 µM DPI and 20 µM ACD018 show strong inhibition, 1 µM ACD005 and 20 µM VAS2870 show reasonably moderate inhibition, 20 µM ACD042 shows no inhibitory capacity similar to the vehicle control. **C** The chemiluminescence values of the above experiment at t = 10 min as a bar graph, comprising 14 Nox4 inhibitors, VAS2870, and standards (DMSO vehicle control, 10 µM DPI, untreated control). (Color figure available online only.)

was verified, and Nox5 activity was detectable in these cells after addition of ionomycin (Fig. 3S), as described before [45]. In this

Fig. 3 Effects of candidate compounds on mitochondrial ROS. **A** ROS measurement of U2-OS cells after treatment with 0.2 µM rotenone for 30 min (except for the untreated control). The rotenone control was otherwise untreated, 0.1% DMSO was used as the vehicle control. **B** DHE signal of 14 selected Nox4 inhibitors, the vehicle control (0.1% DMSO), and VAS2870 at the indicated concentrations. The untreated control (Panel A) has been subtracted as blank for all substances and the control in panel B.

system, Nox5 activity was completely inhibited by DPI. Compound ACD063 displayed intermediate inhibitory activity, and VAS2870 was inactive (Fig. 3S).

An important source of potentially confounding effects is the well-known ability of many plant-derived compounds to unspecifically quench ROS, in many cases by direct chemical interaction (for recent review of ROS quenching mechanisms, see [46]). Such compounds, often referred to as ROS scavengers, neutralise a large variety of chemical ROS entities and therefore are expected to negatively interfere with the activity of most if not all experimental systems described in this communication. To assess the ability of plant-derived compounds to quench mitochondrial ROS production [37], mitochondrial ROS production was stimulated in U2-OS osteosarcoma cells by the addition of rotenone, an inhibitor of ETC complex I [47]. Subsequently, cells were stained with DHE, a redox-sensitive fluorescent dye recorded at 590 nm [37]. Cells were co-stained with Hoechst 33342, and fluorescence was recorded at 465 nm. Fluorescence intensity of Hoechst 33342-stained cells is proportional to cell numbers and was used to normalise DHE fluorescence. Rotenone-induced signal was reduced to background by the addition of ascorbate (Fig. 3A), consistent with previous findings [37]. In this experimental system, the selected compounds had no significant activity as ROS scavengers, with the exception of compound ACD066.

Fig. 4 Nox4 inhibitor assay with Nox4-containing membrane fractions. **A** Representative experiment. *Inserts:* Membrane fractions were prepared from HEK 293 cells transiently transfected with Nox4 and empty vector control, and expression levels of Nox4 and p22^{phox} were determined by Western blot. *Main panel:* Membrane fractions (10 µg protein per well) were used to oxidize AR as described in Materials and Methods; DPI (100 µM) was added at the beginning of the measurement as indicated. In each case, the reaction was started by the addition of 160 µM NADPH, and fluorescence was recorded over 20 min. **B, C** AR measurements were performed using membrane fractions (10 µg protein per well) prepared from HEK 293 cells transiently transfected with Nox4 and empty vector control, as shown in panel A. All inhibitors were present at a concentration of 20 µM except for DPI control (100 µM). 0.1 % DMSO was the vehicle control. The Nox4-dependent oxidation of AR was started by the addition of 160 µM NADPH, and fluorescence was recorded over 20 min. **D** Summary of the signals at t = 10 min after the experiment was started. The stars indicate the significance of the difference between the given substance and the DMSO control from a two-sided Student's test (* p < 0,05; ** p < 0,01; *** p < 0,001). (Color figure available online only.)

Our finding that ACD066 reduced the rotenone-induced DHE signal by roughly 50% suggests a more general mechanism rather than specific inhibition of NADPH oxidase activity for ACD066, which was therefore considered as a general ROS scavenger. Interestingly, the addition of some compounds, such as ACD176 and ACD055, even increased DHE fluorescence (● Fig. 3B); this observation was not investigated further as it is not directly related to Nox inhibition.

For further characterisation of the Nox4 inhibitors identified in this study, we assessed their ability to inhibit Nox4 activity *in vitro*. To this end, membranes were isolated from Nox4-expressing HEK cells (HEK-Nox4) and control HEK cells (HEK-pcDNA), and Nox4 content of the membrane fractions was verified by Western blot (● Fig. 4A). NADPH oxidase activity of these membrane fractions was assayed using AR fluorescence measurements, as described [24,27,34]. HEK-Nox4 derived membranes oxidised AR, and this activity was sensitive to DPI inhibition (● Fig. 4A), as expected. Addition of compounds ACD005, ACD047, ACD053, and ACD212 reduced Nox4 activity *in vitro*, and the inhibitory activity of ACD053 was similar to that of DPI, whereas VAS2870 showed rather weak activity in this assay (● Fig. 4B). Some of the compounds also inhibited the residual activity obtained with membranes from control (HEK-pcDNA) cells (● Fig. 4C), which most likely is not related to Nox4. For normalisation, the values obtained with control-transfected cells were subtracted from the respective values obtained with Nox4-expressing HEK cells. After normalisation, significant Nox4 inhibition was still obtained with compounds ACD053 and ACD047, whereas some compounds failed to inhibit Nox4 activity under these conditions (● Fig. 4D). Addition of ACD066 also significantly reduced AR fluorescence in this assay (● Fig. 4D); however, due to the proven capability of ACD066 to act as general ROS scavenger (see above, ● Fig. 3), this compound was excluded from further analysis. DPI fully inhibited Nox4 activity in intact cells (see above, ● Fig. 1), but after addition of DPI to Nox4-containing membranes, roughly 50% of the

chemiluminescence signal persisted, probably reflecting some background signals in the *in vitro* setting. Conceivably, the high background signal is due to the fact that much higher (roughly 10 fold) cell numbers are required for one measurement in the broken cell assay, compared to assays performed in intact cells; however, other reasons for the high background cannot be excluded.

Of note, the inhibitory potential of VAS2870 was strongly limited in these experiments, and ACD005 showed only an indistinct trend, relative to experiments in intact cells (● Fig. 4D); possible reasons for this are discussed below with the chemoinformatic analysis. The bioavailability and a compound's ability to penetrate into the cell is a big issue, and not all compounds that can inhibit an enzyme *in vitro* are expected to reach their protein target in intact cells. This is of particular importance for Nox4, since it is known that, unlike other members of the Nox family, a substantial proportion of Nox4 is found in intracellular membranes [7,48–50].

We also assessed the activity of selected compounds towards a recombinant fragment of Nox4, which retains the dehydrogenase reaction associated with Nox4 enzymatic activity. The protein fragment Nox4_{304–578} [51] was cloned in a baculoviral vector, expressed in SF9 insect cells as an MBP fusion protein and purified by affinity chromatography [51]. Upon incubation with NADPH, FAD, and NBT, significant dehydrogenase activity was detectable, which was abolished by boiling the recombinant protein prior to the assay (data not shown). The reaction could be triggered by addition of NADPH (Fig. 45) or, alternatively, by the addition of FAD in the presence of NADPH (data not shown), confirming the dependency on both cofactors. Surprisingly, addition of DPI did not lead to sustained inhibition of the diaphorase activity associated with Nox4DH domain; instead, DPI addition in this setting even increased signal intensity (data not shown). The reasons why DPI fails to act as an inhibitor in this setting remain to be clarified. Whereas compound ACD084 significantly inhibited the

Table 2 Selected compounds. Given in the table are the compound numbers as used in the paper in the first column. The second column gives the trivial name as known so far, the third column provides the order numbers from AnalytiCon Discovery. Column four shows the 2D chemical structure, and the last line provides the SMILE (Simplified Molecule Input Line Entry) string for each compound.

Compound Nr.	Name	Order Nr.	Structure	SMILES
ACD 005	–	NP-000520		<chem>O[C@@H]([C@H](OCC1C(C2=CC=C(O)C(OC)=C2)C3=CC(O)=C(OC)C=C3CC1CO)O4)[C@H]([C@@H]([C@H]4CO)O)O</chem>
ACD 018	Bakuchiol	NP-007515		<chem>OC1=CC=C(/C=C/[C@]([C/C=C(C)/C)(C)C=C)C=C1</chem>
ACD 042	Grindelic acid	NP-002752		<chem>CC1(CCC[C@]2(C)[C@]3(O[C@@](C)(CC(O)=O)CC3)C(C)=CCC12)C</chem>
ACD 047	–	NP-000261		<chem>CC(C(OC1C2=CC(OC)=C(O)C=C2)C3=CC=C(O)C(OC)=C3)C1C</chem>
ACD 053	–	NP-002766		<chem>OC(C(OC)=C1)=CC=C1/C=C/C(CC/C=C/C2=CC(O)=C(O)C=C2)=O</chem>
ACD 055	Dihydroyashabushiketol	NP-003489		<chem>O=C(C[C@@H](O)CCC1=CC=CC=C1)CCC2=CC=CC=C2</chem>
ACD 056	Gomisin J	NP-015225		<chem>CC1(C)CC2=C(C(OC)=C(OC)C(O)=C2)C3=C(C=C(O)C(OC)=C3OC)C1</chem>
ACD 063	–	NP-012941		<chem>OC(C=CC(CCCCC(CCC1=CC=C(O)C=C1)OC(C)=O)=C2)=C2O</chem>
ACD 066	–	NP-003620		<chem>O=C(CCCCC1=CC(O)=C(C=C1)O)CCC2=CC=C(O)C(O)=C2</chem>
ACD 067	–	NP-003332		<chem>O=C(CC(O)CCC1=CC=C(O)C(OC)=C1)CCC2=CC=CC=C2</chem>
ACD 069	–	NP-012926		<chem>CC(C(C(O1)C2=CC=C(O)C=C2)C)C1C3=CC(OC)=C(C=C3)OC</chem>
ACD 084	–	NP-003624		<chem>OC(C=CC(CCCCC(CCC1=CC=C(O)C(O)=C1)OC(C)=O)=C2)=C2O</chem>

continued

Table 2 Continued

Compound Nr.	Name	Order Nr.	Structure	SMILES
ACD 176	–	NP-003501		<chem>C[C@](C1CC2=O)(C3=C2C=C(O)C(O)=C3)CCCC1(C)C</chem>
ACD 212	–	NP-014837		<chem>CC(CC1=CC(OC(C2=CC=C(O)C=C2)=O)=C(O)C=C1)C(C)CC3=CC(O)=C(O)C=C3</chem>
VAS 2870	–	–		<chem>C1(N=C(SC2=NC=NC=C2N=NN3CC4=CC=CC=C4)O5)=C5C=CC=C1</chem>

dehydrogenase reaction catalysed by Nox4_{304–578}, compounds ACD005 and ACD063 displayed weak but significant inhibitory activity (**Fig. 4S**); in contrast, compounds ACD069 and VAS2870 were inactive in this assay (**Fig. 4S**). This may indicate that the mode of inhibition displayed by these inhibitors differs from the other compounds described in this communication. As was pointed out by Jaquet et al. [25], inhibition of Nox enzymes including Nox4 can occur at different stages, including depletion of substrate, interference with transcription or complex assembly, masking of the reaction product, etc. For compounds that inhibit Nox4 in intact cells but not in the Nox4DH assay, inhibition apparently involves Nox4 domains not present in the fragment. Accordingly, the mechanism of inhibition may differ between compounds described here and remains to be identified in all cases.

Plant-derived compounds described here can be assigned to several activity profiles: several compounds qualify as highly active Nox4 inhibitors, defined here by an EC₅₀ below 1 µM. This includes ACD005, ACD047, and ACD053, all of which also inhibited Nox2 activity. Two additional compounds identified here, ACD042 and ACD084, are of potential interest, because they did not inhibit the activity of Nox2 nor Nox5. They may be considered as specific Nox4 inhibitors, although their EC₅₀ is slightly higher than for the best Nox4 inhibitors described above. In quantitative terms, their activity is similar to the recently described Nox4 inhibitor VAS2870 [52], which in our hands displayed reasonable activity towards Nox4 and Nox2.

During screening as well as retrospectively, chemoinformatics tools were applied to guide selection of candidates and to characterize our best hits, respectively. In a shape-based alignment, we identified ACD018 and ACD067 as suitable query molecules to enrich hits within inactive compounds during the evaluation showing the highest area under the receiver operator curve value (**Fig. 5S**). A retrospective chemical characterisation was done by

calculation of selected chemical descriptors allowing an estimation of the drug-likeness for oral uptake according to Lipinski [53] (**Table 1S**). ACD005 was the only compound showing clear violation of Lipinski's Rule of Five with more than 10 acceptors, more than 5 donors, and a molecular weight above 500 Da. This candidate Nox4 inhibitor bears a sugar moiety and therefore is very hydrophilic which is reflected in a low logP (o/w). In contrast, high logP (o/w) values are estimated for the compounds ACD018 and ACD212. Although these logP (o/w) values are higher than the criteria of logP ≤ 5, those compounds are classified as drug-like according to Lipinski as one violation is accepted.

The retrospective analysis of the identified hits relies on molecular descriptors calculated *in silico*. Chemical characteristics within the hits presented here give insight in the chemical requirement for Nox4 inhibitors (**Table 2**). So, the occurrence of 6 diarylheptanoids (ACD053, ACD055, ACD063, ACD066, ACD067, and ACD084) and the presence of phenol substructures within 12 out of 14 presented compounds indicate that this substructure is beneficial for Nox4 inhibition. For ROCS, the high AUC values for ACD018 and ACD057 are caused by the presence of 4-substituted phenol substructures. This can be seen as a common characteristic within almost all actives except ACD042 and ACD005. In fact, ACD055, which is the only diarylheptanoid without hydroxylation at the aromatic rings, is less active (EC₅₀ = 1.88 µM) than the chemically related compound ACD067 with a phenol.

For ACD005, the low AUC value is caused by the sugar moiety unique within the test set. ACD005 was characterised by high activity towards Nox4 in the chemiluminescence assay in the assay with intact cells (**Table 1**) but a surprisingly low effect on the isolated membranes (**Fig. 4D**). We assume ACD005 to enter the intact cells although it violates the Lipinski's Rule of Five due to its sugar moiety. The glycosylation is a typical feature of natural products, and Lipinski's Rule of Five might not appropriately reflect the consequences on cellular uptake, as this rule of thumb

was developed from a statistical analysis of non-natural drug molecules [53]. Additionally, the glycosylic bond of ACD005 might be subject of enzymatic hydrolysis depending on the environment within the cells, and the aglycon might be the active form. This hypothesis could explain the reduced activity of ACD005 in the assays performed with isolated membranes and the isolated Nox4 dehydrogenase domain (Nox4_{304–578}) fragment. Besides ACD005, no Lipinski's Rule violations are observed. The two compounds, ACD042 and ACD084, discussed as specific Nox4 inhibitors, showed promising ADMET characteristics. Nox4 inhibitors have been described by others before. Thus, it was shown that extracts of *Piper sarmentosum* inhibit expression of Nox4 at the transcriptional level [26]. In an approach similar to the one chosen here [32], cell-based assays were used to characterise a set of synthetic compounds derived by medicinal chemistry for their ability to inhibit Nox4. Several hits with an IC₅₀ (roughly equivalent to the EC₅₀ described here) in the low micromolar range were reported, and Nox4 activity was entirely inhibited at a concentration of 10 µM for each of their best hits. The compounds described here have comparable activity in cell-based assays, but since they are derived from edible plants, we expect better compatibility with ADMET criteria. Based on a screening campaign of substance libraries with Nox4-containing membranes, K_i values in the two-digit nanomolar range were calculated for the best hits, including GKT137831 [25,33], which is currently being considered for clinical trials [28]. K_i values in general describe the kinetic interaction between enzyme and substrate, and their calculation is rather difficult in such an approach, given the fact that the molar concentration of Nox4 in isolated membranes is unknown. The comparison to the inhibitory activity of compounds described here is further complicated by the fact that EC₅₀ values for these compounds in cell-based assays are currently unknown. In a recent study with a cellular model of Nox4-dependent liver fibrosis, it was shown that inhibition of Nox4 activity in hepatocytes required GKT137831 at a concentration of 20 µM [28], suggesting that the most active compounds (ACD005, ACD047, ACD053, ACD66, ACD067; see **Table 1**) identified here are at least as active as GKT137831, which is considered the best available pharmacological Nox4 inhibitor to date [31].

Acknowledgements

This work was supported by a grant from the Austrian Science Funds (FWF; P23742) and the European Union (integrated project MiMage). We thank V. Jaquet and K.-H. Krause for help with the broken cell assay and technical advice, B. Lener for initial help with the HEK-Nox4-based screening system, and J.E. Fuchs for technical and scientific support for *in silico* characterisation. We are grateful to H. Stuppner, the NCI, and Vasopharm AG for the kind gift of several compounds.

Conflict of Interest

The authors declare no conflict of interest.

Affiliations

- ¹ Institute for Biomedical Aging Research (IBA), Austrian Academy of Sciences, Innsbruck, Austria
- ² Institute of General, Inorganic and Theoretical Chemistry and Center for Molecular Biosciences Innsbruck, University of Innsbruck, Innsbruck, Austria
- ³ Tyrolean Cancer Research Institute at the Medical University of Innsbruck, Innsbruck, Austria
- ⁴ AnalytiCon Discovery GmbH, Potsdam, Germany

References

- Perry BN, Govindarajan B, Bhandarkar SS, Knaus UG, Valo M, Sturk C, Carrillo CO, Sohn A, Cerimele F, Dumont D, Losken A, Williams J, Brown LF, Tan XL, Ioffe E, Yancopoulos GD, Arbiser JL. Pharmacologic blockade of angiopoietin-2 is efficacious against model hemangiomas in mice. *J Invest Dermatol* 2006; 126: 2316–2322
- Lambeth JD. Nox enzymes and the biology of reactive oxygen. *Nat Rev Immunol* 2004; 4: 181–189
- Lambeth JD, Krause KH, Clark RA. NOX enzymes as novel targets for drug development. *Sem Immunopathol* 2008; 30: 339–363
- Geiszt M. NADPH oxidases: New kids on the block. *Cardiovasc Res* 2006; 71: 289–299
- Martyn KD, Frederick LM, von Loehneysen K, Dinanier MC, Knaus UG. Functional analysis of Nox4 reveals unique characteristics compared to other NADPH oxidases. *Cell Signal* 2006; 18: 69–82
- Luo Z, Chen Y, Chen S, Welch WJ, Andresen BT, Jose PA, Wilcox CS. Comparison of inhibitors of superoxide generation in vascular smooth muscle cells. *Br J Pharmacol* 2009; 157: 935–943
- Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: Physiology and pathophysiology. *Physiol Rev* 2007; 87: 245–313
- Royerpokora B, Kunkel LM, Monaco AP, Goff SC, Newburger PE, Baehner RL, Cole FS, Curnutte JT, Orkin SH. Cloning the gene for an inherited human disorder – chronic granulomatous-disease – on the basis of its chromosomal location. *Nature* 1986; 322: 32–38
- Krause KH. Tissue distribution and putative physiological function of NOX family NADPH oxidases. *Jpn J Infect Dis* 2004; 57: S28–29
- Chen K, Craig SE, Keaney Jr. JF. Downstream targets and intracellular compartmentalization in Nox signaling. *Antiox Redox Signal* 2009; 11: 2467–2480
- Geiszt M, Kopp JB, Varnai P, Leto TL. Identification of Renox, an NAD(P)H oxidase in kidney. *Proc Natl Acad Sci USA* 2000; 97: 8010–8014
- Ago T, Kitazono T, Ooboshi H, Iyama T, Han YH, Takada J, Wakisaka M, Ibayashi S, Utsumi H, Iida M. Nox4 as the major catalytic component of an endothelial NAD(P)H oxidase. *Circulation* 2004; 109: 227–233
- Yang S, Madyastha P, Bingel S, Ries W, Key L. A new superoxide-generating oxidase in murine osteoclasts. *J Biol Chem* 2001; 276: 5452–5458
- Ellmark SH, Dusting GJ, Fui MN, Guzzo-Pernell N, Drummond GR. The contribution of Nox4 to NADPH oxidase activity in mouse vascular smooth muscle. *Cardiovasc Res* 2005; 65: 495–504
- Cucoranu I, Clempus R, Dikalova A, Phelan PJ, Ariyan S, Dikalov S, Sorescu D. NAD(P)H oxidase 4 mediates transforming growth factor-beta1-induced differentiation of cardiac fibroblasts into myofibroblasts. *Circ Res* 2005; 97: 900–907
- Gorin Y, Ricono JM, Kim NH, Bhandari B, Choudhury GG, Abboud HE. Nox4 mediates angiotensin II-induced activation of Akt/protein kinase B in mesangial cells. *Am J Physiol Renal Physiol* 2003; 285: F219–229
- Mahadev K, Motoshima H, Wu X, Ruddy JM, Arnold RS, Cheng G, Lambeth JD, Goldstein BJ. The NAD(P)H oxidase homolog Nox4 modulates insulin-stimulated generation of H₂O₂ and plays an integral role in insulin signal transduction. *Mol Cell Biol* 2004; 24: 1844–1854
- Li J, Stouffs M, Serrander L, Banfi B, Bettoli E, Charnay Y, Steger K, Krause KH, Jacani ME. The NADPH oxidase NOX4 drives cardiac differentiation: Role in regulating cardiac transcription factors and MAP kinase activation. *Mol Biol Cell* 2006; 17: 3978–3988
- Uchizono Y, Takeya R, Iwase M, Sasaki N, Oku M, Imoto H, Iida M, Sumimoto H. Expression of isoforms of NADPH oxidase components in rat pancreatic islets. *Life Sci* 2006; 80: 133–139
- Serrander L, Cartier L, Bedard K, Banfi B, Lardy B, Plastre O, Sienkiewicz A, Forro L, Schlegel W, Krause KH. NOX4 activity is determined by mRNA levels and reveals a unique pattern of ROS generation. *Biochem J* 2007; 406: 105–114
- Sampson N, Koziel R, Zenzmaier C, Bubendorf L, Plas E, Jansen-Durr P, Berger P. ROS signaling by NOX4 drives fibroblast-to-myofibroblast differentiation in the diseased prostatic stroma. *Mol Endocrinol* 2011; 25: 503–515

- 22 Schroder K, Zhang M, Benkhoff S, Mieth A, Pliquett R, Kosowski J, Kruse C, Luedike P, Michaelis UR, Weissmann N, Dimmeler S, Shah AM, Brandes RP. Nox4 is a protective reactive oxygen species generating vascular NADPH oxidase. *Circ Res* 2012; 110: 1217–1225
- 23 Li Y, Mouche S, Sajic T, Veyrat-Durebex C, Supale R, Pierroz D, Ferrari S, Negro F, Hasler U, Feraile E, Moll S, Meda P, Deffert C, Montet X, Krause KH, Szanto I. Deficiency in the NADPH oxidase 4 predisposes towards diet-induced obesity. *Int J Obes (Lond)* 2012; 36: 1503–1513
- 24 Laleu B, Gaggini F, Orchard M, Fioraso-Cartier L, Cagnon L, Houngninou-Molango S, Gradia A, Duboux G, Merlot C, Heitz F, Szyndralewicz C, Page P. First in class, potent, and orally bioavailable NADPH oxidase isoform 4 (Nox4) inhibitors for the treatment of idiopathic pulmonary fibrosis. *J Med Chem* 2010; 53: 7715–7730
- 25 Jaquet V, Scapozza L, Clark RA, Krause K-H, Lambeth JD. Small-molecule NOX inhibitors: ROS-generating NADPH oxidases as therapeutic targets. *Antiox Redox Signal* 2009; 11: 2535–2552
- 26 Ugusman A, Zakaria Z, Hui CK, Nordin NA. *Piper sarmentosum* inhibits ICAM-1 and Nox4 gene expression in oxidative stress-induced human umbilical vein endothelial cells. *BMC Complement Alternat Med* 2011; 11: 31
- 27 Jaquet V, Marcoux J, Forest E, Leidal KG, McCormick S, Westermaier Y, Perozzo R, Plastre O, Fioraso-Cartier L, Diebold B, Scapozza L, Nauseef WM, Fieschi F, Krause KH, Bedard K. NADPH oxidase (NOX) isoforms are inhibited by celastrol with a dual mode of action. *Br J Pharmacol* 2011; 164: 507–520
- 28 Jiang JX, Chen X, Serizawa N, Szyndralewicz C, Page P, Schroder K, Brandes RP, Devaraj S, Torok NJ. Liver fibrosis and hepatocyte apoptosis are attenuated by GKT137831, a novel NOX4/NOX1 inhibitor *in vivo*. *Free Radic Biol Med* 2012; 53: 289–296
- 29 Doussiere J, Gaillard J, Vignais PV. The heme component of the neutrophil NADPH oxidase complex is a target for arylidonium compounds. *Biochemistry* 1999; 38: 3694–3703
- 30 O'Donnell BV, Tew DG, Jones OT, England PJ. Studies on the inhibitory mechanism of iodonium compounds with special reference to neutrophil NADPH oxidase. *Biochem J* 1993; 290: 41–49
- 31 Kim JA, Neupane GP, Lee ES, Jeong BS, Park BC, Thapa P. NADPH oxidase inhibitors: a patent review. *Expert Opin Ther Pat* 2011; 21: 1147–1158
- 32 Borbely G, Szabadkai I, Horvath Z, Marko P, Varga Z, Breza N, Baska F, Vantus T, Huszar M, Geiszt M, Hunyady L, Buday L, Orfi L, Keri G. Small-molecule inhibitors of NADPH oxidase 4. *J Med Chem* 2010; 53: 6758–6762
- 33 Wind S, Beuerlein K, Eucker T, Muller H, Scheurer P, Armitage ME, Ho H, Schmidt H, Wingler K. Comparative pharmacology of chemically distinct NADPH oxidase inhibitors. *Br J Pharmacol* 2010; 161: 885–898
- 34 Gaggini F, Laleu B, Orchard M, Fioraso-Cartier L, Cagnon L, Houngninou-Molango S, Gradia A, Duboux G, Merlot C, Heitz F, Szyndralewicz C, Page P. Design, synthesis and biological activity of original pyrazolo-pyridodiazepine, -pyrazine and -oxazine dione derivatives as novel dual Nox4/Nox1 inhibitors. *Bioorg Med Chem* 2011; 19: 6989–6999
- 35 Borbely G, Huszar M, Varga A, Futosi K, Mocsai A, Orfi L, Idei M, Mandl J, Keri G, Vantus T. Optimization of important early ADME(T) parameters of NADPH oxidase-4 inhibitor molecules. *Med Chem* 2012; 8: 174–181
- 36 Lener B, Koziel R, Pircher H, Hutter E, Greussing R, Herndler-Brandstetter D, Hermann M, Unterluggauer H, Jansen-Durr P. The NADPH oxidase Nox4 restricts the replicative lifespan of human endothelial cells. *Biochem J* 2009; 423: 363–374
- 37 Stockl P, Hutter E, Zwerschke W, Jansen-Durr P. Sustained inhibition of oxidative phosphorylation impairs cell proliferation and induces premature senescence in human fibroblasts. *Exp Gerontol* 2006; 41: 674–682
- 38 Caldwell SE, McCall CE, Hendricks CL, Leone PA, Bass DA, McPhail LC. Coregulation of NADPH oxidase activation and phosphorylation of a 48-kD protein(s) by a cytosolic factor defective in autosomal recessive chronic granulomatous disease. *J Clin Invest* 1988; 81: 1485–1496
- 39 Grant JA, Gallardo M, Pickup B. A fast method of molecular shape comparison: A simple application of a Gaussian description of molecular shape. *J Computat Chem* 1996; 17: 1653–1666
- 40 Mills JE, Dean PM. Three-dimensional hydrogen-bond geometry and probability information from a crystal survey. *J Comput Aided Mol Des* 1996; 10: 607–622
- 41 Stielow C, Catar RA, Muller G, Wingler K, Scheurer P, Schmidt HH, Morawietz H. Novel Nox inhibitor of oxLDL-induced reactive oxygen species formation in human endothelial cells. *Biochem Biophys Res Commun* 2006; 344: 200–205
- 42 ten Freyhaus H, Huntgeburth M, Wingler K, Schnitker J, Baumer AT, Vantler M, Bekhte MM, Wartenberg M, Sauer H, Rosenkranz S. Novel Nox inhibitor VAS2870 attenuates PDGF-dependent smooth muscle cell chemotaxis, but not proliferation. *Cardiovasc Res* 2006; 71: 331–341
- 43 Stefanska J, Pawliczak R. Apocynin: molecular aptitudes. *Mediators Inflamm* 2008; DOI: 10.1155/2008/106507
- 44 Liou KT, Shen YC, Chen CF, Tsao CM, Tsai SK. The anti-inflammatory effect of honokiol on neutrophils: mechanisms in the inhibition of reactive oxygen species production. *Eur J Pharmacol* 2003; 475: 19–27
- 45 Banfi B, Tirone F, Durussel I, Knisz J, Moskwa P, Molnar GZ, Krause KH, Cox JA. Mechanism of Ca²⁺ activation of the NADPH oxidase 5 (NOX5). *J Biol Chem* 2004; 279: 18583–18591
- 46 Foti MC, Amorati R. Non-phenolic radical-trapping antioxidants. *J Pharm Pharmacol* 2009; 61: 1435–1448
- 47 Fato R, Bergamini C, Bortolus M, Maniero AL, Leoni S, Ohnishi T, Lenaz G. Differential effects of mitochondrial complex I inhibitors on production of reactive oxygen species. *Biochim Biophys Acta* 2009; 1787: 384–392
- 48 Helmcke I, Heumuller S, Tikkanen R, Schroder K, Brandes RP. Identification of structural elements in Nox1 and Nox4 controlling localization and activity. *Antiox Redox Signal* 2009; 11: 1279–1287
- 49 Lyle AN, Deshpande NN, Taniyama Y, Seidel-Rogol B, Pounkova L, Du P, Papaharalambus C, Lassegue B, Griendling KK. Poldip2, a novel regulator of Nox4 and cytoskeletal integrity in vascular smooth muscle cells. *Circ Res* 2009; 105: 249–259
- 50 Zhang L, Nguyen MV, Lardy B, Jesaitis AJ, Grichine A, Rousset F, Talbot M, Paclet MH, Qian G, Morel F. New insight into the Nox4 subcellular localization in HEK293 cells: first monoclonal antibodies against Nox4. *Biochimie* 2011; 93: 457–468
- 51 Nisimoto Y, Jackson HM, Ogawa H, Kawahara T, Lambeth JD. Constitutive NADPH-dependent electron transferase activity of the Nox4 dehydrogenase domain. *Biochemistry* 2010; 49: 2433–2442
- 52 Kleinschnitz C, Grund H, Wingler K, Armitage ME, Jones E, Mittal M, Barit D, Schwarz T, Geis C, Kraft P, Barthel K, Schuhmann MK, Herrmann AM, Meuth SG, Stoll G, Meurer S, Schrewe A, Becker L, Gailus-Durner V, Fuchs H, Klopstock T, de Angelis MH, Jandeleit-Dahm K, Shah AM, Weissmann N, Schmidt HH. Post-stroke inhibition of induced NADPH oxidase type 4 prevents oxidative stress and neurodegeneration. *PLoS Biol* 2010; 8: e1000479
- 53 Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv Drug Deliv Rev* 2001; 46: 3–26