T. WAKAMATSU, K. NAGAO, H. OHMIYA,* M. SAWAMURA* (HOKKAIDO UNIVERSITY, SAPPORO, JAPAN)

Synthesis of Trisubstituted Alkenylstannanes through Copper-Catalyzed Three-Component Coupling of Alkylboranes, Alkynoates and Tributyltin Methoxide
Angew. Chem. Int. Ed. **2013**, 52, 11620–11623.

Synthesis of Trisubstituted Alkenylstannanes Starting from Alkynoates

R¹ = Ph, *n*-Bu, *n*-Pent, (CH₂)₃phthalimide, (CH₂)₄OTHP, C(Me)₂CH₂CO₂Me, 4-BrC₆H₄, (CH₂)₃OTIPS
R² = Ph, 4-MeOC₆H₄, 4-MeO₂CC₆H₄, 2-MeC₆H₄, Me, CH₂OTHP, CH₂OBn, 4-FC₆H₄, 2-thienyl derivative

Selected examples:

Significance: The authors report a highly regio-selective copper-catalyzed synthesis of trisubstituted alkenylstannanes. Through a three-component coupling of alkylboranes, alkynoates and tributyltin methoxide, these trisubstituted alkenylstannanes are obtained in good yields and with high *syn* selectivity. The appropriate alkylboranes are easily accessible by hydroboration of the corresponding alkenes with the 9-borabicyclo[3.3.1]nonane (9-BBN-H) dimer.

Comment: Standard methods for the synthesis of alkenylstannanes described by Shirakawa and Hiyama include the palladium- or nickel-catalyzed carbostannylation of internal alkynes with organostannanes which are somewhat difficult to prepare.

SYNFACTS Contributors: Paul Knochel, Nadja M. Barl
Synfacts 2014, 10(1), 0084 Published online: 13.12.2013
DOI: 10.1055/s-0033-1340371; Reg-No.: P16113SF

2014 © THIEME STUTTGART • NEW YORK