H. TSUJI,* G. CANTAGREL, Y. UEDA, T. CHEN, L.-J. WAN,* E. NAKAMURA* (THE UNIVERSITY OF TOKYO AND JST-PRESTO, SAITAMA, JAPAN; BEIJING NATIONAL LABORATORY FOR MOLECULAR SCIENCES AND INSTITUTE OF CHEMISTRY, BEIJING, P. R. OF CHINA)

Synthesis of Benzotrifuran and Benzotripyrrole Derivatives and Molecular Orientations on the Surface and in the Solid State

Chem. Asian J. **2013**, *8*, 2377–2382.

Facile Access to Heteroaromatic Triphenylene Analogues

Significance: Planar disc-shaped molecules have attractive properties for applications in materials science, particularly when they exhibit three-fold symmetry. In the present work, the authors demonstrate a versatile approach to the synthesis of benzotrifurans and benzotripyrroles. The C_{3h}-symmetric target molecules are obtained in a triple cyclization with excellent yields over a wide range of substrates (75–97%). The authors also study the 2-D molecular arrangement of the phenyl-substituted benzotrifuran via scanning tunneling microscopy and compare it to its 3-D arrangement in the crystalline state.

Comment: The starting 1,3,5-triethynylbenzenes are prepared by Sonogashira coupling with 1,3,5-trifluoro-2,4,6-triiodobenzene. Reaction with cesium hydroxide in *N,N*-dimethylacetamide (DMA) or with a sodium amide in an amine solvent induces the triple cyclization. For alkynes with *tert*-butyl substitution, the reaction hardly proceeds beyond the second ring closure. Crystal structures were obtained for four benzotrifurans and one benzotripyrrol, and the authors analyze their emission spectra in the solid state and in solution.

SYNFACTS Contributors: Timothy M. Swager, Georgios Markopoulos

Synfacts 2014, 10(1), 0036 Published online: 13.12.2013

DOI: 10.1055/s-0033-1340392; Reg-No.: S14813SF

2014 © THIEME STUTTGART • NEW YORK