Y. S. YOU, T. W. KIM, S. H. KANG* (KOREA ADVANCED INSTITUTE OF SCIENCE AND TECHNOLOGY, DAEJEON, KOREA)

Asymmetric Formation of *tert*-Alkylamines from Serinols by a Dual Function Catalyst

Chem. Commun. **2013**, 49, 9669–9671.

Consecutive Intramolecular Desymmetrization and Kinetic Resolution

Selected examples:

Proposed intermediates in desymmetrization and kinetic resolution:

Significance: This paper describes the consecutive intramolecular desymmetrization and kinetic resolution of 2-substituted *N*-phenoxyserinols by using bisoxazoline- CuCl_2 . The reaction products are obtained in good yield and excellent enantioselectivities (94–99% ee).

Comment: The authors developed a unique asymmetric catalysis system using a single chiral Lewis acid catalyst, which steers two consecutive asymmetric reactions of intramolecular desymmetrization and kinetic resolution. The two successive chemical conversions became unusually enantioselective, resulting in the production of the oxazolidinone benzoates with excellent enantioselectivities, which signifies a powerful and synergistic dual function catalyst effect by a chiral catalyst.

SYNFACTS Contributors: Hisashi Yamamoto, Atsuto Izumiseki
Synfacts 2014, 10(1), 0048 Published online: 13.12.2013
DOI: 10.1055/s-0033-1340432; **Reg-No.:** H15313SF

2014 © THIEME STUTTGART • NEW YORK