W. LI, X. LIU, F. TAN, X. HAO, J. ZHENG, L. LIN, X. FENG* (SICHUAN UNIVERSITY, CHENGDU, P. R. OF CHINA)

Catalytic Asymmetric Homologation of α -Ketoesters with α -Diazoesters: Synthesis of Succinate Derivatives with Chiral Quaternary Centers

Angew. Chem. Int. Ed. **2013**, 52, 10883–10886.

Enantioselective Homologation of α -Keto Esters with α -Diazo Esters

Significance: The Lewis acid catalyzed homologation of carbonyl compounds with diazo compounds can realize synthetically useful carbon chain extension. The authors achieve the asymmetric homologation of acyclic α -keto esters with α -diazo esters by using chiral N,N' -dioxide-ylttrium(III) complexes. Both aryl- and alkyl-substituted α -keto esters are applicable, providing the corresponding succinate derivatives in good yields and enantioselectivities.

SYNFACTS Contributors: Hisashi Yamamoto, Yusuke Ano
Synfacts 2014, 10(1), 0055 Published online: 13.12.2013

DOI: 10.1055/s-0033-1340454; **Reg-No.:** H16113SF

Comment: The use of bulky adamantyl α -diazo esters can suppress the formation of undesired by-products. Steric hindrance on the 2,6-positions of the phenyl ring in the ligand is also essential to improve both enantioselectivity and reactivity. The attack of α -diazo ester occurs from re -face of the coordinating α -keto ester preferably due to the obstruction of si -face by the aryl group in the N,N' -dioxide ligand.

Category

Metal-Catalyzed
Asymmetric
Synthesis and
Stereoselective
Reactions

Key words

diazo esters

α -keto esters

yttrium