X. LI, Y. ZHU, J. SHAO, B. WANG, S. ZHANG, Y. SHAO, X. JIN, X. YAO, R. FANG, X. SHAO*
(LANZHOU UNIVERSITY, P. R. OF CHINA)

Non-Pyrolytic, Large-Scale Synthesis of Trichalcogenasumanene: A Two-Step Approach
Angew. Chem. Int. Ed. **2014**, *53*, 535–538.

‘Chalc’ing up a New Sumanene

Significance: Sumanenes are nanocarbon buckybowls with a six-membered benzene ring at their center. The authors report a new synthesis of sulfur- or selenium-substituted sumanenes from a triphenylene that avoids the very high temperature FVP methods (1000 °C) previously used. The new method employs much milder conditions and achieves much higher yields than the only other reported synthesis of trithiosumanene (K. Imamura et al. *Chem. Commun.* **1999**, 1859), and also provides access to the previously unknown triselenasumanene.

Comment: The reported synthesis circumvents the challenge of simultaneous insertion of chalcogens into the bay positions of triphenylene by first accessing 1,2-dichalcogenin rings. These six-membered rings presumably have less ring strain than the monochalcogen rings and can be converted into the desired trichalcogenosumanenes by using copper powder in a solid-state reaction. This reaction can be performed on multi-gram scale.

SYNFACTS Contributors: Timothy M. Swager, Graham T. Sazama
Synfacts 2014, 10(3), 0255 Published online: 17.02.2014
DOI: 10.1055/s-0033-1340784; Reg-No.: S01214SF