SYNLETT Spotlight 466

Allyl Alkyl Carbonates

Compiled by Martin Riemer

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Martin Riemer was born in 1985 in Templin, Germany. He studied chemistry at the University of Potsdam and received his diploma in 2011. Currently, he is working towards his Ph.D. under the supervision of Professor Dr. Bernd Schmidt. His current research interests focus on applications of palladium-catalysed reactions for the synthesis of (poly)phenolic natural products.

Department of Chemistry, University Potsdam, Karl-Liebknecht-Str. 24–25, 14476 Golm, Germany
E-mail: martin.riemer@uni-potsdam.de

Dedicated to my family

Introduction

Allyl methyl carbonate is the simplest allyl alkyl carbonate. It was first synthesised by Hermann Otto Laurenz Fischer in 1929.¹ In general, a synthesis of these carbonates is possible in high yields starting from allyl alcohols, which can be converted with dialkyl dicarbonates² or alkyl chloroformates³ under basic conditions (e.g., BuLi, pyridine) into the corresponding allyl alkyl carbonates. Allyl carbonates are highly versatile reagents, and they can be used both for nucleo- and electrophilic reactions.

Abstracts

(A) Allylic Alkylation of Nucleophiles; C–C Bond Formation

Tsuji⁴ and Trost⁵ pioneered the palladium-catalysed allylation of nucleophiles. The allylic alkylation is a versatile method to construct C–C bonds, especially products with bulky quaternary carbon centres.^{6,7} The allyl carbonate and palladium form an η^3 -allylpalladium complex, which is attacked by a nucleophile. In general, C–H acidic compounds are used, but nucleophiles like diphenylmethane are also suitable.⁸

(B) Allylic Alkylation of Nucleophiles; C–Het Bond Formation

Various heteroatoms – with aliphatic as well as aromatic substituents – can be allylated by allyl alkyl carbonates^{9–11} using catalysis by palladium or iron complexes.¹² While the reaction of 1,1-dimethylallyl bromide with phenol leads to the unexpected *n*-product due to an S_N' reaction,¹³ the analogue carbonate leads to the desired *iso* product.¹⁴

(C) Asymmetric Allylation of Nucleophiles; C–C Bond Formation

The Trost asymmetric allylic alkylation, often referred to as AAA, is the enantioselective version of the Tsuji–Trost reaction. The AAA is catalysed by palladium or molybdenum. The enantioselectivity can be introduced by a chiral ligand, for example by a tetradentate Trost ligand.¹⁵ Furthermore, branched asymmetric allylation products can be generated by iridium catalysis. The enantioselectivity can be introduced as described above.¹⁶

(D) Reductive Allylation of Alkyl Halides

The direct allylation of alkyl halides results in a $C(sp^3)-C(sp^3)$ coupling. This catalysed reaction proceeds via an allyl alkyl cobalt intermediate. Manganese acts as reducing agent for the allyl copper complex.¹⁷

(E) Barbier-Type Allylation

Allyl ethyl carbonate can be used for the allylation of aldehydes and ketones in good yield. Furthermore, crotylation, prenylation, and intramolecular allylation are also possible with the corresponding carbonate. The Barbier-type allylation is mediated by a bimetallic system of titanium/palladium and highly accelerated by manganese dust. Initially, palladium undergoes an oxidative addition with allyl carbonate. Single electron transfer (SET) of the η^3 -allylpalladium complex forms a palladium(I) intermediate. This species fragments further to an allyl radical, which can form the nucleophilic η^3 -allyltitanocene(IV) complex.¹⁸

(F) Alder-ene Reaction

1,4-Dienes can be formed by the Alder-ene reaction of allyl carbonates and alkynes. The *E/Z*-selectivity could be increased by the use of a permethylated cyclopentadienyl ruthenium complex.¹⁹

(G) Allylation of Styrenes via a Heck-Type Reaction

The iridium-catalysed reaction of 2-vinylanilines and allyl carbonates leads to *Z,E*-dienes. This method is a *cis*-selective supplement to the Heck reaction, which affords the *trans* products. The authors discuss an amine-assisted iridium-catalysed vinyl C–H bond activation to form the reactive intermediate.²⁰

(H) Direct C–H Allylation of Arenes

The allylation of arenes is catalysed by a permethylated cyclopentadienyl ruthenium complex. The reaction proceeds via C–H activation and is directed by *N,N*-diisopropylacetamide.²¹

References

- (1) Fischer, H. O. L.; Feldmann, L. *Chem. Ber.* **1929**, *62*, 858.
- (2) Lemke, M.-K.; Schwab, P.; Fischer, P.; Tischer, S.; Witt, M.; Noehringer, L.; Rogachev, V.; Jäger, A.; Kataeva, O.; Fröhlich, R.; Metz, P. *Angew. Chem. Int. Ed.* **2013**, *52*, 11651.
- (3) Trost, B. M.; Miller, J. R.; Hoffman, C. M. Jr. *J. Am. Chem. Soc.* **2011**, *133*, 8165.
- (4) Tsuji, J.; Takahashi, H.; Morikawa, M. *Tetrahedron Lett.* **1965**, *6*, 4387.
- (5) Trost, B. M.; Fullerton, T. J. *J. Am. Chem. Soc.* **1973**, *95*, 292.
- (6) Dickschat, A. T.; Behrends, F.; Surmiak, S.; Weiß, M.; Eckert, H.; Studer, A. *Chem. Commun.* **2013**, *49*, 2195.
- (7) Maki, K.; Kanai, M.; Shibasaki, M. *Tetrahedron* **2007**, *63*, 4250.
- (8) Sha, S.-C.; Zhang, J.; Carroll, P. J.; Walsh, P. J. *J. Am. Chem. Soc.* **2013**, *135*, 17602.
- (9) Schmidt, B.; Nave, S. *Adv. Synth. Catal.* **2006**, *348*, 531.
- (10) Dieskau, A. P.; Plietker, B. *Org. Lett.* **2011**, *13*, 5544.
- (11) Deska, J.; Kazmaier, U. *Chem. Eur. J.* **2007**, *13*, 6204.
- (12) Plietker, B. *Angew. Chem. Int. Ed.* **2006**, *45*, 6053.
- (13) Patent: US4515801 A1, Bayer Aktiengesellschaft, **1985**.
- (14) Beare, K. D.; McErlean, C. S. P. *Tetrahedron Lett.* **2013**, *54*, 1056.
- (15) Trost, B. M.; Zhang, Y. *Chem. Eur. J.* **2011**, *17*, 2916.
- (16) Gärtner, M.; Jäkel, M.; Achatz, M.; Sonnenschein, C.; Tverskoy, O.; Helmchen, G. *Org. Lett.* **2011**, *13*, 281.
- (17) Qian, X.; Auffrant, A.; Felouat, A.; Gosmini, C. *Angew. Chem. Int. Ed.* **2011**, *50*, 10402.
- (18) Millán, A.; Campaña, A. G.; Bazdi, B.; Miguel, D.; Álvarez de Cienfuegos, L.; Echavarren, A. M.; Cuerva, J. M. *Chem. Eur. J.* **2011**, *17*, 3985.
- (19) Trost, B. M.; Martos-Redruejo, A. *Org. Lett.* **2009**, *13*, 1071.
- (20) He, H.; Liu, W. B.; Dai, L. X.; You, S. L. *J. Am. Chem. Soc.* **2009**, *131*, 8346.
- (21) Wang, H.; Schröder, N.; Glorius, F. *Angew. Chem. Int. Ed.* **2013**, *52*, 5386.