J. TURCONI,* F. GRIOLET, R. GUEVEL, G. ODDON, R. VILLA, A. GEATTI, M. HVALA, K. ROSSEN, R. GÖLLER, A. BURGARD* (SANOFI, SISTERON, MARCY L'ETOILE AND NEUVILLE SUR SAÔNE, FRANCE; SANOFI-AVENTIS, GARESSIO, ITALY; SANOFI-AVENTIS, FRANKFURT AM MAIN, GERMANY)

Semisynthetic Artemisinin, the Chemical Path to Industrial Production

Org. Process Res. Dev. 2014, 18, 417–422.

Industrial Production of Artemisinin

Category

Synthesis of Natural Products and Potential Drugs

Key words

artemisinin

artemisinic acid

diastereoselective hydrogenation

photooxidation

SYNFACT
of the month

Significance: Artemisinin is an antimalarial agent extracted from the leaves of *Artemisia annua*. The quest for a more reliable and economic supply of the drug resulted in the large-scale synthesis depicted, which is based on two key steps: (1) the diastereoselective hydrogenation of artemisinic acid (**A**) and (2) the photooxidation of dihydroartemisinic acid derivative **D**.

SYNFACTS Contributors: Philip Kocienski
Synfacts 2014, 10(6), 0555 Published online: 16.05.2014
DOI: 10.1055/s-0033-1341264; Reg-No.: K02214SF

Comment: A major breakthrough was the discovery of a highly efficient biosynthetic route to artemisinic acid (**A**) in engineered *Saccharomyces cerevisiae* (C. J. Paddon et al. *Nature* 2013, 496, 528). This commercial synthesis delivers 365 kg batches of artemisinin starting from 600 kg of **A** (55% overall) and is expected to produce 60 tons per annum.