Retrorectal cystic hamartoma: this cyst has a “tail” to tell

A 50-year-old man presented with a 6-week history of lumbosacral back pain and altered bowel habit without hematochezia. Abdominopelvic computed tomography (CT) revealed a 5-cm rectal mass with obliteration of the fat plane between the mass and the tip of the coccyx that was interpreted as representing a neoplasm (► **Fig. 1 a**). Following a normal digital rectal exam, colonoscopy revealed a subtle posterior mid-rectal wall indentation, suspected to represent either an intramural process or extrinsic compression. The overlying mucosa appeared normal (► **Fig. 1 b**). A curvilinear endoscopic ultrasound examination excluded a sub-epithelial lesion, instead revealing a 3.4 × 3.3 cm multiloculated cystic mass extrin-

sic but adjacent to the rectal wall with no evidence of lymphadenopathy. This lesion was located distal and posterior to the inferior border of the prostate gland and comprised approximately 15 loculi. Each locule was thin-walled in the absence of septations. Many of the loculi contained intracystic echogenic material (► **Video 1**). Surgical resection of the retrorectal cystic mass mandated posterior dissection of the coccyx laterally. Surgical pathology confirmed a multiloculated presacral cystic mass measuring 6.0 × 4.3 × 3.2 cm (► **Fig. 1 c**). The cystic loculi were lined by squamous epithelium, transitional type epithelium, and columnar epithelium with focal goblet cells and no evidence of malignancy (► **Fig. 1 d**). The

findings were diagnostic of cystic hamartoma (tailgut cyst).

Until adverse event data for lower gastrointestinal endoscopic ultrasound (EUS) become available, the current recommendation is to avoid fine-needle aspiration (FNA) of perirectal cysts due to concerns regarding abscess formation even with prophylactic antibiotic use, as highlighted in one patient who subsequently required percutaneous drainage [1]. Complete intact surgical excision is advised to avoid the potential risk of needle-tract seeding, infection, and fistula formation [2].

Endoscopy_UCTN_Code_CCL_1AF_2AH

Competing interests: None

Video 1

A multiloculated extramural cystic structure with occasional intracystic locular echogenic material suggestive of mucin or debris.

Fig. 1 a Preoperative computed tomography (CT) scan of the pelvis in a 50-year-old man with lumbosacral back pain and altered bowel habit without hematochezia. b Retroflexed endoscopic view of the presented lesion preoperatively. c, d The surgical and histopathological examination findings.

F. C. Gleeson, M. J. Levy

Division of Gastroenterology
and Hepatology, Mayo Clinic College
of Medicine, Rochester, Minnesota,
United States of America

References

- 1 *Mohamadnejad M, Al-Haddad MA, Sherman S et al.* Utility of EUS-guided biopsy of extramural pelvic masses. *Gastrointest Endosc* 2012; 75: 146–151
- 2 *Mathis KL, Dozois EJ, Grewal MS et al.* Malignant risk and surgical outcomes of presacral tailgut cysts. *Br J Surg* 2010; 97: 575–579

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1344130>
Endoscopy 2013; 45: E191–E192
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author**F. C. Gleeson**

Division of Gastroenterology and Hepatology
Mayo Clinic College of Medicine
200 First Street SW
Rochester
MN 55905
USA
Fax: +507-266-3939
gleeson.ferga@mayo.edu