Endoscopic treatment of rectal pocket syndrome

Fig. 1 Rectal bulge with central orifice.

A 70-year-old woman was admitted with rectal pain and soiling 3 years after a Longo's stapled hemorrhoidopexy [1]. Physical examination and laboratory tests were negative. Colonoscopy showed a 30-mm bulge with a 3-mm orifice in the lower rectum. Stools could be observed through the orifice (● **Fig. 1**). A pelvic magnetic resonance imaging scan highlighted a stool-laden diverticulum of the mesorectal region, 35×20×25 mm in size (● **Fig. 2**). An endoscopic ultrasound color-Doppler examination showed poor vascularization of the septum between the diverticulum and the rectal lumen. Endoscopic incision of the cavity wall was performed using a biliary pre-cut needle, with incisions at 3, 6, and 9 o'clock (● **Fig. 3**). Stools were then evacuated by digital squeezing. The procedure was complicated by bleeding, which was successfully treated by endoscopic clipping. The patient was discharged after a brief observation period. At 1-month follow-up she was asymptomatic, and lower endoscopy showed a fully opened diverticulum with no residual stool (● **Fig. 4**). The "rectal pocket syndrome," caused by entrapment of the tied purse string by fired staples, is a rare complication (2.5%) of stapled hemorrhoidopexy and usually requires surgical opening of the pocket [2]. Only two case reports have been published previously in the literature, one of which was managed by releasing the tied suture and the other by creating a divert-

Fig. 2 Pelvic magnetic resonance imaging showed a stool-laden diverticulum in the mesorectal region.

Fig. 3 Endoscopic incision of the cavity.

Fig. 4 The fully opened diverticulum (1 month follow-up).

ing stoma [3,4]. To our knowledge the present case is the first report of endoscopic treatment for rectal pocket syndrome. The procedure appears to be feasible, safe, and effective. The idea was taken from experience of treatment of esophageal rings by means of sagittal incision [5].

Endoscopy_UCTN_Code_TTT_1AQ_2AJ

Competing interests: None

A. Amato¹, A. Vannelli², F. Radaelli¹, S. Paggi¹, E. Rondonotti¹, G. Spinzi¹

¹ Division of Gastroenterology, Valduce Hospital, Como, Italy

² Division of Surgery, Valduce Hospital, Como, Italy

References

- 1 Longo A. Treatment of haemorrhoidal disease by reduction of mucosal and haemorrhoidal prolapse with a circular-suturing device: a new procedure. In: Proceedings of the 6th World Congress of Endoscopic Surgery; 1998 June 3–6; Rome, Italy. Rome: Monduzzi Editori; 1998: 777–784
- 2 Pescatori M, Aigner F. Stapled transanal rectal mucosectomy ten years after. *Tech Colo-proctol* 2007; 11: 1–6
- 3 Cipriani S, Pescatori M. Acute rectal obstruction after PPH stapled haemorrhoidectomy. *Colorectal Dis* 2002; 4: 367–370
- 4 Da Rin F, Faccini M, Pinardi L et al. An uncommon complication after rectal mucosal prolapsectomy. *Osp Ital Chir* 2006; 12: 214–216
- 5 DiSario JA, Pedersen PJ, Bichiş-Canoutas C et al. Incision of recurrent distal esophageal (Schatzki) ring after dilation. *Gastrointest Endosc* 2002; 56: 244–248

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1344419>
Endoscopy 2013; 45: E282–E283
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

A. Amato, MD
 Divisione di Gastroenterologia
 Ospedale Valduce
 Via Dante 11
 22100 Como
 Italy
 Fax: +39-031-308047
arnalam@alice.it