Endoscopic detorsion for sigmoid volvulus using unsedated water-immersion colonoscopy

We report a simple method of endoscopic detorsion for sigmoid volvulus using unsedated water-immersion colonoscopy, which we currently use in the majority of our cases.

Importantly, volvulus detorsion with a colonoscope should be attempted only in patients with an absence of necrotic findings. Endoscopic detorsion is performed without fluoroscopic guidance. A cap or hood attached to the tip of the colonoscope, which maintains a distance between the instrument and the colonic wall, so keeping the luminal direction in view, is used to aid insertion [1]. The water-immersion method involves water infusion in lieu of air insufflation as the principal modality to decrease pain during insertion of the colonoscope [2–4]. Water is infused into the rectum through the biopsy port of the scope using two 50-mL disposable syringes or intermittently infused using a water-jet system. Removal of residual luminal air diminishes the boundary and improves the view. Water accumulates around the tip of the colonoscope and collapses the colon as previously described [2,4].

The colonoscope is inserted with a twisting motion, according to the torsion of the sigmoid colon, and is typically passed under direct observation through the normal-caliber colon to the point of obstruction. This point classically appears as a termination of the lumen in a “whirl sign” [5]. Care needs to be taken to avoid air insufflation while gently attempting to pass the scope through the twisted segment (► **Fig. 1 a**). If this process is successful, the distended proximal segment (► **Fig. 1 b**) is decompressed using endoscopic suction, which often results in spontaneous detorsion. When liquid or soft stools flow out, the success of endoscopic detorsion is confirmed.

Fig. 1 Endoscopic appearances of a sigmoid volvulus showing: **a** the mucosa at the point of twisting; **b** the dilated lumen in the twisted segment of sigmoid colon.

In our experience the success rate of this treatment has been 100%, therefore we believe an endoscopic detorsion is a reasonable treatment for elderly patients with sigmoid volvulus.

Endoscopy_UCTN_Code_TTT_1AQ_2AJ

Competing interests: None

**S. Sugimoto^{1,2}, T. Mizukami³, T. Ito¹,
Y. Tsunoda¹, S. Imamura¹, T. Tamura¹,
S. Nagakubo¹, Y. Morohoshi¹, Y. Koike¹,
Y. Fujita¹, H. Komatsu¹**

¹ Department of Gastroenterology,
Yokohama Municipal Citizens' Hospital,
Yokohama, Japan

² Division of Gastroenterology and Hepa-
tology, Department of Internal Medicine,
School of Medicine, Keio University,
Tokyo, Japan

³ Endoscopy Center, NHO Kurihama
Medical and Addiction Center, Yokosuka,
Japan

References

- 1 Kondo S, Yamaji Y, Watabe H et al. A random-
ized controlled trial evaluating the useful-
ness of a transparent hood attached to the
tip of the colonoscope. *Am J Gastroenterol*
2007; 102: 75–81
- 2 Mizukami T, Hibi T. How I teach my trainees
“water navigation colonoscopy”. *Am J Clin*
Med 2010; 7: 144–146
- 3 Leung FW, Amato A, Ell C et al. Water-aided
colonoscopy: a systematic review. *Gastro-*
intest Endosc 2012; 76: 657–666
- 4 Mizukami T, Yokoyama A, Imaeda H et al. Col-
lapse-submergence method: simple colono-
scopic technique combining water infusion
with complete air removal from the rectosig-
moid colon. *Dig Endosc* 2007; 19: 43–47
- 5 Sugimoto S, Mizukami T, Morohoshi Y et al.
Sigmoid volvulus associated with Chilaiditi's
syndrome. *Intern Med* 2013; 52: 515–516

Bibliography

DOI [http://dx.doi.org/
10.1055/s-0033-1344567](http://dx.doi.org/10.1055/s-0033-1344567)
Endoscopy 2013; 45: E263–E264
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

S. Sugimoto, MD
Division of Gastroenterology and Hepatology
Department of Internal Medicine
School of Medicine, Keio University
35 Shinanomachi, Shinjuku-ku
Tokyo 160-8582
Japan
Fax: +81-3-3353-6247
sugimoto-kei@umin.ac.jp