A rare cause of dysphagia: hairy esophagus

Fig. 1 Esophageal endoscopic view in a 61-year-old man with progressive dysphagia, showing the intact skin graft in the proximal esophagus with the epithelial surface bearing hair.

A 61-year-old man was admitted to our clinic due to progressive dysphagia over the past 4 weeks. He had a history of laryngeal cancer. After failure of radiation therapy 9 months previously, total laryngectomy and bilaterally selective neck dissection had been carried out. Pectoralis major myocutaneous flap reconstruction resulted in the postoperative complication pharyngocutaneous fistula, 3 months after the operation. Esophageal endoscopic examination revealed an intact skin graft in the hypopharynx and proximal esophagus, with the epithelial surface bearing hair, as well as a proximal esophageal stricture (● Fig. 1). Apart from the flap, cervical computed tomography revealed no pathological concerns

(● Fig. 2). Endoscopic dilation of the stricture resulted in improvement of symptoms.

Current treatment of laryngeal and pharyngeal tumors involves a combination of radiation therapy, chemotherapy, and radical surgical resection. With extensive surgical resection, deltopectoral or pectoralis major flaps are frequently used to restore/reconstruct both the anatomy and function of the operative area [1,2]. The reconstructed tube is formed by rotating the skin flap, with the flap placed in the cervical region with the raw surface on the spine and the skin turned on itself, so that the outer epithelial surface becomes the inner lining. However, the epithelial surfaces of flaps may bear hair, particularly in males, and may result in a “hairy esophagus” [3], which further could give rise to the sensation of dysphagia.

Although hair growth has been reported to cease spontaneously within 6–12 months of surgery, on rare occasions it may persist. Where the hair growth in the endopharynx-esophagus is exuberant, patients present with progressive dysphagia, hair-spitting, and choking episodes due to the masses of hair [3]. The endoscopic solution is complex, and bougienage dilatation and hair removal using forceps and/or scissors have been reported [3,4]. Radiation therapy has been used to treat secondary dysphagia due to excessive hair growth [2,3].

Fig. 2 Sagittal (a) and horizontal (b) cervical computed tomography scans showing nothing of note apart from the myocutaneous flap.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AH

Competing interests: None

H. Korkmaz¹, Ç. Elsürer², T. Temel³

¹ Selcuk University, Faculty of Medicine, Departments of Internal Medicine, Division of Gastroenterohepatology, Selcuklu, Konya, Turkey

² Selcuk University, Faculty of Medicine, Department of Otorhinolaryngology, Selcuklu, Konya, Turkey

³ Eskişehir Osmangazi University, Faculty of Medicine, Department of Internal Medicine, Division of Gastroenterology, Eskişehir/Turkey

References

- 1 Piazza C, Tglietti V, Nicolai P. Reconstructive options after total laryngectomy with sub-total or circumferential hypopharyngectomy and cervical esophagectomy. *Curr Opin Otolaryngol Head Neck Surg* 2012; 20: 77–88
- 2 Agha FP, Wimbish KJ. Hirsute esophagus: Clinical and roentgen features. *Gastroenterol Radiol* 1984; 9: 297–300
- 3 McLean GL, Laufer I. “Hairy esophagus.” A complication of pharyngo-esophageal reconstructive surgery in two cases. *Am J Radiol* 1979; 132: 269–270
- 4 Perez LC, Fernandes MR, Aguilera XAC et al. Hairy esophagus: A complication of pharyngo-esophageal reconstructive surgery. *Rev Esp Enferm Dig* 2008; 100: 294–295

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1344572>
Endoscopy 2013; 45: E321
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

H. Korkmaz
Selcuk Üniversitesi
Selçuklu Tıp Fak
Alaeddin Keykubad Kampüsü
42075 Selçuklu
Konya
Turkey
Fax: +332-241-6065
husein68@hotmail.com