Pancreatic stent eroding into the stomach

Fig. 1 One end of a stent protruding into the stomach in a 32-year-old alcoholic man with acute necrotizing pancreatitis.

Fig. 2 The other end of the stent was at the minor papilla.

Fig. 4 Endoscopic view of the stomach following stent removal.

Fig. 3 **a** Fluoroscopic image taken with the endoscope tip close to the gastric end of the stent.
b Fluoroscopic image taken with the endoscope tip close to the papillary end of the stent.

A 32-year-old alcoholic man diagnosed as having acute necrotizing pancreatitis and being managed conservatively developed high-grade fever not responsive to intravenous antibiotics. A percutaneous catheter inserted into the large, infected acute necrotic collection drained frank pus. The patient gradually improved and contrast-enhanced computed tomography (CECT) 3 weeks later revealed a marked reduction in the size of the collection. The transgastric route of the percutaneous catheter was visible on the CECT scan. The catheter continued to drain 300–500 mL of clear pancreatic juice and 4 weeks later the patient underwent endoscopic retrograde

pancreatography (ERP). This revealed a complete pancreas divisum with partial disruption of the dorsal duct. A bridging 5-Fr stent was placed through the minor papilla and the patient showed marked improvement, with the external drainage of pancreatic juice subsiding after 2 weeks. The percutaneous catheter was thereafter removed. After another 4 weeks, the patient underwent another ERP for removal of the stent and to document healing of the ductal disruption. During the passage of the duodenoscope into the stomach a stentlike foreign body was seen in the stomach, protruding through the posterior wall (● Fig. 1). The transpapil-

lary stent placed a few weeks earlier was seen at the papilla but the flanges had migrated inwards (● Fig. 2). Fluoroscopic examination revealed that the transpapillary stent had migrated into the stomach through the pancreatic duct disruption, following the route created by the transgastric percutaneous catheter (● Fig. 3). The stent was removed via the minor papilla (● Fig. 4) and a pancreatogram revealed persistence of ductal disruption. A bridging 5-Fr transpapillary stent was placed. ERP was repeated 4 weeks later and revealed healing of both the ductal disruption and the fistulous opening in the stomach.

Transpapillary stents and nasopancreatic drains have been successfully used for healing of pancreatic ductal disruptions. Stent treatment may be associated with significant complications such as infection, stent block, duct perforation, migration, and stent-induced ductal changes [1].

Endoscopy_UCTN_Code_CPL_1AK_2AD

Competing interests: None

S. S. Rana, H. S. Dhaliwal, V. Sharma, V. Chaudhary, D. K. Bhasin

Department of Gastroenterology, Post Graduate Institute of Medical Education and Research (PGIMER), Chandigarh, India

Reference

- 1 Bhasin DK, Rana SS, Nanda M et al. Comparative evaluation of transpapillary drainage with nasopancreatic drain and stent in patients with large pseudocysts located near tail of pancreas. *J Gastrointest Surg* 2011; 15: 772–776

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1344774>
Endoscopy 2013; 45: E361–E362
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Dr S. S. Rana

Department of Gastroenterology
Post Graduate Institute of Medical Education
and Research (PGIMER)
Sector 12
Chandigarh – 160 012
India
Fax: +91-172-2744401
drsurrinderrana@yahoo.co.in