Endoscopic diagnosis and treatment of a giant duodenal lipoma presenting with gastrointestinal bleeding

Fig. 1 Endoscopic view of a 3-cm long polypoid lesion with multiple ulcerations in the second part of the duodenum in a 66-year-old man with fatigue and acute upper gastrointestinal bleeding.

Fig. 2 Endoscopic ultrasound showing a hyperechoic lesion (11 × 19 mm) originating from the submucosa in the duodenal wall.

A 66-year-old man presented with fatigue and acute upper gastrointestinal bleeding (hemoglobin 92 g/L). He had melena but no signs of hematemesis. Upper gastrointestinal endoscopy revealed a 35 × 15-mm large polypoid lesion with multiple ulcerations in the second part of the duodenum (● Fig. 1). No biopsy samples were taken due to the risk of bleeding. The diagnosis was unclear and the patient underwent endoscopic ultrasound, which demonstrated a hyperechoic lesion measuring 11 × 19 mm in diameter in the submucosa in the duodenal wall with intact muscularis propria (● Fig. 2), suggestive of a lipoma. A subsequent capsule endoscopy excluded distal causes of bleeding in the small intestine. Next, the lesion was removed endoscopically using an endoloop and snare without any complication (● Fig. 3). The resected lesion exhibited multiple ulcerated areas (● Fig. 4). Histological examination demonstrated a duodenal lipoma with large blood vessels in contact with these ulcerated areas (● Fig. 5).

Duodenal lipomas are extremely rare and constitute only one in 600 benign tumors of the gastrointestinal tract [1]. Duodenal lipomas are usually asymptomatic but larger ones can, in rare cases, cause abdominal pain, intestinal obstruction, or hemorrhage [2, 3]. Symptomatic duodenal lipomas should be removed. The current recommendation is endoscopic excision, unless this is technically difficult and war-

Fig. 3 a The lesion was removed using an endoloop and snare (endocut mode effect 2 and forced coagulation effect 2, 30 W, VIO 300 D; ERBE Elektromedizin, Tübingen, Germany). b The resection site 24 hours after polypectomy.

Fig. 4 The resected specimen showing multiple ulcerated areas.

Endoscopy_UCTN_Code_CCL_1AB_2AZ

Competing interests: None

Henrik Thorlacius¹, Håkan Weiber¹, Otto Ljungberg², Jörgen Nielsen³, Ervin Toth³

¹ Section of Surgery, Department of Clinical Sciences, Malmö, Skåne University Hospital, Lund University, Malmö, Sweden

² Section of Pathology, Department of Clinical Sciences, Malmö, Skåne University Hospital, Lund University, Malmö, Sweden

³ Section of Gastroenterology, Department of Clinical Sciences, Malmö, Skåne University Hospital, Lund University, Malmö, Sweden

rants surgical excision. Nonetheless, this unusual case with a duodenal lipoma causing upper gastrointestinal bleeding underlines the clinical importance of endoscopic ultrasound in the workup of patients with unclear submucosal lesions in the gastrointestinal tract.

Fig. 5 Histological section from the lesion showing adipose cells compatible with duodenal lipoma and large blood vessels in contact with the ulceration (arrow).

References

- 1 Mayo CW, Pagtaluman RJG, Brown DJ. Lipoma of the alimentary tract. *Surgery* 1963; 53: 598–603
- 2 Tung CF, Chow WK, Peng YC et al. Bleeding duodenal lipoma successfully treated with endoscopic polypectomy. *Gastrointest Endosc* 2001; 54: 116–117
- 3 Blanchet MC, Arnal E, Paparel P et al. Obstructive duodenal lipoma successfully treated by endoscopic polypectomy. *Gastrointest Endosc* 2003; 58: 938–939

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1344825>
Endoscopy 2013; 45: E385–E386
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Henrik Thorlacius

Department of Clinical Sciences, Malmö
 Section of Surgery
 Skåne University Hospital
 Lund University
 S-205 02 Malmö
 Sweden
 Fax: 46-40-336207
henrik.thorlacius@med.lu.se