Calcium polystyrene sulfonate bezoar in the ileum: diagnosis and treatment with double-balloon endoscopy

An 86-year-old man, admitted for trans-arterial chemoembolization of a hepatoma, developed fever and acute renal failure following the procedure. He was given antibiotic therapy and oral calcium polystyrene sulfonate (30g daily) for hyperkalemia. After 1 week, the patient developed bilious vomiting with abdominal pain. An abdominal X-ray disclosed diffuse dilatation of the small bowel (● Fig. 1) and computed tomography was suspicious for a bezoar in the ileum along with intestinal obstruction (● Fig. 2). A surgeon was consulted but surgical therapy was declined due to the high surgical risk. The patient received conservative medical therapy but his intestinal obstruction failed to resolve. A decision was made to carry out retrograde enteroscopy to remove the bezoar. When the ileum was entered, we found a large, tubular-shaped, firm yellowish bezoar occupying the intestinal lumen (● Fig. 3 and ● Video 1). There were also some ileal ulcers. Irrigation with water and fragmentation using a snare helped mobilize the bezoar. Multiple small brownish granules were seen after the bezoar was fragmented (● Fig. 4) and the aspirated fluid contained resin granules. Given the endoscopic findings and the drug history of the patient, he was diagnosed as having a calcium polystyrene sulfonate resin-associated bezoar. Despite the efforts to remove the bezoar, the patient died of multiple organ failure 1 month later.

Calcium polystyrene sulfonate is an exchange resin used to treat hyperkalemia. A few cases of resin-related bezoars with intestinal obstruction have been reported [1–3]. Such bezoars most often form in critically ill infants [1–3] or in debilitated elderly patients with decreased bowel mobility and prolonged usage of the agent. The treatment of polystyrene sulfonate-related intestinal obstruction is surgery; only one case of non-surgical management has been reported [3]. The present case report documents the endoscopic findings related to resin-associated bezoar. In addition, we have found enteroscopy may be a useful tool in the treatment of intestinal obstruction resulting from the presence of such a bezoar.

Fig. 1 Abdominal X-ray showing diffuse small-bowel dilatation in an 86-year-old man with a drug-induced bezoar.

Fig. 2 Abdominal computed tomography (CT) showing radiopaque material (arrows) in the ileum and which caused the intestinal obstruction.

Video 1

A calcium polystyrene sulfonate resin-associated bezoar in the small intestine removed by endoscopic fragmentation.

Endoscopy_UCTN_Code_TTT_1AP_2AD

Competing interests: None

Fig. 3 Endoscopic view showed a large, tubular and firm, yellowish structure occupying the ileal lumen.

Fig. 4 Post-fragmentation endoscopic view of the bezoar. Multiple small granules (arrows) were observed.

Tien P. Lai¹, Chia W. Yang², Fu Y. Siaop¹, Hsu H. Yen²

¹ Department of Emergency Medicine, Changhua Christian Hospital, Changhua, Taiwan, Republic of China

² Department of Gastroenterology, Changhua Christian Hospital, Changhua, Taiwan, Republic of China

References

- 1 Metlay LA, Klionsky BL. An unusual gastric bezoar in a newborn: Polystyrene resin and candida albicans. *J Pediatr* 1983; 102: 121 – 123
- 2 Garcia-Pardo G, Martinez-Vea A, Auguet T et al. Intestinal obstruction complicating calcium polystyrene sulphonate therapy. *Nephrol Dial Transplant* 1996; 11: 751
- 3 Koneru P, Kaufman RA, Talati AJ et al. Successful treatment of sodium PolyStyrene sulfonate bezoars with serial water-soluble contrast enemas. *J Perinatol* 2003; 23: 431 – 433

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1344835>
Endoscopy 2013; 45: E378–E379
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Hsu H. Yen
 Endoscopy Center
 Department of Gastroenterology
 Changhua Christian Hospital
 Changhua City
 135 Nanhsiao Street Changhua
 500 Taiwan
 China
 91646@cch.org.tw