Intramural hematoma of the colon after polypectomy – an unusual case occurrence

A 43-year-old man was admitted to our hospital for polypectomy because of multiple colonic polyps. He had no history of hemorrhage or abdominal trauma, and was not being treated with anticoagulant therapy. His laboratory results, including coagulation testing, were all normal. A ball-like polyp was identified in the descending colon (● Fig. 1a) and was removed using a snare to leave a 10-mm peduncle. The remaining four small polyps (≤ 5 mm), located in the sigmoid colon and rectum, were removed using forceps.

The following morning, the patient ascended and descended a staircase for approximately 40 minutes. He later complained of having passed bloody stools, with a volume of approximately 150 mL, 32 hours after the polypectomy. We performed another colonoscopy and found bleeding at the site of the peduncle in the descending colon. Three clips were placed to stop the bleeding (● Fig. 1b).

Later the same day, 40 hours after polypectomy, the patient again passed bloody stools, so we immediately performed a third colonoscopy. During this procedure we observed an intramural hematoma (70×50×40 mm) near the site of the peduncle (● Fig. 1c) and noted that the surrounding mucosa was ruptured and bleeding. We therefore injected 10 mL of 1:2000 epinephrine into the hematoma and two clips were used to repair the rupture. Following treatment, the hematoma reduced in size and the bleeding ceased (● Fig. 1d). The patient was kept under observation for 3 days until he passed normal-colored stools.

Significant intramural hematoma can occur in any part of the gastrointestinal tract [1]. The small bowel, particularly the duodenum, is the site most frequently involved; however intramural hematoma of the colon is a relatively rare event [2]. Spontaneous intramural hematomas are

often the result of blunt trauma [3]. The typical causes of nontraumatic spontaneous hematoma are blood dyscrasia, anticoagulant therapy, and other hematologic diseases. Intramural hematoma after polypectomy is rarely observed.

Endoscopy_UCTN_Code_CPL_1AJ_2AC

Competing interests: None

Shaoping Yu, Guangyao Wu, Zongquan Wen, Haiyan Huang

Department of Gastroenterology, Dongguan Kanghua Hospital, Dongguan City, Guangdong Province, China

References

- 1 Lee SH, Lee JH, Park DH et al. Intramural colonic hematoma: complication of anticoagulation with heparin. *Gastrointest Endosc* 2005; 62: 783–784
- 2 Liu Y, Yang S, Tong Q. Spontaneous intramural hematoma of colon. *Clin Gastroenterol Hepatol* 2012; 10: e38
- 3 Li ZL, Wang ZJ, Han JG. Spontaneous perforation of a rectal hematoma. *World J Gastroenterol* 2012; 18: 2438–2440

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1359122>
Endoscopy 2014; 46: E46
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Shaoping Yu, MD
Department of Gastroenterology
Dongguan Kanghua Hospital
1000# Dongguan Avenue
Dongguan City
Guangdong Province
China
Fax: +86-769-23095553
yushaopingmd@163.com

Fig. 1 Colonoscopic views in a 43-year-old man with multiple colonic polyps showing: **a** a ball-like polyp in the descending colon that was removed using a snare; **b** the three clips that were placed on the post-polypectomy peduncle after the patient passed bloody stools 32 hours later; **c** the bleeding intramural hematoma (70×50×40 mm) found near the peduncle during a third colonoscopy that was performed 40 hours after the polypectomy when the patient continued to show evidence of bleeding; **d** the hematoma, which had reduced in size and stopped bleeding, following treatment with epinephrine.