Shiitake mushroom-induced ileus managed using double-balloon enteroscopy

Fig. 1 Abdominal X-ray showing distended jejunum and an air-fluid level.

Fig. 2 Abdominal computed tomography (CT) scan showing a mass with a peculiar shape (arrow).

Fig. 3 Endoscopic view of the jejunum showing the impacted mushroom.

Fig. 4 Abdominal X-ray showing relief of the ileus. The fragmented mushroom had migrated to the descending colon (arrow).

Detection of the cause of dietetic ileus is often difficult. High quality computed tomography (CT) scanning has improved our ability to precisely diagnose a large variety of intra-abdominal and intraluminal disease processes, including the pres-

ence of foreign bodies [1]. Here we report a case of ileus caused by ingestion of shiitake mushroom that could be diagnosed before treatment and could be treated endoscopically without surgery. The patient was a 65-year-old man who had no surgical history. He had a history of stroke and had paralysis on the left side of the body. He presented with abdominal pain and distension that had lasted for 1 week. Abdominal radiography revealed a dilated small bowel with an air-fluid level suggesting intestinal obstruction (● Fig. 1). Abdominal CT scan showed a low density mass and mild distension of the jejunum proximal to the mass. The shape of the mass was irregular and the contrast CT scan suggested a shiitake mushroom (● Fig. 2). We therefore questioned the patient in more detail about his recent diet, and found that he had eaten a meal 4 days previously that

included shiitake mushroom. Based on the dietary history and the CT scan, we diagnosed ileus caused by shiitake mushroom, and we performed double-balloon small-bowel endoscopy on the third day after admission. A large piece of shiitake mushroom was found impacted in the proximal jejunum (● Fig. 3). We crushed and cut away the shiitake mushroom using a snare (SD-5U-1; Olympus, Tokyo, Japan). The entire jejunal mucosa was normal, without ulcers or strictures. After the treatment, abdominal X-ray film showed that the fragmented shiitake mushroom had migrated to the descending colon (● Fig. 4). The patient left the hospital without complications.

Endoscopy_UCTN_Code_TTT_1AP_2AD

Competing interests: None

Meiko Shozushima¹, Ken Ohata¹, Kouichi Nonaka^{1,2}, Nobuyuki Matsuhashi³

¹ Division of Gastroenterology, NTT Medical Center Tokyo, Japan

² Department of Gastroenterology and Hepatology, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

³ Department of Gastroenterology, Kanto Medical Center NTT EC – Gastroenterology, Tokyo, Japan

Reference

- 1 Miura T, Kimura N, Nakamura J et al. Rice cake ileus – a rare and ethnic but important disease status in east-southern Asia. *Intern Med* 2011; 50: 2737–2739

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1359138>
Endoscopy 2013; 45: E437
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Kouichi Nonaka, MD
Division of Gastroenterology
NTT Medical Center Tokyo
5-9-22 Higashi-gotanda, Shinagawa-ku
Tokyo 141-8625
Japan
Fax: +81-3-34486541
nonaka513@gmail.com