



## In vivo appearances of gallbladder carcinoma under magnifying endoscopy and probe-based confocal laser endomicroscopy after endosonographic gallbladder drainage



**Fig. 1** a,b Computed tomography showing the lumen-apposing stent drainage of the gallbladder to the first part of the duodenum.



**Fig. 2** Gallbladder adenocarcinoma in an 87-year-old man, seen at follow-up endoscopy 3 months after endosonographic gallbladder drainage with a lumen-apposing stent: **a** white light endoscopic view; **b** magnified narrow band imaging (NBI) appearance.

An 87-year-old man with multiple medical co-morbidities was admitted for right upper quadrant pain and fever. Computed tomography showed a distended gallbladder with gallstones and pericholecystic fluid compatible with a diagnosis of acute cholecystitis. There was no gallbladder mass. Since the patient had sepsis and was unfit for surgery, endoscopic ultrasound (EUS) drainage of the gallbladder was performed with a lumen-apposing stent (10×15 mm, AXIOS; Xlumena, Mountain View, California, USA) as an alternative to percutaneous cholecystostomy (Fig. 1 a, b). The patient had an uneventful recovery.


Follow-up cholecystoscopy performed 3 months later showed clearance of all stones. However, a 2-cm polypoid lesion was noted at the fundus of the gallbladder. Magnifying narrow band imaging (NBI) showed enlarged and irregular mucosal glands with dilated and corkscrew-appearance microvasculature that was suspicious for malignancy (Fig. 2). Probe-based confocal laser endomicroscopy (CLE) (GastroFlex; Mauna Kea Technologies, France) showed darkened and irregular columnar cells with loss of villous architecture (Fig. 3 a, b). Miniprobe EUS examination (UM-DP12–25R; Olympus, Tokyo, Japan) showed suspicion of tumor

involvement of the gallbladder muscularis propria (Fig. 4). Final histological findings confirmed the presence of a gallbladder adenocarcinoma (Fig. 5). The patient was then treated conservatively as he was too frail to undergo any major surgery.


To our knowledge, this is the first description of a gallbladder adenocarcinoma discovered endoscopically. This was made possible through the use of endosonographic drainage that allowed endoscopic assessment of the gallbladder [1,2]. Magnifying NBI endoscopy has revolutionized the diagnosis of early gastrointestinal neoplasms and is pivotal to performance of endoscopic mucosal resection or submucosal dissection [3]. CLE allows in vivo assessment of cellular architecture and has been shown to be associated with high sensitivity and specificity for diagnosis of Barrett's metaplasia and biliary malignancy [4]. EUS-guided gallbladder drainage in this patient opened the way for application of the above instruments in aiding diagnosis that was not suspected on CT.

Endoscopy\_UCTN\_Code\_CCL\_1AZ\_2AC


**Competing interests:** None



**Fig. 3** a Confocal laser endomicroscopy (CLE) image of the normal-looking gallbladder mucosa, showing villous architecture with columnar epithelium. b CLE image of the gallbladder adenocarcinoma, displaying marked darkened, variably sized glands.



**Fig. 4** Miniprobe examination with endoscopic ultrasound (EUS) showed tumor involvement of the muscularis propria of the gallbladder.



**Fig. 5** A complex cribriform malignant gland is present in the stroma supporting the diagnosis of adenocarcinoma (hematoxylin and eosin [H&E], original magnification  $\times 400$ ).

**Anthony Y. B. Teoh<sup>1</sup>, Anthony W. H. Chan<sup>2</sup>, Philip W. Y. Chiu<sup>1</sup>, James Y. W. Lau<sup>1</sup>**

<sup>1</sup> Department of Surgery, Prince of Wales Hospital, The Chinese University of Hong Kong, Hong Kong SAR

<sup>2</sup> Department of Anatomical and Cellular Pathology, Prince of Wales Hospital, The Chinese University of Hong Kong, Hong Kong SAR

## References

- 1 de la Serna-Higuera C, Pérez-Miranda M, Gil-Simón P et al. EUS-guided transenteric gallbladder drainage with a new fistula-forming, lumen-apposing metal stent. *Gastrointest Endosc* 2013; 77: 303–308
- 2 Itoi T, Binmoeller KF, Shah J et al. Clinical evaluation of a novel lumen-apposing metal stent for endosonography-guided pancreatic pseudocyst and gallbladder drainage (with videos). *Gastrointest Endosc* 2012; 75: 870–876
- 3 Endoscopic Classification Review Group. Update on the Paris Classification of superficial neoplastic lesions in the digestive tract. *Endoscopy* 2005; 37: 570–578
- 4 Nakai Y, Isayama H, Shinoura S et al. Confocal laser endomicroscopy in gastrointestinal and pancreatobiliary diseases. *Dig Endosc* 28. 08. 2013. doi: 10.1111/den.12152 [Epub ahead of print]

## Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0033-1359139>  
*Endoscopy* 2014; 46: E13–E14  
 © Georg Thieme Verlag KG  
 Stuttgart · New York  
 ISSN 0013-726X

## Corresponding author

**Anthony Y. B. Teoh, MD**

Department of Surgery, Prince of Wales Hospital  
 The Chinese University of Hong Kong  
 Shatin, Hong Kong  
 Fax: +852-2637-7974  
[anthonyteoh@surgery.cuhk.edu.hk](mailto:anthonyteoh@surgery.cuhk.edu.hk)