Successful treatment of esophageal fistulas with endoscopic injection of alpha-cyanoacrylate monomer

Fig. 1 The endoscopic appearance in each of the four patients showing in each case the site of the fistula (yellow arrow): **a** before treatment, with the fistula opening visible; **b** after treatment, with the fistula successfully closed. Note: although in patient #4 the fistula was still slightly open, no contrast leakage was observed.

The application of glue is an established treatment for fistulas [1,2]. We consider that the alpha-cyanoacrylate monomer (α -CA) is ideal for glue embolization of intractable fistulas in the gastrointestinal tract because the α -CA glue spreads faster, so creating a bond more rapidly than the other polymer glues, such as n-butyl cyanoacrylate polymer and 2-octyl cyanoacrylate polymer [3]. However, there have been few reports of successful fistula closure using α -CA glue [4].

We herein report on four patients with intractable esophageal fistulas who were successfully treated with endoscopic injection of α -CA. The causes of the fistulas were as follows: an aortoesophageal fistula that developed following a road traffic accident (Patient #1); anastomotic leakage after esophagectomy for esophageal cancer (Patients #2 and #3); perforation of a colon conduit after esophagectomy for esophageal cancer (Patient #4). In the image-guided therapy suite, a mixture of α -CA (Aron alpha A; Sankyo, To-

kyo, Japan) and oily contrast agent (Lipiodol; Guerbet, Tokyo, Japan) in a ratio of 0.3 mL:1.7 mL was endoscopically injected through the fistula using a dispersion tube and a 2.5-mL glue syringe. We used 50% glucose to push the solution from the syringe into the dispersion tube. Because the presence of moisture causes the glue to set, exposure to normal levels of humidity in the air will start polymerization occurring within seconds. However, by making the glue syringe containing the α -CA and Lipiodol air-tight, we were able to inject the solution safely during endoscopy in the image-guided therapy suite.

The procedure was repeated every 1 or 2 weeks until the fistula was closed. In all four patients, the esophageal fistulas were successfully closed with four or six endoscopic injections of α -CA (● Fig. 1). This method is a feasible and safe procedure that may be effective for the treatment of nonhealing esophageal fistulas.

Endoscopy_UCTN_Code_TTT_1AO_2AI

Competing interests: None

Toshiyasu Ojima, Mikihiro Nakamori, Masaki Nakamura, Masahiro Katsuda, Takeshi Iida, Keiji Hayata, Katsunari Takifuji, Makoto Iwahashi, Shuichi Matsumura, Tomoya Kato, Junya Kitadani, Hiroki Yamaue

Second Department of Surgery,
Wakayama Medical University,
Wakayama, Japan

References

- 1 Cameron R, Binmoeller KF. Cyanoacrylate applications in the GI tract. *Gastrointest Endosc* 2013; 77: 846–857
- 2 Yoon JH, Lee HL, Lee OY et al. Endoscopic treatment of recurrent congenital tracheo-esophageal fistula with Histoacryl glue via the esophagus. *Gastrointest Endosc* 2009; 69: 1394–1396

- 3 Katada K, Sano H, Katoh Y et al. Ethyl 2-cyanoacrylate as an embolic agent for cranial arteriovenous malformations. An experimental study. *Acta Radiol Suppl* 1986; 369: 623–626
- 4 Iwata T, Chung K, Toda M et al. Malignant esophagobronchial fistula with severe esophageal stenosis, successfully treated by transbronchial occlusion with cyanoacrylate glue. *Endoscopy* 2012; 44 (Suppl. 02): E317–E318

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1359159>
Endoscopy 2014; 46: E62–E63
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Hiroki Yamaue, MD
Second Department of Surgery
Wakayama Medical University
School of Medicine
811-1, Kimiidera
Wakayama 641-8510
Japan
Fax: +81-73-4466566
yamaue-h@wakayama-med.ac.jp