Endoscopic salvage technique for spontaneous dislocation and tumor ingrowth of a partially covered, self-expandable metallic stent after endoscopic ultrasound-guided choledochoduodenostomy

Fig. 1 Radiograph 7 months after endoscopic ultrasound-guided choledochoduodenostomy in a 67-year-old woman, showing spontaneous dislocation of a partially covered self-expandable metallic stent.

Fig. 2 Endoscopic image 7 months after endoscopic ultrasound-guided choledochoduodenostomy showing the partially covered self-expandable metallic stent trimmed using argon plasma coagulation.

Endoscopic ultrasound-guided choledochoduodenostomy (EUS-CDS) has been established as an alternative method of biliary drainage after failure of endoscopic retrograde cholangiopancreatography (ERCP) [1,2]. Because of its prolonged stent patency and reduced bile leakage, use of the covered self-expandable metallic stent (CSEMS) has recently been favored. We present a case of spontaneous dislocation of a partially covered SEMS (PCSEMS) after EUS-CDS that was successfully rescued by endoscopic intervention.

Fig. 3 Radiograph showing spontaneous dislocation of the partially covered self-expandable metallic stent and stricture of the stent due to tumor ingrowth.

A 67-year-old woman with metastatic cancer of the pancreatic head was admitted to our hospital with obstructive jaundice. She underwent EUS-CDS using a PCSEMS (WallFlex stent, 10×60 mm; Boston Scientific Japan, Tokyo, Japan) without complication (▶ **Video 1**). Seven months later, she developed acute cholangitis. The PCSEMS had dislocated, as was confirmed on computed tomography (▶ **Fig. 1**). Our first attempt was to try to extract the PCSEMS using a snare, but this failed because of tumor ingrowth into the uncovered portion. In our next attempt, the PCSEMS was partially trimmed using argon plasma coagulation with an electro-surgical generator (ICC 200; Erbe Elektromedizin, Tübingen, Germany) at 80 W and with a flow rate of 2 L/min (▶ **Fig. 2**), and we attempted guidewire cannulation through the partially fragmented PCSEMS. A 0.025-inch guidewire was advanced through the ERCP catheter under fluoroscopic guidance (▶ **Video 2**). ERC revealed a stricture of the PCSEMS caused by tumor ingrowth (▶ **Fig. 3**, ▶ **Video 2**). An ERCP catheter and a biliary dilation catheter

could not be passed through the stricture (▶ **Video 2**). In our third attempt, we dilated the stricture using a 6-Fr wire-guided diathermic dilator (Cysto-Gastro-Set; Endo-Flex, Voerde, Germany) using an EGS-100 electrosurgical generator (Olympus, Tokyo, Japan; 30 W in pulse cut slow mode) (▶ **Fig. 4**, ▶ **Video 2**). Finally, a 6-Fr uncovered SEMS (Zilver 635 stent; 10 mm×60 mm, Cook-Japan, Tokyo, Japan) was placed without complication (▶ **Fig. 5**, ▶ **Video 2**).

Video 1

Endoscopic ultrasound-guided choledochoduodenostomy using a partially covered self-expandable metallic stent.

Video 2

Endoscopic salvage technique for a severe stricture of a partially covered self-expandable metallic stent that was dilated using a 6-Fr diathermic dilator, followed by placement of an uncovered self-expandable metallic stent.

Fig. 4 Radiograph showing a 6-Fr diathermic dilator successfully advanced through the severe stricture over a guidewire under fluoroscopic guidance.

Fig. 5 Radiograph showing placement of the uncovered self-expandable metallic stent in the stricture.

Spontaneous dislocation of a PCSEMS after EUS-CDS is a very rare complication [1,2]. In the case described here, we successfully mitigated this situation using endoscopic intervention. In particular, a diathermic

dilator is useful for dilating severe strictures [3,4]. The findings described here suggest that more attention should be paid to the possibility of PCSEMS dislocation when performing EUS-CDS.

Endoscopy_UCTN_Code_CPL_1AL_2AD

Competing interests: None

Hiroshi Kawakami, Masaki Kuwatani, Kazumichi Kawakubo, Taiki Kudo, Yoko Abe, Kimitoshi Kubo, Naoya Sakamoto

Department of Gastroenterology and Hepatology, Hokkaido University Graduate School of Medicine, Sapporo, Japan

References

- 1 Itoi T, Isayama H, Sofuni A et al. Stent selection and tips on placement technique of EUS-guided biliary drainage: transduodenal and transgastric stenting. *J Hepatobiliary Pancreat Sci* 2011; 18: 664–672
- 2 Kawakubo K, Isayama H, Kato H et al. Multi-center retrospective study of endoscopic ultrasound-guided biliary drainage (EUS-BD) for malignant biliary obstruction in Japan. *J Hepatobiliary Pancreat Sci* 11. 09. 2013. doi: doi: 10.1002/jhbp.27 [Epub ahead of print]
- 3 Kawakami H, Kuwatani M, Eto K et al. Resolution of a refractory severe biliary stricture using a diathermic sheath. *Endoscopy* 2012; 44 (Suppl. 02): E119–120
- 4 Kawakami H, Kuwatani M, Kawakubo K et al. Transpapillary dilation of refractory severe biliary stricture or main pancreatic duct by using a wire-guided diathermic dilator (with video). *Gastrointest Endosc* 07. 09. 2013. doi: 10.1016/j.gie.2013.07.055 [Epub ahead of print]

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1359163>
Endoscopy 2014; 46: E58–E59
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Hiroshi Kawakami, MD, PhD

Department of Gastroenterology and Hepatology
 Hokkaido University Graduate School of Medicine
 Kita 15, Nishi 7, Kita-ku
 Sapporo 060-8638
 Japan
 Fax: +81-11-7067867
hiropon@med.hokudai.ac.jp