African Medicinal Plants with Antidiabetic Potentials: A Review

Authors

Aminu Mohammed^{1,2}, Mohammed Auwal Ibrahim^{1,2}, Md. Shahidul Islam¹

Affiliations

¹ Discipline of Biochemistry, School of Life Sciences, University of KwaZulu-Natal (Westville Campus), Durban, South Africa

² Department of Biochemistry, Faculty of Science, Ahmadu Bello University, Zaria, Nigeria

Key words

- Africa
- antidiabetic effects
- diabetes mellitus
- medicinal plants
- North Africa
- East Africa
- West Africa
- Central Africa
- Southern Africa

Abstract

▼
Diabetes mellitus is one of the major health problems in Africa. The conventional oral synthetic antidiabetic drugs available to manage the disease are costly and not readily affordable to the majority of the affected population. Interestingly, the continent is endowed with a tremendous number of medicinal plants that have been explored for their folkloric treatment of diabetes mellitus. Scientific investigations have validated the antidiabetic potentials of a number of these medicinal plants but there is no repository with information on these scientifically investigated plants as a guide for future research. In this review article, all of the *in vivo* antidiabetic studies

conducted between January 2000 and July 2013 on African plants are systematically compiled with a closer look at some relevant plants from the continent's subregions. Plants of the Asteraceae and Lamiaceae families are the most investigated, and West Africa has the highest number of investigated plants. Although promising results were reported in many cases, unfortunately, only a few studies reported the partial characterization of bioactive principles and/or mechanisms of action. It is hoped that government agencies, pharmaceutical industries, and the scientific community will have a look at some of these plants for future research and, if possible, subsequent commercialization.

Introduction

▼
Diabetes mellitus (DM) is a heterogeneous group of metabolic disorders characterized by persistent hyperglycemia [1] and derangement in the metabolism of carbohydrates, fats, and proteins as a result of defects in insulin secretion and/or insulin action [2].

Recent data from the International Diabetes Federation (IDF) indicates that DM affects over 366 million people worldwide and this is likely to increase to 552 million or even more by the year 2030 [3]. In Africa, more than 14 million people have diabetes, accounting for about 4.3% of adults and is responsible for about 401 276 deaths in 2012 in the region [4]. West Africa recorded the highest number of DM cases with Nigeria (3.2 million diabetics) and Côte d'Ivoire (421 023 diabetics) occupying first and second positions, respectively. In Southern Africa, South Africa tops the list (2.0 million diabetics) followed by the Democratic Republic of Congo (737 000 diabetics). Kenya was listed as the fifth country in Africa and the first from the eastern region of Africa

(720 730 diabetics), while Cameroon (517 860 diabetics) recorded the highest figure from the central region. North Africa had the least number of diabetics among the African subregions with no single nation in the top ten list of African countries with DM [4].

At present, different approaches are used to control DM using modern synthetic antidiabetic drugs in addition to lifestyle modification. This includes sulphonylureas (glibenclamide), glucosidase inhibitors (acarbose), and biguanide (metformin). However, these synthetic oral hypoglycemic agents have characteristic profiles of serious side effects, which include hypoglycemia, weight gain, gastrointestinal discomfort, nausea, liver and heart failure, and diarrhea [5] in addition to being rather costly and not affordable by the majority of African populations. These limitations coupled with an exponential increase in the prevalence of DM motivate researchers to scientifically validate the folkloric use of a number of antidiabetic African medicinal plants as possible alternative therapies. This is partly because herbs and natural products form an important compo-

received August 22, 2013
revised Dec. 31, 2013
accepted January 1, 2014

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1360335>
Published online February 17, 2014
Planta Med 2014; 80: 354–377
© Georg Thieme Verlag KG
Stuttgart · New York ·
ISSN 0032-0943

Correspondence

Dr. Md. Shahidul Islam
School of Life Sciences
University of KwaZulu-Natal
(Westville Campus)
Durban 4000
South Africa
Phone: + 27 3 12 60 87 17
Fax: + 27 3 12 60 79 42
islamd@ukzn.ac.za

Fig. 1 Map of Africa showing the different subregions. (Color figure available online only.)

nent of the health care delivery system in African countries [6]. According to the World Health Organization (WHO) [7], 80% of the population in many African countries depend almost entirely on traditional medicines, herbal medicines in particular, for their primary health care needs [8,9]. This is attributed to the perceived effectiveness of the plant-based therapies as well as the availability of these medicinal plants because the continent accounts for about 25% of the total number of higher plants in the world where more than 5400 medicinal plants were reported to have over 16300 medicinal uses [10].

In Africa, herbal medicines are usually provided by a traditional healer, who utilizes natural products in curing many diseases. They have different local methods to diagnose DM in their patients, as they do not rely on laboratory investigations. This is achieved through identifying symptoms like frequent urination, sexual dysfunction, swollen legs, hands and stomach, obesity, fatigue, and profuse sweating during the consultation process. In some cases, they direct the patients to urinate on locally prepared formulations and return after a couple of days with the results of a diagnosis. At present, DM is among the diseases which are most extensively treated with traditional medicines using medicinal plants. This is evident by the propensity of the ethnobotanical surveys for medicinal plants used for the management of DM from different African subregions that include West [11, 12], East [13–15], North [16–18], Southern [19,20], and Central Africa [21]. Interestingly, scientific investigations have confirmed the efficacy of a number of these plant-derived formulations on DM, but presently there is no comprehensive review and/or repository that exists in the literature of these scientifically investigated African antidiabetic medicinal plants that cover the whole of Africa. In this article, we conducted an exhaustive review of all scientifically investigated African antidiabetic medicinal plants whose results have been published between January 2000 and July 2013. The scientific community, government agencies, and

Fig. 2 Subregional distributions of investigated African medicinal plants with antidiabetic effects.

pharmaceutical industries may use this as a possible guide for future research.

Results and Discussion

A map of Africa indicating the subregions of the continent as used in this review is presented in **Fig. 1**. A total of 185 plants species from 75 families in Africa have been investigated for antidiabetic effects. The information obtained on these plants includes scientific and common names, families, parts of the plant used, solvent used, and whether the crude extracts or fractions were used in the course of investigation. From the results, plants from the West African subregion account for 51.69% of all the plants investigated for antidiabetic potentials in Africa over the period mentioned (**Fig. 2**). More than 90% of all documented plants from this region emanate from Nigeria, with few data from Ghana, Senegal, Benin, Togo, and Côte d'Ivoire (**Table 1**). Reports on the antidiabetic effects of North African plants account for 21.91% (**Fig. 2**) and originated from Morocco, Egypt, Algeria, Tunisia, Sudan, and Libya (**Table 3**). On the other hand, 12.92% of antidiabetic African plants were reported from Southern Africa (**Fig. 2**) with most of the studies originating from South Africa (**Table 4**). The remaining parts, East (**Table 2**) and Central (**Table 5**) Africa, recorded 7.87 and 5.61%, respectively, of African medicinal plants with antidiabetic effects. The results also indicated that plants from the Asteraceae and Lamiaceae families received a lot of attention in all parts of Africa (**Fig. 3**). On the other hand, analysis of the investigated parts of the plant indicated that the leaf was the most scientifically investigated part (**Fig. 4**).

More importantly, promising results were reported in many cases but, unfortunately, only a few studies reported detailed characterization of bioactive principles. Data available indicated that only six plants from northern and two from western regions received partial characterization of a possible active ingredient that could be responsible for their antidiabetic effects (**Table 6**). In order to provide a full view of the antidiabetic potentials of these African medicinal plants, the scientifically investigated medicinal plants are categorized into subregions (**Fig. 1**) and discussed more thoroughly. The discussions are based on the subregions and the criteria used for highlighting a plant that has potency of the reported antidiabetic activity, except in the case of

Table 1 List of scientifically investigated antidiabetic medicinal plants from West Africa.

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day)/ Extract/Fraction	Type of effects	Type of DM/ Model used	Country	References
<i>Acacia albidia</i> Del.	Ana Tree	Mimosaceae	Root	200 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[139]
<i>Acalypha wilkesiana</i> Müll. Arg.	Copperleaf	Euphorbiaceae	Leaf	100, 200, and 300 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[140]
<i>Adansonia digitata</i> L.	Baobab	Bombacaceae	Stem bark	100, 200, and 400 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[141]
<i>Azelaia Africana</i> Smith.	Counter wood	Fabaceae	Root	62.5, 125, 250, 500, and 1000 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[142]
<i>Ageratum conyzoides</i> L.	Billygoat weed	Asteraceae	Seed	100, 200, 400 and, 500 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[143, 144]
<i>Alchornea cordifolia</i> Müll. Arg.			Leaf	200, 400, and 800 mg/kg bw Butanol fraction	Antihyperglycemic	Type 1/Animal	Nigeria	[145]
<i>Allium cepa</i> L.	Onion	Liliaceae	Bulb	200, 250, and 300 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[85, 146–150]
<i>Allium sativum</i> L.	Garlic	Liliaceae	Bulb	200, 250, and 300 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[85, 150]
<i>Aloe perryi</i> Baker.	Cashew	Liliaceae	Leaf	2 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[34]
<i>Anacardium occidentale</i> L.		Anacardiaceae	Leaf	34, 200, 300, and 400 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1 and 2/Animal	Nigeria	[22–25, 151]
<i>Anisopous mannii</i> N. E. Br.	–	Asclepiadaceae	Stem	100, 200, 300, and 400 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[152, 153]
<i>Anthocheista djalensis</i> A. Chev.	Cabbage tree	Loganiaceae	Leaf/Stem/Root	1 g/kg bw Methanol extract and its fractions	Antihyperglycemic	Type 1/Animal	Nigeria	[154]
<i>Axonopus compressus</i> P. Beauv.	Blanket grass	Fabaceae	Leaf	250, 500, and 1000 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[155]
<i>Azadirachta indica</i> A. Juss.	Neem	Meliaceae	Leaf	70 and 400 mg/kg bw Ethanol extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria/ Ghana	[26–31]
<i>Bauhinia rufescens</i> Lam.	–	Caesalpinaceae	Leaf	200, 300, and 400 mg/kg bw Methanol extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[156, 157]
<i>Bridelia ferruginea</i> Benth.	–	Euphorbiaceae	Root bark	250 mg/kg bw Methanol extract	Antihyperglycemic and hypoglycemic	Type 1/Animal	Nigeria	[158]
<i>Carica papaya</i> L.	Pawpaw	Caricaceae	Seed	100, 200, 300, and 400 mg/kg bw Aqueous extract	Antihyperglycemic and hypolipidemic	Type 1/Animal	Nigeria	[159, 160]
<i>Carum carvi</i> L.	Caraway	Apiaceae	Fruit	5, 10, 20, 40, and 80 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[161–164]
<i>Cassia italic</i> Mill.	Italian senna	Caesalpinaceae	Leaf	200 mg/kg bw Aqueous/ethanol extract	Antihyperglycemic and antioxidative	Type 1/Animal	Nigeria	[165, 166]
<i>Catharantus roseus</i> L.	Madagascar periwinkle	Apocynaceae	Leaf	Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[77, 167]
<i>Cela pentandra</i> L. Gaertn.	Silk-cotton tree	Bombacaceae	Stem bark	250, 400, 800, and 1500 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[168]
<i>Chrysophyllum cainito</i> L.	Star apple	Anacardiaceae	Leaf	10, 20, and 30 g/l Aqueous extract in drinking water	Antihyperglycemic	Type 1/Animal	Côte d'Ivoire	[169]
<i>Cinchona calisaya</i> Wedd.	–	Rubiaceae	Stem bark	50 and 100 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[170]

continued

Table 1 Continued

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day)/ Extract/Fraction	Type of effects	Type of DM/ Model used	Country	References
<i>Cissampelos mucronata</i> A. Rich.	Heart-leaved vine	Menispermaceae	Leaf	200, 400, and 800 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[171]
<i>Cissampelos owariensis</i> P. Beauv.	Velvet leaf	Menispermaceae	Leaf	100 and 200 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[172]
<i>Citrus aurantium</i> L.	Bitter orange	Rutaceae	Fruit	400 and 800 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[173]
<i>Citrus paradise</i> Macfad.	Grapefruit	Rutaceae	Seed	100, 300, and 600 mg/kg bw Aqueous extract	Antihyperglycemia	Type 1/Animal	Nigeria	[174]
<i>Clausena lansium</i> Lour. Skeels.	Wampee	Rutaceae	Stem bark	100 mg/kg bw Methanol and dichloro methane extract	Antihyperglycemic and insulinotropic	Type 1/Animal	Nigeria	[175]
<i>Cnestis ferruginea</i> D. C.	–	Connaraceae	Leaf	250 mg/kg bw Methanol and ethyl acetate extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[176]
<i>Combretum micranthum</i> G. Don.	–	Combretaceae	Leaf	100, 200, and 400 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[177]
<i>Commelina Africana</i> L.	–		Leaf	500 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[144]
<i>Curcumin longa</i> L.	Curcuma	Zingiberaceae	Root	250 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[178]
<i>Daniella oliveri</i> Bull. Misc.	Daniella	Caesalpinaceae	Root	250 mg/kg bw Aqueous extract	Antihyperglycemic and inhibition of glycolytic enzymes	Type 1/Animal	Nigeria	[179, 180]
<i>Detarium microcarpum</i> Guill. & Perr.	Sweet detar	Caesalpinaceae	Leaf	Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[153]
<i>Ficus asperifolia</i> L.	Sand paper tree	Moraceae	Stem	400, 800, and 1200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[181]
<i>Ficus exasperate</i> Vahl.	White fig tree	Moraceae	Leaf	100, 200, and 300 mg/kg bw Aqueous extract	Antihyperglycemic and insulinotropic	Type 1/Animal	Nigeria	[182, 183]
<i>Ganoderma lucidum</i> Curtis. P. Karst.	Hemlock varnish shelf	Ganodermataceae	Fruit	50 mg/kg bw Ethyl acetate and butanol fractions	Antihyperglycemic	Type 1/Animal	Nigeria	[184]
<i>Gongronema latifolium</i> Benth.	Anaranth globe	Asclepiadaceae	Leaf	2, 25, 75, 100, 200, and 400 mg/kg bw Aqueous extract	Antihyperglycemic, antihyperlipidemic and insulinotropic	Type 1/Animal	Nigeria	[32–39, 147]
<i>Hibiscus sabdariffa</i> L.	Red sorrel	Malvaceae	Calyces	0.5 mg/ml Aqueous and ethanol extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[40–42]
<i>Holarrhena floribunda</i> G. Don.	False rubber tree	Apocynaceae	Leaf	100, 250 and 500 mg/kg bw Ethanol extract and its solvent fractions	Antihyperglycemic	Type 1/Animal	Côte d'Ivoire	[185]
<i>Homalium letestui</i> Pellegr.	–	Flacourtiaceae	Root	500, 750, and 1000 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[186]
<i>Hunteria umbellata</i> K. Schum. Hallier f.	Flantueh	Apocynaceae	Stem	50, 100, and 200 mg/kg bw Ethanol extract	Antihyperglycemic and insulinotropic	Type 1/Animal	Nigeria	[187]
<i>Hymenocardia acida</i> Tul.	Red-heart	Phyllanthaceae	Leaf	250, 500, 1000, and 2000 mg/kg bw Methanol extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[188]
<i>Hyptis suaveolens</i> Poit.	Mint weed	Lamiaceae	Leaf	750 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[189]
<i>Indigofera pulchra</i> L.	Indigofera	Papilionaceae	Leaf	50, 100, 200, 250, 500 and 1000 mg/kg bw Butanol fraction	Antihyperglycemic and hypoglycemic	Type 1/Animal	Nigeria	[43–46]
<i>Irvingia gabonensis</i> Aubry-Lecomte ex O'Rorke. Baill.	Wild mango	Asteraceae	Stem bark	200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[190] continued

Table 1 Continued

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day)/ Extract/Fraction	Type of effects	Type of DM/ Model used	Country	References
<i>Khaya senegalensis</i> Desr. A. Juss.	African mahogany	Meliaceae	Stem	50, 100, and 150 mg/kg bw Aqueous extract/oil	Antihyperglycemic	Type 1/Animal	Nigeria	[191, 192]
<i>Leptadenia hastate</i> Pers.	–	Leguminosae	Leaf	300 mg/kg bw Ethanol extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[153, 193]
<i>Loranthus micranthus</i> L.	Mistletoe	Loranthaceae	Leaf	250 and 400 mg/kg bw Methanol extract and its fractions	Antihyperglycemic	Type 1/Animal	Nigeria	[194, 195]
<i>Mangifera indica</i> L.	Mango	Anacardiaceae	Leaf	0.5 and 1.0 g/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[196]
<i>Mammea africana</i> Sabine.	African apple	Cuttiferae	Stem bark	30, 60, and 90 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[197]
<i>Melanthera scandens</i> Schumacher. Roberty.	–	Asteraceae	Leaf	37, 74, and 111 mg/kg bw Ethanol extract and its fractions	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[198]
<i>Mimosa invisa</i> Mart.	Sleeping plant	Fabaceae	Leaf	Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[153]
<i>Momordica charantia</i> L.	Bitter melon	Cucurbitaceae	Leaf	250, 400, and 500 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[199, 200]
<i>Morinda lucida</i> Benth.	Brimstone tree	Rubiaceae	Stem bark	50, 100, 200, and 240 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[201, 202]
<i>Moringa oleifera</i> Lam.	Horseradish	Moringaceae	Leaf	100, 200, and 300 mg/kg bw Aqueous Extract	Antihyperglycemic	Type 1/Animal	Nigeria	[203]
<i>Musa sapientum</i> L var. paradisiacal. Sucker.	Banana	Musaceae		5, 10, 250, and 500 mg/kg bw Methanol extract	Antihyperglycemic and GIT transit time	Type 1/Animal	Nigeria	[204, 205]
<i>Musanga cecropioides</i> R.Br. & Tedlie.	Umbrella tree	Cecropiaceae	Stem bark	250, 500, and 1000 mg/kg bw Aqueous and ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[206]
<i>Nauclaea latifolia</i> S. M.	Bishop's head	Rubiaceae	Root/Stem/ Leaf	200, 400, and 1000 mg/kg bw Ethanol and hexane extracts	Antihyperglycemic and su- crose and maltase inhibitions	Type 1/Animal	Benin/Ni- geria	[47–51]
<i>Newbouldia laevis</i> P. Beauv.	African Border Tree	Bignoniaceae	Leaf	100, 200, and 400 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[207]
<i>Ocimum gratissimum</i> L.	African/Clove basil	Lamiaceae	Leaf	250, 400, 500, 600, 800, 1000, and 1500 mg/kg bw Methanol extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[53–60]
<i>Ocimum suave</i> Wild.	Hoary basil	Lamiaceae	Leaf	800 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[208]
<i>Oxytenanthera abyssinica</i> A. Rich Munro.	Bindura bamboo	Gramineae	Leaf	25 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Benin	[50]
<i>Parinari microphylla</i>	Boxleaf azara	Chrysobalanaceae	Seed	500 mg/kg bw Ethanol extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[209]
<i>Parinari excels</i> (Guinea Plum)			Bark	100 and 300 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Senegal	[210]
<i>Parkia biglobosa</i> Jacq.	African locust bean	Fabaceae	Seed	6 g/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[211]
<i>Parquetina nigrescens</i> Afzel. Bullock.		Asclepiadaceae	Leaf	1000 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[212]
<i>Persea Americana</i> Mill.	Avocado	Lauraceae	Seed	450 and 900 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[213]
<i>Phyllanthus amarus</i> L.	Stone breaker	Euphorbiaceae	Whole plant	150, 300, 500, 600, and 1000 mg/ kg bw Aqueous/hydroalcoholic extract	Antihyperglycemic and insulinotropic	Type 1/Animal	Nigeria/ Togo	[61–64] continued

Table 1 Continued

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day)/ Extract/Fraction	Type of effects	Type of DM/ Model used	Country	References
<i>Phyllanthus niruri</i> L.	Gulf leaf flower	Euphorbiaceae	Whole plant	240 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[214]
<i>Picralima nitida</i> Stapf.	Picralima	Apocynaceae	Pulp/Seed	250 and 648 mg/kg bw Ethanol and aqueous extract	Antihyperglycemic and hypoglycemic	Type 1/Animal	Nigeria/ Benin	[65, 66]
<i>Raphia hookeri</i> G.Mann & H.Wendl.	Wine palm	Palmaeae	Stem	50, 100, and 200 mg/kg bw Aqueous extract	Antihyperglycemic and hypolipidemic	Type 1/Animal	Nigeria	[215]
<i>Rauvolfia vomitoria</i> Afzel.	Swizzle stick	Apocynaceae	Fruit	400 and 800 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[173]
<i>Sansevieria senegambica</i> Baker.	African flax	Agavaceae	Root	100, 200, and 300 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[216]
<i>Sarcocephalus latifolius</i> Sm. E. A. Bruce.	African peach	Rubiaceae	Root	250 mg/kg bw Aqueous extract	Antihyperglycemic and inhi- bition of glycolytic enzymes	Type 1/Animal	Nigeria	[179, 180]
<i>Senna occidentalis</i> L.	Stink weed	Caesalpinaceae	Leaf	200, 300, and 450 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[217]
<i>Senna alata</i> L. Roxb.	Candle bush	Leguminosae	Leaf	250 mg/kg bw Methanol extract	Antihyperglycemic and hypoglycemia	Type 1/Animal	Nigeria	[158]
<i>Senna siamea</i> Lam.	Thailand shower	Caesalpinaceae	Leaf/Stem bark	250, 500, 1000, 2000, and 3000 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[218, 219]
<i>Setaria megaphylla</i> Steud. Dur. & Schinz.	Ribbon grass	Phyllanthaceae	Root	200, 400, and 600 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[220]
<i>Sida acuta</i> Burm. f.	Wire weed	Malvaceae	Leaf	200 and 400 mg/kg bw Ethanol and methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[221]
<i>Sphagnetocola trilobata</i> L. Pruski.	Creeping oxeye	Asteraceae	Leaf	50 mg/kg bw Aqueous extract	Antihyperglycemic and antioxidative	Type 1/Animal	Nigeria	[222]
<i>Sphenocentrum jollyanum</i> Pierre.	Sorghum bicolor	Menispermaceae	Root	50, 100, and 200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[223]
<i>Stachytarpheta angustifolia</i> Mill. Vahl.	Devil's coach whip	Verbenaceae	Whole part	250, 500, 750, and 1000 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[224]
<i>Telfairia occidentalis</i> Hook. f.	Fluted pumpkin	Cucurbitaceae	Seed	2, 100, and 250 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[225, 226]
<i>Terminalia catappa</i> L.	Bengal almond	Combretaceae		0.6 ml/20 kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Côte d'Ivoire	[227]
<i>Treulia Africana</i> Decne. fruit	African bread- fruit	Moraceae	Root	200 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Nigeria	[228]
<i>Triplochiton scleroxylon</i> K. Schum.	African white wood	Sterculiaceae	Stem bark	Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[229]
<i>Vernonia amygdalina</i> Del.	Bitter leaf	Compositae	Leaf	50, 100, 200, 250, 400, and 500 mg/kg bw Hexane and ethylacetate extract	Antihyperglycemic, insulino- tropic, and antioxidant	Type 1/Animal	Nigeria	[67–82]
<i>Vernonia colorata</i> Schreb.	Star-flowered bitter-tea	Compositae	Leaf	100 and 300 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Senegal	[230]
<i>Vitex doniana</i> Wild.	African black plum sweet	Verbenaceae	Stem bark	100 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Nigeria	[62]
<i>Viscum album</i> L.	Mistletoe	Loranthaceae	Whole plant	2, 50, and 100 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Nigeria	[34, 231]
<i>Zingiber officinale</i> L. Roscoe.	Ginger	Zingiberaceae	Rhizome	200, 250, and 300 mg/kg bw Aqueous extract	Antihyperglycemic and α -amylase inhibition	Type 1/Animal	Nigeria	[83–89]

Table 2 List of scientifically investigated antidiabetic medicinal plants from East Africa.

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day)/ Extract/Fraction	Type of effects	Type of DM/ Model used	Country	Reference
<i>Aspilia pluriseta</i> Schweinf.	Dwarf aspilia	Asteraceae	Root	50, 100, and 200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Kenya	[232]
<i>Bidens pilosa</i> L.	Spanish needle	Asteraceae	Leaf	50, 100, and 200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Kenya	[232]
<i>Caylusea abyssinica</i> Fresen. Fisch. & Mey.	–	Resedaceae	Leaf	100, 200, and 300 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Ethiopia	[133]
<i>Catha edulis</i> Vahl.	Bushman's tea	Celastraceae	Root	50, 100, and 200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Kenya	[232]
<i>Erythrina abyssinica</i> Lam.	Red-hot-poker	Fabaceae	Stem bark	50, 100, and 200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Kenya	[232]
<i>Ficus sycomorus</i> L.	Fig-mulberry	Moraceae	Stem bark	50, 100, and 150 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Kenya	[233]
<i>Moringa stenopetala</i> Baker f.	Cabbage tree	Moringaceae	Leaf	500 mg/kg bw Butanol fraction	Antihyperglycemic and hypoglycemic	Type 1/Animal	Ethiopia	[137]
<i>Pappea capensis</i> Eckl. & Zeyh.	Jacket plum	Sapindaceae	Leaf/Stem bark	100 and 200 mg/kg bw Aqueous and ethylacetate extract	Antihyperglycemic	Type 1/Animal	Kenya	[234]
<i>Pentas schimperi</i> A. Rich.	–	Rubiaceae	Leaf	500 and 1000 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Ethiopia	[235]
<i>Strychnos heningsii</i> Gilg.	Red bitter berry	Loganiaceae	Leaf	50, 100, and 200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Kenya	[232]

West Africa where citations were used as a criteria in addition to the potency of the results.

West Africa

Anacardium occidentale L. (Anacardiaceae), or cashew, is perhaps one of the most cited plants from West Africa. The antidiabetic activity of a stem-bark methanol extract was investigated in a fructose-fed type 2 diabetes (T2D) model of rats [22]. Treatment with 200 mg/kg body weight (bw)/day of the extract given orally significantly ameliorated the changes in plasma glucose, lipid profile, malonyldialdehyde, urea, and creatinine induced by an enriched fructose diet, but showed no effect on plasma alkaline phosphatase levels. Extract treatment reduced plasma glucose levels by almost 40% in fructose-fed type 2 diabetic rats. In another study, oral administration of an ethanolic extract of inner bark and fractions at various doses administered caused a significant decrease in blood glucose levels in a type 1 diabetes (T1D) model of rats [23]. The crude extract decreased blood glucose by 36.8% at 700 mg/kg bw when different fractions at 300, 30, and 200 mg/kg bw/day indicated a glycemic decrease of 18.4%, 15.6%, and 17.3%, respectively. Furthermore, bioactivity-guided fractionation of the ethanolic extract led to fractions that displayed diverse polyphenolic compounds, which are known for their hypoglycemic effect. The methanolic leaf extract, orally administered at 400 mg/kg bw/day, decreased the blood glucose levels of alloxanized rats by 20.8% after 4 hours of treatment compared to 47.63% for tolbutamide, a standard antidiabetic drug [24]. More recently, Ukwenya et al. [25] reported that administration of a methanolic leaf extract at a dose of 300 mg/kg bw recovered the beta cell damage in a T1D model of rats. Although an extensive antidiabetic study on this plant is yet to be done, the preliminary data indicates that the stem bark contains more therapeutically active antidiabetic phytochemicals than the other parts of the plant.

Another highly cited antidiabetic plant is *Azadirachta indica* A. Juss. (Meliaceae), which is commonly referred to as neem tree. The hypoglycemic and antihyperglycemic effects of the leaves in a T1D model of rats have been investigated [26,27]. Oral treatment of the aqueous extract at 400 mg/kg bw/day was found to decrease fasting blood glucose by 54% compared to the control. Further studies were conducted by Akinlola et al. [28], who reported that the ethanolic extract orally administered at 500 mg/kg bw/day prevented intestinal lesions and decreased hyperglycemia (87.5%) in an STZ-induced T1D model of rats. Moreover, in 2010, Akinlola et al. [29] evaluated the chronic treatment of diabetic rats with an *A. indica* leaf ethanolic extract at 500 mg/kg bw/day orally on blood glucose, pancreatic islet histopathology, and oxidative status of the pancreas. The results obtained were quite promising, the fasting blood glucose of the extract-treated group dropped to 50%, the number of β -cells was improved and, similarly, islet histology showed a marked improvement with a significant decrease in oxidative stress. In 2011, this research group also indicated that the oral administration of *A. indica* leaf extract at 500 mg/kg bw/day ameliorated the renal damage in an STZ-induced T1D model of rats [30]. In another study, an orally administered ethanolic leaf extract at 400 mg/kg bw/day was reported to decrease fasting blood glucose by 50% and further prevented the alterations posed by DM on immunological and hematological parameters [31] which have clinical significance in the control of atherosclerosis and other diabetes-associated vascular complications.

Gongronema latifolium Benth. (Asclepiadaceae) is another plant that received much attention as a hypoglycemic and antihyperglycemic agent in Africa. Almost all of its parts are claimed to have an antidiabetic effect. Treatment with the aqueous leaf extract at various dosages was found to decrease the fasting blood glucose in an STZ-induced T1D model of rats [32–35]. Fasting blood glucose dropped by 30.4% compared to untreated diabetic animals. Akah et al. [36] reported that treatment with both aque-

Table 3 List of scientifically investigated antidiabetic medicinal plants from North Africa.

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day)/ Extract/Fraction	Type of effects	Type of DM/Model used	Country	Reference
<i>Ajuga reptans</i> L.	Herb ivy	Labiatae	Whole part	10 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Morocco/ Tunisia	[89–93]
<i>Allium cepa</i> L.	Onion	Liliaceae	Bulb	100 and 400 mg/kg bw Aqueous extract	Antihyperglycemic, antihyper- lipidemic, and antioxidative	Type 1 and 2/Animal and DM patients	Sudan	[94–96]
<i>Anabasis articulata</i> Forsk. Moq.	Jointed anabis	Chenopodiaceae	Leaf	400 mg/kg bw Methanol extract	Antihyperglycemic and hypoglycemic	Type 1/Animal	Algeria	[236]
<i>Artemisia herba-alba</i> Asso.	White wormwood	Lamiaceae	Aerial part	2 g/kg bw Hydroalcoholic extract	Antihyperglycemic	Type 2/Animal	Algeria	[237]
<i>Balanites aegyptiaca</i> L.	Desert date/Hegleg	Balanitaceae	Fruit	80 mg/kg bw Ethanol and aqueous extracts	Antihyperglycemic, hypoglyce- mic, and α -amylase inhibition	Type 1/Animal	Egypt	[97, 98]
<i>Capparis spinosa</i> L.	Flinders rose	Capparidaceae	Fruit	20 and 1500 mg/kg bw Aqueous extract	Antihyperglycemic and anti- hyperlipidemic	Type 2/Animal	Morocco	[103, 238]
<i>Carum carvi</i> L.	Caraway	Apiaceae	Fruit/Oil	2 ml and 20 mg/kg bw Oil and aqueous extract	Antihyperglycemic and anti- hyperlipidemic	Type 1/Animal	Morocco/ Egypt	[99–102]
<i>Centaurium erythraea</i> Rafn.	Bitter herb	Gentianaceae	Aerial part/Leaf	200 mg/kg bw Aqueous extract	Antihyperglycemic and antioxidative	Type 1 and 2/Animal	Algeria/ Morocco	[239]
<i>Chamaemelum nobile</i> L.	Chamomile	Asteraceae	Aerial part	20 mg/kg bw Aqueous extract	Antihyperglycemic, repress glucoseogenesis, and improve insulin sensitivity	Type 2/Animal	Morocco	[104]
<i>Cinnamomi cassia</i> Nees & T. Nees, J. Presl.	Cinnamon twigs	Lauraceae	Stem bark	200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Egypt	[240]
<i>Cleome droserifolia</i> De- lile. Forsk.	Cleome herb	Capparaceae	Leaf	0.31 g/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Egypt	[241]
<i>Cuminum cyminum</i> L.	Cumin	Apiaceae	Oil	2 ml/kg bw	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Egypt	[102]
<i>Curcumin longa</i> L.	Curcuma	Zingiberaceae	Root	300 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	Egypt	[110]
<i>Cynara corrigera</i> L.	Wild artichoke	Asteraceae	Root	1.5 g/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Libya	[242]
<i>Eucalyptus globulus</i> Labill.	Waxy bloom	Myrtaceae	Leaf	200 and 400 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Algeria	[243]
<i>Trigonella foenum grae- cum</i> L.	Fenugreek	Leguminosae	Seed	1.5 g/kg bw Methanol extract	Antihyperglycemic and α -amylase inhibition	Type 1/Animal	Egypt	[98]
<i>Globularia alypum</i> L.	Black thorn	Globulariaceae	Leaf	20 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Morocco	[244]
<i>Guiera senegalensis</i> J.F. Gmel.	Moshi medicine	Combretaceae	Leaf	200 and 400 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Sudan	[243]
<i>Inula viscosa</i> L.	False yellow head	Asteraceae	Aerial part	20 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Morocco	[245]
<i>Lepidium sativum</i> L.	Pepper grass	Cruciferae	Stem	20 mg/kg bw Aqueous extract	Antihyperglycemic and hypoglycemic	Type 1/Animal	Morocco	[246]
<i>Magnifera indica</i> L.	Mango	Anacardiaceae	Leaf	250 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Egypt	[247]
<i>Morus alba</i> L.	White mulberry	Moraceae	Leaf/Root bark	200 and 400 mg/kg bw Ethanol extract	Antihyperglycemic, antihyper- lipidemic, and antioxidative	Type 1/Animal	Egypt	[105–107] continued

Table 3 List of scientifically investigated antidiabetic medicinal plants from North Africa.

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day)/ Extract/Fraction	Type of effects	Type of DM/Model used	Country	Reference
<i>Nigella sativa</i> L.	Black seed	Ranunculaceae	Seed	2 g and 300 mg/kg bw Petroleum ether extract	Antihyperglycemic and enhancement of GLUT4 expression	Type 1 and 2/Animal	Egypt	[108–111]
<i>Ocimum basilicum</i> L.	Sweet basil	Lamiaceae	Leaf	20 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Morocco/ Egypt	[102, 248]
<i>Panax ginseng</i> L.	Korean ginseng	Araliaceae	Root	22.5 mg/rat	Antihyperglycemic, anti- oxidative, and insulinotropic	Type 1/Animal	Egypt	[249]
<i>Psidium guajava</i> L.	Guava	Myrtaceae	Leaf	250 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Egypt	[247]
<i>Rubus fruticosus</i> L.	Blackberry	Rosaceae	Leaf	100 mg/kg bw Aqueous extract	Antihyperglycemic and hypoglycemic	Type 1/Animal	Egypt	[244]
<i>Salvia officinalis</i> L.	Garden sage	Lamiaceae	Leaf	200 and 400 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Algeria	[243]
<i>Spergularia purpurea</i> Pers. G. Don.	Purple sand spurry	Caryophyllaceae	Whole plant	10 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Morocco	[250, 251]
<i>Suaeda fruticosa</i> L. Forssk.	Shrubby seablite	Chenopodiaceae	Aerial part	0.8 mg/kg bw/min Aqueous extract	Antihyperglycemic and hypoglycemic	Type 1/Animal	Morocco	[252]
<i>Thymelaea hirsute</i> L. Endl.	Spur flax	Thymelaceae	Aerial part	3 mg/kg bw Aqueous extract	Antihyperglycemic and hypoglycemic	Type 1/Animal	Morocco	[253]
<i>Thymus vulgaris</i> L.	Thyme	Lamiaceae	Oil	2 ml/kg bw	Antihyperglycemic and hypoglycemic	Type 1/Animal	Egypt	[102]
<i>Trigonella foenum graecum</i>	Fenugreek	Leguminosae	Seed	1.5 g/kg bw Methanol extract	Antihyperglycemic, hypo- glycemic, and α -amylase inhibition	Type 1/Animal	Egypt	[98]
<i>Triticum repens</i> L.	Couch grass	Gramineae	Root	20 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Morocco	[254]
<i>Zingiber officinale</i> L. Roscoe	Ginger	Zingiberaceae	Rhizome	4 ml/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1/Animal	Egypt	[255]
<i>Ziziphus spina-christi</i> L.	Christ's Thorn Jujube	Rhamnaceae	Leaf	100 and 450 mg/kg bw Butanol fraction	Antihyperglycemic, insulino- tropic, and α -amylase in- hibition	Type 1 and 2/Animal	Egypt	[112–114]

Table 4 List of scientifically investigated antidiabetic medicinal plants from Southern Africa.

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day) Extract/Fraction	Type of effects	Type of DM/Model used	Country	Reference
<i>Azela afficana</i> SM. ex Pers.	Counter wood	Fabaceae	Root	100 and 200 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	South Africa	[256]
<i>Allium cepa</i> L.	Onion	Liliaceae	Bulb	0.5 and 2.0% Aqueous extract	Antihyperglycemic	Type 1/Animal	South Africa	[257]
<i>Aloe excels</i> Berger.	–	Aloaceae	–	–	Antihyperglycemic	Type 1/Animal	Zimbabwe	[258]
<i>Aloe ferox</i> Mill.	Bitter aloe	Xanthorrhoeaceae	Leaf	300 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	South Africa	[259]
<i>Aloe greatheadii</i> var. <i>davyana</i>	Spotted aloe	Asphodelaceae	Leaf	300 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	South Africa	[259]
<i>Artemisia afra</i> Jacq.	African worm-wood	Asteraceae	Leaf	50 and 100 mg/kg bw Aqueous extract	Antihyperglycemic, anti-oxidative, and insulinotropic	Type 1/Animal	South Africa	[115, 116]
<i>Brachylaena discolor</i> D. C.	Coast silver oak	Asteraceae	Leaf	50 and 150 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	South Africa	[260]
<i>Bryophyllum pinnatum</i> Lam.	Life plant	Crassulaceae	Leaf	400 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	South Africa	[117]
<i>Camellia sinensis</i> L.	White tea	Theaceae	–	0.5 g/100 ml Aqueous extract	Antihyperglycemic	Type 1/Animal	South Africa	[261]
<i>Clausena anisata</i> Wild.	Horse wood	Rutaceae	Leaf/Root	100, 200, 400, and 800 mg/kg bw Methanol extract	Antihyperglycemic	Type 1/Animal	South Africa	[262]
<i>Catharantus roseus</i> L. G. Don.	Madagascar periwinkle	Apocynaceae	Leaf	500 mg/kg bw Methanol extract	Antihyperglycemic and hypoglycemic	Type 1/Animal	South Africa	[263]
<i>Euclea undulata</i> Thunb. var. <i>myrtina</i>	Fire fighter's blessing	Ebenaceae	Root	25 and 50 mg/kg bw Acetone extract	Antihyperglycemic	Type 1/Animal	South Africa	[264]
<i>Hypoxis hemerocallidea</i> Fisch. & C. A. Mey.	African potato	Hypoxidaceae	Corm	50, 100, 200, 400, and 800 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	South Africa	[265, 266]
<i>Leonotis leonurus</i> L. R. Br.	Throw-hort	Lamiaceae	Leaf	125 250, and 500 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	South Africa	[267]
<i>Momordica charantia</i> L.	Bitter melon	Cucurbitaceae	Whole plant	50, 100, 200, 400, and 800 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	South Africa	[268, 277]
<i>Prosopis glandulosa</i> Torr.	Honey mesquite	Fabaceae	Pods	100 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 1 and 2/Animal	South Africa	[269]
<i>Psidium guajava</i> L.	Guava	Myrtaceae	Leaf	Aqueous extract	Antihyperglycemic	Type 1/Animal	South Africa	[270]
<i>Raphia gentiliana</i> De Wild.	–	Arecaceae	Fruit	0.2 g/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	DR Congo	[118]
<i>Rhus chirindensis</i> Baker F.	Red currant	Anacardiaceae	Stem bark	50, 100, 200, 400, and 800 mg/kg bw Aqueous extract	Anti-hyperglycemic	Type 1/Animal	South Africa	[271]
<i>Sclerocarya birrea</i> A. Rich. Hochst.	Jelly plum	Anacardiaceae	Stem bark	100, 200, 400, and 800 mg/kg bw Methanol and dichloromethane extract	Anti-hyperglycemic	Type 1/Animal	South Africa	[119–122]
<i>Strychnos henningsii</i> Gilg.	Red bitter berry	Loganiaceae	Stem bark	125, 250, and 500 mg/kg bw Aqueous extract	Antihyperglycemic, anti-oxidative, and hypoglycemic	Type 2/Animal	South Africa	[272]
<i>Sutherlandia frutescens</i> R. Br. var. <i>incana</i> E. MEY.	Cancer brush	Fabaceae	Leaf	2.5 g/100 ml Aqueous extract	Antihyperglycemic, antihyperlipidemic, and prevention of insulin resistance	Type 1/Animal	South Africa	[123–125]

Table 5 List of scientifically investigated antidiabetic medicinal plants from Central Africa.

Scientific name	Common name	Family	Part(s) used	Dosage (orally/day)/ Extract/Fraction	Type of effects	Type of DM/ Model used	Country	Reference
<i>Anacardium occidentale</i> L.	Cashew	Anacardiaceae	Leaf	35, 175, and 250 mg/kg bw Methanol and its solvent fractions	Antihyperglycemic	Type 1/Animal	Cameroon	[273]
<i>Bersama engleriana</i> Gurke.	Winged-bersama	Melanthaceae	Leaf	300 and 600 mg/kg bw Aqueous extract	Antihyperglycemic and antihyperlipidemic	Type 2/Animal	Cameroon	[126, 127]
<i>Canarium schweinfurthii</i> Engl.	Bush candle tree	Burseraceae	Stem bark	150 and 300 mg/kg bw Methanol and dichloromethane extract	Antihyperglycemic	Type 1/Animal	Cameroon	[274]
<i>Cela pentandra</i> L.	Silk-cotton tree	Bombacaceae	Stem bark	40 and 75 mg/kg bw	Antihyperglycemic	Type 2/Animal	Cameroon	[268]
Gaertn.	Watermelon	Cucurbitaceae	Seed	Methylene chloride/methanol extracts 50 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Cameroon	[226]
<i>Cucumeropsis mannii</i> Naudin.	White seed melon	Cucurbitaceae	Seed	50 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Cameroon	[226]
<i>Cucurbita moschata</i> Duchesne ex Poir.	Butternut squash	Leguminosae	Pods	50 mg/kg bw Aqueous extract	Antihyperglycemic	Type 1/Animal	Cameroon	[226]
<i>Dichrostachys glomerata</i> Chiov.	Chinese lantern	Cucurbitaceae	Seed	400 mg Ethanol extract	Antihyperglycemic and antihyperlipidemic	Type 2/DM Patients	Cameroon	[125]
<i>Dracena arborea</i> Wild.	Dragon tree	Dracaenaceae	Root	500 Aqueous and 100 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Cameroon	[130]
<i>Kalanchoe crenata</i> Andr. Haw.	Never ride	Crassulaceae	Whole plant	50 and 68 mg/kg bw Methanol extract	Antihyperglycemic, antihyperlipidemic, and antioxidative	Type 1/Animal	Cameroon	[131]
<i>Lagenaria siceraria</i> L.	Bottle gourd	Cucurbitaceae	Seed	50 mg/kg bw Ethanol extract	Anti-hyperglycemic	Type 1/Animal	Cameroon	[226]
<i>Sclerocarya birrea</i> A. Rich. Hochst.	Jelly plum	Anacardiaceae	Stem bark	150 and 300 mg/kg bw Methanol and dichloromethane extract	Antihyperglycemic and insulinotropic	Type 1 and 2/Animal	Cameroon	[275, 276]
<i>Telfaria occidentalis</i> Hook. f.	Fluted pumpkin	Cucurbitaceae	Seed	50 mg/kg bw Ethanol extract	Antihyperglycemic	Type 1/Animal	Cameroon	[259]
<i>Terminalia superba</i> Engl. & Diels	Ofram tree	Combretaceae	Stem bark	150 and 300 mg/kg bw Methanol and dichloromethane extract	Antihyperglycemic	Type 1/Animal	Cameroon	[274]

Fig. 3 Percentage of distributions in families of African medicinal plants with antidiabetic effects.

Fig. 4 Frequency of plant part used in anti diabetic studies.

ous and methanolic extracts and fractions at 800 mg/kg bw showed a significant antihyperglycemic effect in an experimentally-induced T1D model of rats. The highest glycemic reduction recorded was 43% at 32 h post-treatment of the crude aqueous extract compared to 35% for the methanolic extract. The methanolic fraction showed the highest decrease in glycemia (30%) at 400 mg/kg bw after 32 hours of treatment. Furthermore, these extracts showed a protective effect on the activity of some cardiac enzymes, which are crucial in the management of DM, in the same model of diabetic rats [37]. In an attempt to investigate the possible mechanism of action, Adebajo et al. [38] reported the insulinotropic and glucose lowering effects of the combined root and stem bark methanolic extracts (1:1) and various fractions when administered orally at a dose of 100 mg/kg bw/day. It was observed that the effect of the combined extracts was far better

than the individual actions of the fractions. Both the extract and fractions caused an insulin release and lowered the blood glucose levels better than glibenclimide in glucose-loaded rats and INS-1 cell lines. In another study, oral administration of ethanolic roots and twig extracts at 200 and 400 mg/kg bw/day showed a protective effect against alterations on the markers of kidney functions in a T1D model of rats [39]. Fasting blood glucose dropped by more than 60% for a single twig administered via orogastric intubation compared to 43% for the root ethanolic extracts.

The hypoglycemic and hypolipidemic effects of *Hibiscus sabdariffa* L. (Malaceae) calyces have been reported [40]. The authors showed that the oral administration of the aqueous extract at 0.5 mg/ml/day alleviated the oxidative stress in a T1D model of rats [41], which is comparable to vitamin C and glibenclimide. The extract decreased fasting blood glucose by almost 70%,

Table 6 Scientifically investigated African antidiabetic medicinal plants with identified possible bioactive compounds.

Scientific name	Common name	Family	Part(s) used	Possible compound(s) present/isolated	Country	Reference
<i>Anacardium occidentale</i>	Cashew	Anacardiaceae	Leaf/n-hexane/diethylether fractions	Polyphenols, terpenoids, alkaloids, and flavonoids	Nigeria	[23]
<i>Balanites aegyptiaca</i>	Desert date/Hegleg	Balanitaceae	Fruit/chloroform: methanol: water	Diosgenin	Egypt	[97,98]
<i>Carum carvi</i>	Caraway	Apiaceae	Fruit oil	D-limonene, benzyl alcohol, O-cresol, isomenthone, methyl chavicol, D-carvone, perillaldehyde, and β -patchoullene	Morocco/Egypt	[99–102]
<i>Gongronema latifolium</i>	Amaranth globe	Asclepiadaceae	Root stem/methanol extract	α and β -amyrin cinnamates, lupenyl cinnamates, lupenyl acetate, and two other unknown triterpenoids Y and Z	Nigeria	[38]
<i>Morus alba</i>	White mulberry	Moraceae	Root bark/water-methanol fractions	Morusin, cyclomorusin, neocyclomorusin, kuwanon E, 2-arylbenzo furan, moracin M, betulinic acid and methyl ursolate, and two other triterpenes, betulinic acid and methyl ursolate, mulberroside A, 5,7,2'-trihydroxy flavanone-4'-O- β -D-glucoside, and albanols A and B	Egypt	[105–107]
<i>Trigonella foenum graecum</i>	Fenugreek	Leguminosae	Seed/chloroform: methanol: water fractions	Diosgenin	Egypt	[98]
<i>Ziziphus spina-christi</i>	Christ's Thorn Jujube	Rhamnaceae	Leaf/butanol fraction	Chritinin-A	Egypt	[112–114]

which is comparable to ascorbic acid and glibenclimide. To investigate the possible mode of action, Adedayo and Ganiyu [42] revealed that the extract inhibited the activities of the two key enzymes involved in carbohydrate digestion, alpha glucosidase and alpha amylase, which are very crucial in the management of T2D mellitus.

Using an alloxan-induced T1D rat model, Tanko et al. [43,44] demonstrated the glucose lowering effect of the hydromethanolic extract of the *Indigofera pulchra* L. (Papilionaceae) leaf when administered orally at 250, 500, and 1000 mg/kg bw/day. Surprisingly, the lower dosage that was administered decreased the glucose levels by 50% after 8 hours post-treatment. The daily intraperitoneal administration of ethyl acetate and *n*-butanol fractions of this extract also showed a hypoglycemic effect [45, 46] and was able to restore the DM-induced hematological alterations.

Nauclea latifolia S.M. (Rubiaceae) is among the most widely used traditional plants in the management of DM in the different parts of Nigeria. In an attempt to investigate this claim by traditional herbalists, Gidado et al. [47] showed that the oral treatment of the aqueous leaf extract of *N. latifolia* at 200 mg/kg bw/day significantly decreased blood glucose levels in a T1D model of rats, which was 45% within a 4-hour period of treatment. Further studies by this group indicated that both the aqueous and ethanolic leaf extracts at 100, 200, and 400 mg/kg bw/day significantly decreased the blood glucose levels in a dose-dependent manner when administered orally. At 400 mg/kg bw, aqueous and ethanolic leaf extracts lowered fasting blood glucose by 31.7% and 36.1%, respectively, [48] and were able to inhibit maltase and sucrase activities *in vitro* but not *in vivo* [49]. Other researchers reported that stem and root ethanolic extracts administered intraperitoneally at 25 mg/kg bw/day have potent immunosuppressive effects on T cell proliferation in an STZ-induced T1D model of rats [50] and could improve the antioxidant status and hormonal changes in another diabetes rat model [51].

Ocimum gratissimum L. (Lamiaceae) is used widely as a condiment or spice in different cultural settings in Africa. It is traditionally used for the management of DM [52]. Intraperitoneal administration of 400 mg/kg bw/day of *O. gratissimum* methanolic leaf extract significantly decreased the blood glucose level in both normal and T1D models of rats by 56% and 69%, respectively [53, 54]. Oral treatments at 500, 1000, and 1500 mg/kg bw/day of the aqueous leaf extract dropped the fasting blood glucose by more than 50% [55]. Mohammed et al. [56] also showed that the administration of 500 mg/kg bw of this extract caused a reduction in blood glucose by 81.3% 24 hours after administration. Moreover, the *O. gratissimum* leaf ethanolic extract treated at 200 mg/kg bw/day prevented the alteration of germinal epithelium, distortion of seminiferous tubules, as well as vacuolation of seminiferous tubules in an STZ-induced T1D model of rats [57,58]. More recently, Oguanobi et al. [59] reported that oral administration of the *O. gratissimum* leaf extract at 100, 200, and 300 mg/kg bw/day had a blood glucose lowering effect and the ability to alleviate derangements in serum and biliary bilirubin, cholesterol, and electrolytes in a neonatal STZ-induced T2D rat model [60].

Phyllanthus amarus L. (Euphorbiaceae) (stone breaker or gulf flower) is another highly cited antidiabetic plant from West Africa. The hypoglycemic potential of the aqueous leaf extract of *P. amarus* was investigated in an alloxan-induced T1D rat model. The extract at a dose of 260 mg/kg bw produced a significant ($p < 0.05$) reduction in blood glucose level by 112%, 61%, and 31% at 24 hours, 7 days, and 14 days of oral administration, respectively. Furthermore, the reduction was dose-dependent and dropped by 82%, 41%, and 16% after 24 hours, 7 days, and 14 days of oral administration, respectively [61]. Owolabi et al. [62] reported that oral treatment of the aqueous extract at 100 mg/kg bw/day decreased fasting blood glucose by 46.53% compared to 66.6% for insulin treatment in a T1D model of rats. In another study from Togo, Povi et al. [63] indicated that *P. amarus* whole plant aqueous and hydroalcoholic extracts at 500 and 1000 mg/kg bw/day had both hypoglycemic and hypolipidemic effects

after 15 days of oral administration. Aqueous leaf and seed extracts at 150, 300, and 600 mg/kg bw/day were also shown to have antihyperglycemic, antihyperlipidemic, and cardioprotective effects as well as an insulin sensitizing effect in a T2D model of rats [64].

The hypoglycemic effect of the seed aqueous extract of *Picralima nitida* Stapf. (Apocynaceae) has been investigated in a T1D model of rats [65]. The extract at 648 mg/kg bw decreased the fasting blood glucose by about 19.46% and 75.5% in normoglycemic and alloxanized rats within 3 and 6 hours, respectively. However, a contradictory finding was reported by Igboasoiyi et al. [66] where the seed (250 mg/kg bw/day), but not fruit pulp, extract showed no hypoglycemic effect.

The folkloric use of *Vernonia amygdalina* Del. (Compositae) (bitter leaf) in the management of DM is widely documented and this corroborates with the propensity of antidiabetic studies conducted in both type 1 and type 2 animal models of diabetes [67–74] and human subjects [75]. In all studies reported, *V. amygdalina* was found to significantly reduce the hyperglycemia in T1D models of rats. The alterations on the markers of kidney functions were prevented by the leaf ethanolic extract administered at 400 mg/kg bw/day by gastric intubation and the fasting blood glucose was decreased by more than 80% [76]. Oral administration of this extract was also reported to prevent the macrovascular complications associated with DM [77]. In an attempt to further investigate the hypoglycemic effect of this plant, Akah et al. [78] reported that the hexane/ethyl acetate fraction obtained from the leaf methanolic extract possessed an antidiabetic effect when treated orally at 80, 160, and 320 mg/kg bw/day. The histological and hematological results showed no alterations on the full tissue architecture and other parameters analyzed. It was also reported that *V. amygdalina* leaf ethanolic extract at 100 mg/kg bw/day prevented the alteration of germinal epithelium, distortion of seminiferous tubules, as well as vacuolation of seminiferous tubules in an STZ-induced T1D model of rats [58]. The synergistic or antagonistic effects of this plant with other traditionally claimed antidiabetic plants have also been documented in numerous studies [79–82].

Another plant that has received much attention from researchers and has been used in the management of DM is *Zingiber officinale* L. (Zingiberaceae) (ginger). Its hypoglycemic effect has been reported in STZ and glucose-induced diabetic rat models [83–85]. The glucose-lowering effect began after 30 minutes of intraperitoneal administration of a rhizome aqueous extract at 2, 4, and 8 mg/kg bw. At 200, 250, and 300 mg/kg bw, the decreases in the blood glucose levels were recorded as 51.4%, 56.9%, and 56.7%, respectively. In another study, Iranloye et al. [86] investigated the antihyperglycemic and antioxidant effects of an orally administered aqueous extract of *Z. officinale* at 500 mg/kg bw/day in both type 1 and fructose-fed T2D models of rats. The extracts decreased blood glucose by 48.23% in type 1, and 83.5% in T2D models of rats. The extract also showed the high radical scavenging ability and improvement of insulin biosynthesis in this experiment. It has also been reported that *Z. officinale* at doses of 250 and 500 mg/kg bw/day could inhibit oxidative stress and inflammation by enhancing antioxidant enzymes and TNF- α activity in an STZ-induced T1D model of rats [87]. Arikawe et al. [88], in their studies, indicated that this extract could prevent the diabetes and insulin resistant-associated effects on spermatogenesis in an experimentally induced diabetes rat model.

North Africa

Ajuga iva L. (Labiatae) is among the most frequently investigated antidiabetic plants from northern Africa, especially Morocco and Algeria. The hypoglycemic and hypolipidemic effects of the aqueous extract of *A. iva* have been investigated in a number of T1D models of rats [89–91]. Oral administration of the extract at 10 mg/kg bw/day reduced the plasma glucose levels by 69.73% after a 6-hour post-treatment period and 87.3% in a subchronic dosing of 28 days in diabetic rats compared to 21.4% and 18.4% in normoglycemic rats, respectively. Furthermore, the *in vivo* antioxidant effects of this plant's extracts have also been reported in an STZ-induced T1D model of rats [92]. The 4-week supplementation of 0.5% aqueous extract of *A. iva* prevented oxidative damages by decreasing lipid peroxidation, and improved the activities of plasma and tissue antioxidant enzymes in experimentally induced diabetic rats. Hamden et al. [93] reported that the phytoecdysteroids rich extract of *A. iva* prevented the diabetes-associated microvascular complications in alloxan-induced diabetic rats when administered orally for fifteen days.

Allium cepa L. (Liliaceae) (onion) is widely distributed throughout the African region and is among the most cited plants from West and North Africa. The antidiabetic effect of *A. cepa* was evaluated in both animal models and human subjects [94]. Fresh crude slices (100 g) of *A. cepa* were given to type 1 and 2 diabetic human subjects. In type 1 diabetic subjects, a 50% reduction in the blood glucose level was observed at 4 hours post-treatment compared to insulin-treated (70.8%) patients. In type 2 diabetic subjects, about 20% reduction was observed compared to 37.5% in insulin-injected subjects [94]. It was also reported by El-Demerdash et al. [95] that *A. cepa* at a dose of 1 mL or 0.4 g/100 g bw restored the biochemical and antioxidant status altered by alloxan injection in a T1D model of rats. The concentration of thiobarbituric acid reactive substances and the activity of glutathione S-transferase in plasma, liver, testes, brain, and kidneys were significantly increased in alloxan diabetic rats. These increases were completely prevented in *A. cepa*-treated rats. Various compounds were isolated and correlated positively as being responsible for the hypoglycemic effect of *A. cepa*. Phenolics and sulphur compounds, such as cysteine and allyl propyldisulphide, have been associated with this effect [94]. Some studies linked the observed hypoglycemic effects of this plant to the essential oils [96].

In Egyptian traditional medicine, *Balanites aegyptiaca* L. (Balaniaceae) is popularly used as an oral hypoglycemic agent. The protective effect of orally treated aqueous and ethanolic extracts at 80 mg/kg bw/day against liver damage, and hypoglycemic and hypolipidemic effects in alloxan-induced type 1 diabetic rats were investigated [97]. Liver glycogen, serum insulin, leptin, and testosterone levels were increased in treated rats while glucagon, total lipids, total cholesterol, triglyceride level, and transaminase activities were significantly decreased. In another study, oral administration of *B. aegyptiaca* fruit aqueous extract at 1500 mg/kg bw/day decreased fasting blood glucose by 24% compared to the diabetic control. The dose-dependent inhibition of alpha amylase and glucose-6-phosphatase activities with an increase in glucose-6-phosphate dehydrogenase and phospho-fructokinase activities were reported [98]. The TLC and HPLC fingerprints of this extract led to the identification of a marker compound, diosgenin [98].

It has been reported that *Carum carvi* L. (Apiaceae) has been used for medicinal purposes since ancient times [99]. Hypoglycemic and hypolipidemic effects of the aqueous fruit extract orally administered at 20 mg/kg bw/day have been evaluated in an STZ-in-

duced T1D model of rats [100, 101]. The extract significantly decreased the blood glucose levels by more than 50% compared to nondiabetic rats within two weeks of administration but did not increase the insulin levels of normal rats. Moreover, treatment with the extract significantly decreased the serum lipid profile levels. *C. carvi* has been classified as a good hypoglycemic agent with high radical scavenging activity [102]. Some of the active compounds isolated from this plant include d-limonene, benzyl alcohol, o-cresol, isomenthone, methyl chavicol, d-carvone, perillaldehyde, and β -patchoullene [102].

In Morocco, *Chamaemelum nobile* L. (Asteraceae) is used locally for the treatment of DM and its complications. The antihyperglycemic effect of the aqueous fruit extract administered at 20 mg/kg bw/day has been reported in an obese T2D model of mice [103]. The postprandial hyperglycemia dropped significantly by more than 80%, and the increase in body weight was completely prevented in mice treated with this extract. In another study, Lemhadri et al. [104] reported that the extract-treated animals demonstrated a decrease in endogenous glucose production compared to the diabetic control group. It also improved insulin sensitivity in peripheral tissues, which was confirmed by an increased glucose utilization in an STZ-induced diabetic model of mice [104].

Morus alba L. (Moraceae) (white mulberry) is another important plant used in the management of DM in northern Africa. The hypoglycemic effect of root bark flavonoid-rich fractions has been investigated [105]. The blood glucose levels of treated animals at 600 mg/kg bw for ten days dropped by about 50% compared to that of the diabetic control group. Insulin levels were significantly increased, while the peroxide levels were significantly decreased in this study. Morusin, cyclomorusin, neocyclomorusin, kuwanon E, 2-arylbenzofuran, moracinM, betulinic acid, and methylursolate were the compounds isolated from *M. alba*. El-Sayyad et al. [106] investigated the effect of the *M. alba* leaf extract on both micro- and macrovascular complications associated with DM. In their study, treatment with the extract at 100 mg/kg bw/day prevented the increase in maternal serum glucose, alterations in lipid profiles, and creatine phosphokinase activity, as well as retinal neurotransmitters including acetylcholine, adrenaline, noradrenaline, serotonin, histamine, dopamine, and gamma amino butyric acid. Furthermore, cataract and retinopathy were also prevented in the treated groups. In another study, mulberroside A, 5,7,2'-trihydroxyflavonone-4-O- β -D-glucoside, and albanols A and B were also isolated from fractions derived from the ethanolic root bark of *M. alba* [107]. The authors also indicated that oral administration of these fractions at 500 mg/kg bw/day significantly prevented the oxidative damage induced by hyperlipidemia in rats.

Nigella sativa L. (Ranunculaceae), popularly used as a spice, has also been reported to be used traditionally in the treatment of diabetes and was investigated in a number of studies [108]. The hypoglycemic potentials of *N. sativa* oil from the seed have been reported [109]. The oil significantly reduced the blood glucose levels in an STZ-induced T1D model of rats within six weeks of oral administration. It was reported by the authors that the blood glucose lowering effect might be stimulated by extra hepatic tissues rather than by insulin release. In another study, *N. sativa* seed extract decreased the blood glucose levels by almost 81% and normalized fructosamine, hemoglobin, and albumin levels in experimentally induced diabetic animals within 30 days of oral administration of 300 mg/kg bw [110]. The extract was able to ameliorate the diabetes-associated oxidative damages in the

same study. The regulation of hepatic glycolytic and gluconeogenic enzyme activities were considered as a possible mode of action in this study. Benhaddou-Andaloussi et al. [111] have recently demonstrated the antidiabetic effect of *N. sativa* seed ethanolic extract when administered at 2 g/kg bw/day, and which is mediated through an insulin-sensitizing action by enhancing acetyl-CoA carboxylase phosphorylation, a major component of the insulin-independent AMPK signaling pathway, and by enhancing muscle GLUT4 expression.

The antidiabetic effect of the butanol leaf extract and christinin A (a major saponin glycoside) of *Ziziphus spina-christi* L. (Rhamnaceae) has been investigated in both type 1 and type 2 diabetic models of rats [112, 113]. In type 2 but not in the T1D model, pretreatment of both the extract and the isolated compound at 100 mg/kg bw/day indicated a clear improvement in the oral glucose tolerance test and mediated glucose-induced insulin release. Both the extract and christinin A caused a significant decrease in blood glucose levels by 24% and 22%, respectively, after 60 minutes, and increased serum insulin levels in a T2D model of rats. The extract was also reported to cause no damage to the kidneys, liver, or hematological parameters after 30 days of oral administration. Michel et al. [114] have also demonstrated the hypoglycemic effect of *Z. spina-christi* leaf ethanolic extract when fed orally at 200 mg/kg bw/day. A significant increase in serum insulin and C-peptide levels were observed in extract-treated animals. The extract also ameliorated the oxidative damages and prevented the protein glycosylation induced by diabetes. The activities of liver glucose-6-phosphatase and alpha amylase (IC₅₀ of 0.3 mg/mL) were inhibited by this extract, but significantly increased the activity of glucose 6-phosphate dehydrogenase. HPLC and spectrophotometric determination revealed a flavonoid as a marker compound named christinin A in this plant.

Southern Africa

Artemisia afra Jacq. (Asteraceae) is mostly identified by its aromatic odor. It is widely available and is being used to treat DM. In 2011, Afolayan and Sunmonu [115] reported that the orally administered aqueous leaf extract at 50 and 100 mg/kg bw/day significantly decreased the blood glucose levels by more than 50% compared to the diabetic control, with a concomitant increase in insulin levels. The extract also showed a high antioxidant effect by increasing the levels of antioxidant enzymes and decreasing lipid peroxidation. Similar effects of the extract on antioxidant defense systems in the liver and kidneys were also observed within three weeks of oral administration at the same doses in diabetic animals [116].

Bryophyllum pinnatum Lam. (Crassulaceae), popularly known as "good luck" or "life plant" is widely used in the management of DM by the majority of the African populace. Ojewole [117] has reported the antidiabetic, antinociceptive, and anti-inflammatory effects of *B. pinnatum* in rats using fresh egg albumin-induced pedal (paw) edema, and in an STZ-induced T1D model of rats. *B. pinnatum* leaf aqueous extract treated orally at 400 mg/kg bw significantly and dose-dependently decreased fasting blood glucose in diabetic rats by almost 50% within 8 hours. A similar effect was observed in albumin-induced acute inflammation of the rat hind paw.

Raphia gentiliana De Wild. (Arecaceae) is one of the most popular plants used in the treatment of several disease ailments in the Democratic Republic of Congo. Its fruits are commonly consumed as food. The hypoglycemic effect of the aqueous fruit extract has been investigated in normoglycemic human subjects and glu-

cose-induced hyperglycemic animals [118]. After one and two hours post-treatment at 200 mg/kg bw, the fasting blood glucose dropped by 27% and 56%, respectively, in diabetic mice. In human subjects, the glycemic index (signifying glucose absorbed into the blood after a meal) and load index (total glucose content in normal subjects) recorded were -3.60% and -1.36% , respectively, which are within the recommended range. The results indicate the preventive role of *R. gentiana* fruit in glucose absorption, which could be associated with the active principles present in the fruit extract.

The stem bark, roots, and leaves of *Sclerocarya birrea* A. Rich. Hochst. (Anacardiaceae) are widely used in South Africa and African countries as folk medicine in the treatment DM. The hypoglycemic effect of the stem bark aqueous extract has been investigated in normal and STZ-induced diabetic rats [119, 120]. *S. birrea* stem bark aqueous extract at 800 mg/kg bw significantly and dose-dependently decreased the fasting blood glucose in both normal and diabetic-treated rats, with a maximum reduction capacity of 50.16%, which was comparable with that of chlorpropamide (62.44%) after 8 hours post-oral treatment. Furthermore, Musabayane et al. [121] previously established the beneficial effect of the *S. birrea* stem bark aqueous extract treated orally at various doses (60, 120, and 240 mg/kg bw) on markers of kidney and cardiovascular functions in diabetic rats. It significantly decreased the blood glucose levels and the levels of Na^+ and K^+ ion excretion rates, which were not altered by short-term or prolonged exposure to the extract. The same research group reported that the ethanolic stem bark extract at the same dosages improved blood glucose, the glomerular filtration rate, and mean arterial blood pressure in an STZ-induced T1D model of rats [122].

Sutherlandia frutescens R. BR., variety incana E. MEY., (Fabaceae) is among the most common and widely used plants in the southern part of Africa for the treatment of DM and its associated complications. The hypoglycemic effect of *S. frutescens* has been investigated [123]. The shoots aqueous extract significantly prevented the STZ-induced hyperglycemic condition in mice when administered orally at various dosages (50–800 mg/kg bw). In another study, Chadwick et al. [124] reported that *S. frutescens* is a potential agent in the management of DM, especially type 2 DM. The aqueous leaf extract-treated rats showed an increase in glucose uptake and utilization by peripheral tissues with a decrease in intestinal glucose absorption. Glucose uptake was carried out using [^3H] deoxy-glucose. The [^3H] deoxy-glucose count increased significantly in muscle and epididymal fat tissues when treated with the *S. frutescens* shoot aqueous extract in a diet-induced T2D model of rats, and the results were comparable with metformin-treated rats. In a different study, *S. frutescens* prevented insulin resistance and showed a hypolipidemic effect in diabetic rats [125]. The plasma free fatty acid dropped significantly. Similarly, the homeostatic model assessment (HOMA-IR) and quantitative insulin sensitivity check index (QUICKI) demonstrated that oral treatment of the extract at 50 mg/kg bw prevented the development of insulin resistance in a high-fat diet-fed insulin-resistance model of rats.

Central Africa

Bersama engleriana (Melianthaceae) is commonly available in almost all African regions and has been traditionally used in the management of DM. To evaluate such a claim, preliminary hypoglycemic effects of both aqueous and methanolic leaf extracts at 300 and 600 mg/kg bw/day were reported in normoglycemic rats

[126]. At the 600 mg dose, blood glucose levels dropped by 37.7% and 49.11% for aqueous and methanolic extracts, respectively. To expand upon this study, Watcho et al. [127] investigated the effect of this plant on STZ/nicotinamide T2D models of rats. The leaf aqueous and methanolic extracts treated orally at 400 and 600 mg/kg bw/day significantly and dose-dependently decreased the blood glucose levels and lipid profile with an increase in HDL cholesterol levels. At the 600 mg dose, the ethanolic extract demonstrated a higher reduction of blood glucose levels (80.31%) compared to the aqueous extract (67.74%). The decrease in organ weight recorded in diabetic-untreated rats was completely prevented in the extract-treated groups.

Dichrostachys glomerata Chiov. (Cucurbitaceae) is a spice used in the management of DM and its associated complications. Because of its strong antioxidant action, the effect of *D. glomerata* on various cardiovascular disease risk factors in obese normoglycemic and obese type 2 diabetic human subjects has been reported [128]. Dried pods were supplied daily in the form of capsules containing 400 mg *D. glomerata* 30–60 minutes before lunch and dinner. The results of the study indicated a decrease in body weight by 7.91% in obese normoglycemic and 5.97% in obese type 2 diabetic subjects. Similarly, the reduction of BMI, waist and hip circumference, body fat, blood pressure, blood cholesterol, triglycerides, glucose, and glycosylated hemoglobin was higher in the normoglycemic subjects compared to the obese type 2 diabetic subjects. The results confirmed the traditional claims that *D. glomerata* could ameliorate the complications associated with DM and other related cardiovascular diseases.

Although rarely investigated, an important complication associated with DM in men is infertility or erectile dysfunction [129]. As a result, Wankeu-Nya et al. [130] investigated the possible antidiabetic effect of aqueous and ethanolic root bark extracts of *Dracena arborea* Wild. (Dracaenaceae), which has been widely acclaimed for its aphrodisiac action in Cameroonian traditional medicine. The antihyperglycemic effect was observed in an STZ-induced T1D model of rats, with no such effect on normoglycemic rats. Oral treatment at 500 mg/kg bw of aqueous extract and 100 mg/kg bw of ethanolic extract for three weeks ameliorated the severe damages of the testes morphology and spermatogenesis, as observed in the diabetic-untreated rats. Although an increase in blood glucose was observed in both the aqueous (15.39%) and ethanolic extract (19.04%) -treated animals, this increase was lower than that of the untreated animals ($>60.34\%$). Flavonoids, sterols, and saponins were some of the compounds qualitatively determined to be present in both extracts.

Kalanchoe crenata Andr. Haw. (Crassulaceae) is among the most widely used plants in Cameroon and other central African countries for therapeutic purposes. Kamgang et al. [131] have reported that the aqueous-ethanol extract significantly, but not dose-dependently, decreased the blood glucose levels of diet-induced type 2 diabetic rats within six hours and four weeks of oral treatment at 200 mg/kg bw/day. The percentage of decrease recorded after four weeks was 52%. The diabetic-treated rats also showed an improvement in insulin sensitivity, a decrease in body weight, and reduced water intake. In another study, Fondjo et al. [132] investigated the antidyslipidemic and antioxidant effects of *K. crenata* whole plant methanolic extract in an STZ-induced T1D model of rats. The extract treated orally at 50 and 68 mg/kg bw/day showed a decrease in serum, liver, and kidney malondialdehyde levels, with an increase in activities of antioxidant enzymes. The glycemic reduction in treated animals was 35% and 44% for 50 and 68 mg/kg bw, respectively, after the 6-week post-treat-

ment period. All diabetic-treated animals showed a decrease in lipid parameters, with an increase in HDL cholesterol levels and an overall reduction of the atherogenic index by 31%.

East Africa

Caylusea abyssinica Fresen. Fisch. & Mey. (Resedaceae) is popularly used in different East African countries for the management of DM, especially in Ethiopian folklore medicine. Tamiru et al. [133] reported the hypoglycemic effect of the methanolic leaf extract of *C. abyssinica* in a normal, glucose-loaded, and STZ-induced T1D model of rats when administered orally at 100, 200, and 300 mg/kg bw. In an oral glucose tolerance test, the extract-treated rats indicated a better glucose handling ability than the diabetic controls. A reduction of 52.2%, 62.3%, and 52.8% in glycemia was achieved at the fourth hour of treatment at 100, 200, and 300 mg/kg bw, respectively.

The hypoglycemic effect of five Kenyan medicinal plants in alloxanized mice has been reported [134]. These plants include *Strychnos henningsii* Gilg. (Loganiaceae), *Erythrina abyssinica* Lam. (Fabaceae), *Aspilia pluriseta* Schweinf. (Asteraceae), *Bidens pilosa* L. (Asteraceae), and *Catha edulis* (Vahl) Forssk. ex Endl. (Celastraceae). All the extracts showed a significant and dose-dependent blood glucose lowering activity within the 4-h post-treatment period at 50, 100, and 150 mg/kg bw. The hypoglycemic effect of *C. edulis* was much better compared to others when given at a dose of 150 mg/kg bw and was as effective as insulin. Polyphenols were the major active components detected in the various parts of the plants under this study. Based on the toxicity study conducted for various parts of these plants, at higher dosages, aqueous extracts from the stem bark of *E. abyssinica*, the root bark of *C. edulis*, and *B. pilosa* leaves were nephrotoxic as well as hepatotoxic. *S. henningsii* leaves were moderately toxic while *A. pluriseta* root bark was reported as safe during these studies.

Momordica charantia L. (Cucurbitaceae) is one of the plants commonly used as food and in therapeutic purposes by both diabetic and healthy people, and has been known in traditional medicine for its glucose-lowering action worldwide [135]. Its fruit has a distinguished bitter taste, which is more pronounced as it ripens; hence, it is named bitter melon. A study reported by Matheka et al. [136] demonstrated that the oral administration of fresh fruit juice extract of *M. charantia* by gastric gavage at 10 ml/kg bw decreased blood glucose levels significantly by about 30 and 10% after 30 and 90 minutes, respectively, in a T1D model of rats.

Moringa stenopetala Baker f. (Moringaceae) is acclaimed for its glucose-lowering ability in Ethiopian traditional medicine. Nardos et al. [137] reported the antidiabetic effect of the leaf extracts and fractions of *M. stenopetala* in an alloxan-induced T1D model of rats. The extracts and fractions intraperitoneally administered at 300 mg/kg bw decreased blood glucose levels significantly by nearly 20% after an 8-day post-treatment period. The ethanolic extract was safe up to 5 g/kg bw. Furthermore, Toma et al. [138] demonstrated both antihyperglycemic and antihyperlipidemic effects with a daily oral administration of butanol fraction from the leaf ethanolic extract of *M. stenopetala* for four weeks in an alloxan-induced T1D rat model.

Conclusion

Apart from the folkloric claims, it is evident from the above reviewed studies that Africa is blessed with an abundance of anti-diabetic plants resources based on scientific findings. However, due to the variations in the scientific investigations in terms of analyzed antidiabetic parameters, doses, and durations used, it is difficult to precisely identify the plant(s) with the best reported activity, but our close analysis of the reports seem to suggest that *O. gratissimum*, *A. occidentale*, *V. amygdalina*, *G. latifolium*, *A. indica*, *C. carvi*, *M. alba*, and *A. iva* are the most active because they received much attention as is evident by numerous studies and, thus, possibly contain the most bioactive antidiabetic phytochemicals among all the plants. The methods mostly utilized for the extractions of various parts via organic solvent extractions include maceration/cold extraction, soxhlet, distillation, percolation, and sequential extraction. Moreover, it is evident that very few studies were reported to involve human subjects. Most studies used either T1D or T2D animal models. Unfortunately, perhaps due to limited research resources, most of the studies are preliminary in nature (though with promising results) and do not include detailed isolation and characterization of the bioactive compounds and/or the mechanisms of antidiabetic actions. Government agencies and/or pharmaceutical industries should support more research activities in this area in order to commercially utilize these antidiabetic medicinal plants for a solution to the continent's myriad of economic problems.

Methodology

Relevant literatures were collected by searching the major scientific databases including Pubmed, ScienceDirect, Medline, and Google Scholar for medicinal plants of African origin that have been studied and investigated for their antidiabetic therapeutic potentials *in vivo*. Some articles were found through tracking citations from other publications or by directly accessing the journals' website. They were considered on the basis of the geographical region of their origin. The literature considered were those available covering the period January 2000 to July 2013. The keyword combinations for the search were antidiabetic, antihyperglycemia, hypoglycemia, medicinal plant, and Africa. Supplementary information was obtained by using another keyword combination such as plant, hypoglycemia, and Africa. A total of 313 articles were retrieved in this review, out of which 256 research articles that reported *in vivo*, and not *in vitro*, activity were selected and presented in this review. Following the search, the plants were categorized and presented based on their regional origins.

Acknowledgements

This study was supported by the Competitive Research Grant from the Research Office, University of KwaZulu-Natal (UKZN), Durban, an Incentive Grant for Rated Researchers, and Grant Support for Women and Young Researchers from the National Research Foundation (NRF), Pretoria, South Africa. The first author was awarded a PhD study scholarship by the Education Trust Fund desk office, Ahmadu Bello University, Zaria, Nigeria.

Conflict of Interest

There is no conflict of interest within this article.

Reference

- Neelesh M, Sanjay J, Sapna M. Antidiabetic potential of medicinal plants. *Acta Pol Pharm* 2010; 67: 113–118
- Maritim AC, Sander RA, Watkins JB. Diabetes, oxidative stress, and antioxidants: a review. *J Biochem Mol Toxicol* 2003; 17: 24–38
- Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. *Diabet Res Clin Pract* 2011; 94: 311–321
- International Diabetes Federation (Brussels). Diabetes at a glance, 2012, Africa (AFR). Available at https://www.idf.org/sites/default/files/IDF_AFR_5E_Update_FactSheet_0.pdf. Accessed February 13, 2013.
- Michael PK, Asim AB, Robert SB. The utility of oral diabetes medications in type 2 diabetes of the young. *Curr Diabet Rev* 2005; 1: 83–92
- Cragg GM, Newman DJ. Natural products: a continuing source of novel drug leads. *Biochim Biophys Acta* 2013; 1830: 3670–3695
- Zhang X. Legal status of traditional medicines and complementary/alternative medicine. A worldwide review. *World Health Organ Tech Rep* 2001; 2: 5–41
- Zhang BB, Moller DE. New approaches in the treatment of type 2 diabetes. *Curr Opin Chem Biol* 2000; 4: 461–467
- Calixto JB. Efficacy, safety, quality control, marketing and regulatory guidelines for herbal medicines (phytotherapeutic agents). *Braz J Med Biol Res* 2000; 33: 179–189
- Van Wyk BE, deWet H, Van Heerden FR. An ethnobotanical survey of medicinal plants in the south-eastern Karoo, South Africa. *S Afr J Bot* 2008; 74: 696–704
- Adebayo AG. Inventory of antidiabetic plants in selected districts of Lagos State, Nigeria. *J Ethnopharmacol* 2009; 121: 135–139
- N'guessa K, Kouassi K, Kouadio K. Ethnobotanical study of plants used to treat diabetes, in traditional medicine, by Abbey and Krobou people of Agboville (Côte-d'Ivoire). *Am J Sci Res* 2009; 4: 45–58
- Keter LK, Mutiso PC. Ethnobotanical studies of medicinal plants used by traditional health practitioners in the management of diabetes in lower eastern province, Kenya. *J Ethnopharmacol* 2012; 139: 74–80
- Moshi MJ, Mbwambo ZH. Experience of Tanzanian traditional healers in the management of non-insulin dependent diabetes mellitus. *Pharm Biol* 2002; 40: 552–560
- Rutebemberwa E, Lubega M, Katureebe SK, Oundo A, Kiweewa F, Mukanga D. Use of traditional medicine for the treatment of diabetes in Eastern Uganda: a qualitative exploration of reasons for choice. *BMC Int Health Hum Rights* 2013; 13: 2–7
- Rachid A, Rabah D, Farid L, Zohra SF, Houcine B, Nacéra B. Ethnopharmacological survey of medicinal plants used in the traditional treatment of diabetes mellitus in the North Western and South Western Algeria. *J Med Plants Res* 2012; 6: 2041–2050
- Bnouham M, Mekhfi H, Legssyer A, Ziyat A. Ethnopharmacology forum medicinal plants used in the treatment of diabetes in Morocco. *Int J Diabet Metab* 2002; 10: 33–50
- Eddouks M, Maghrani M, Lemhadri A, Ouahidi ML, Jouad H. Ethnopharmacological survey of medicinal plants used for the treatment of diabetes mellitus, hypertension and cardiac diseases in the south-east region of Morocco (Tafilalet). *J Ethnopharmacol* 2002; 82: 97–103
- Afolayan AJ, Sunmonu TO. *In vivo* Studies on Antidiabetic Plants Used in South African Herbal Medicine. *J Clin Biochem Nutr* 2010; 47: 98–106
- Mahop TM, Mayet M. Enroute to biopiracy? Ethnobotanical research on antidiabetic medicinal plants in the Eastern Cape Province, South Africa. *Afr J Biotechnol* 2007; 6: 2945–2952
- Din N, Dibong SD, Mpondo E, Priso RJ, Kwin NF, Ngoye A. Inventory and identification of plants used in the treatment of diabetes in Douala Town (Cameroon). *Eur J Med Plants* 2011; 1: 60–73
- Olatunji LA, Okwusidi JI, Soladoye AO. Antidiabetic effect of *Anacardium occidentale* stem-bark in fructose-diabetic rats. *Pharm Biol* 2005; 43: 589–593
- Abdullahi S, Olatunji GA. Antidiabetic activity of *Anacardium occidentale* in alloxan – diabetic rats. *J Sci Technol* 2010; 30: 35–41
- Fagbohun TR, Odunfunwa KT. Hypoglycemic effect of methanolic extract of *Anacardium occidentale* leaves in alloxan-induced diabetic rats. *Niger J Physiol Sci* 2010; 25: 87–90
- Ukwenya VO, Ashaolu JO, Adeyemi DO, Akinola OB, Caxton-Martins EA. Antihyperglycemic activities of methanolic leaf extract of *Anacardium occidentale* (Linn.) on the pancreas of streptozotocin-induced diabetic rats. *J Cell Anim Biol* 2012; 6: 207–212
- Koffour GA, Amoatengor P, Okai CA, Fiagbe NIY. Hypoglycemic effects of whole and fractionated *Azadirachta indica* (Neem) seed oils on alloxan-induced diabetes in New Zealand white rabbits. *J Ghana Sci Assoc* 2011; 13: 34–39
- Henry DA, Itemobong SE, Ito FU, Patrick EE, Isong NB. Effect of aqueous extract of *Azadirachta indica* (Neem) leaves on some indices of pancreatic function in alloxan-induced diabetic Wistar rats. *Pharmacologia* 2012; 3: 420–425
- Akinola OB, Zatta L, Dosumu OO, Akinola OS, Adelaja AA, Dini L, Caxton-Martins EA. Intestinal lesions of streptozotocin-induced diabetes and the effects of *Azadirachta indica* treatment. *Pharmacology* 2009; 3: 872–881
- Akinola OB, Caxton-Martins EA, Dini L. Chronic treatment with ethanolic extract of the leaves of *Azadirachta indica* ameliorates lesions of pancreatic islets in streptozotocin diabetes. *Int J Morphol* 2010; 28: 291–302
- Akinola OB, Zatta L, Dosumu OO, Akinola OS, Dini L, Caxton-Martins EA. Ameliorative effects of ethanolic leaves extract of *Azadirachta indica* on renal histologic alterations streptozotocin-induced diabetic rats. *Am J Chin Med* 2011; 39: 903–916
- Itemobong SE, Item JA, Henry DA, Ito FU, Oboso EE, Patrick EE. Effects of ethanol extract of *Azadirachta indica* leaves on some immunological and haematological parameters of diabetic Wistar rats. *Afr J Pharm Pharmacol* 2010; 4: 104–108
- Nwanjo HU, Okafor MC, Oze GO. Anti-lipid peroxidative activity of *Gongronema latifolium* in streptozotocin-induced diabetic rats. *Niger J Physiol Sci* 2006; 21: 61–65
- Johnkennedy N, Adamma E, Augustin I, Ifeoma UH. Influence of *Gongronema latifolium* on some biochemical parameters in alloxan induced diabetes. *Int J Anal Pharm Biomed Sci* 2012; 1: 13–17
- Ibegbulem CO, Chikezie PC. Hypoglycemic properties of ethanolic extracts of *Gongronema latifolium*, *Aloe perryi*, *Viscum album* and *Allium sativum* administered to alloxan induced diabetic albino rats (*Rattus norvegicus*). *J Biol Chem Res* 2012; 29: 16–25
- Francis VU, Grace AE, Jones OA, Frank EE. Hypoglycemic effect of *Gongronema latifolium* extracts in rats. *J Nat Sci Res* 2013; 3: 37–44
- Akah PA, Uzodinma SU, Okolo CE. Antidiabetic activity of aqueous and methanol extract and fractions of *Gongronema latifolium* (Asclepiadaceae) leaves in alloxan diabetic rats. *J Appl Pharm Sci* 2011; 1: 99–102
- Edet EE, Akpanabiatu MI, Eno AE, Umoh IB, Itam EH. Effect of *Gongronema latifolium* crude leaf extract on some cardiac enzymes of alloxan-induced diabetic rats. *Afr J Biochem Res* 2009; 3: 366–369
- Adebajo AC, Ayoola MD, Odediran SA, Aladesanmi AJ, Schmidt TJ, Verspohl EJ. Evaluation of ethnomedical claim III: anti-hyperglycaemic activities of *Gongronema latifolium* root and stem. *J Diabet* 2013; 5: 336–343
- Orok UE, Eneji EG, Luke OU, Eyo RA, Sampson EV, Iwara IA, Oko OM. Effect of ethanolic root and twig extracts of *Gongronema latifolium* (uta-zi) on kidney function of streptozotocin induced hyperglycemic and normal Wistar rats. *J Med Med Sci* 2012; 3: 291–296
- Sini JM, Umar IA, Inuwa HM. The beneficial effect of extract of *Hibiscus sabdariffa* calyces in alloxan-diabetic rats: Hypoglycaemic and hypolipidaemic activities. *J Med Plants Res* 2011; 5: 2182–2186
- Sini JM, Umar IA, Inuwa HM. The beneficial effect of extract of *Hibiscus sabdariffa* calyces in alloxan-diabetic rats: Reduction of free radical load and enhancement of antioxidant status. *J Pharmacogn Phytother* 2011; 3: 141–149
- Adedayo OA, Ganiyu O. Aqueous extracts of roselle (*Hibiscus sabdariffa* Linn.) varieties inhibit α -amylase and α -glucosidase activities *in vitro*. *J Med Food* 2013; 16: 88–93
- Tanko Y, Abdelaziz MM, Adelaie AB, Fatihu MY, Musa KY. Effects of hydromethanolic leaves extract of *Indigofera pulchra* on blood glucose levels of normoglycemic and alloxan-induced diabetic Wistar rats. *Int J Appl Res Nat Prod* 2009; 1: 13–18
- Tanko Y, Hayatu Z, Mohammed A, Goji ADT, Musa KY, Yerima M. Effect of Residual Aqueous Portion of Hydro-methanolic Leaves Extract of *Indigofera pulchra* on blood glucose Levels of alloxan-induced diabetic Wistar rats. *Int J Anim Vet Adv* 2009; 1: 18–21
- Tanko Y, Abdelaziz MM, Adelaie AB, Fatihu MY, Musa KY. Effects of ethyl acetate portion of *Indigofera pulchra* leaves extract on blood glucose levels of alloxan-induced diabetic and normoglycemic Wistar rats. *Asian J Med Sci* 2009; 1: 10–14

- 46 Tanko Y, Mabrouk MA, Adelaiye AB, Fatihu MY, Musa KY. Anti-diabetic and some haematological effects of ethylacetate and n-butanol fractions of *Indigofera pulchra* extract on alloxan-induced diabetic Wistar rats. *J Diabet Endocrinol* 2011; 2: 1–7
- 47 Gidado A, Ameh DA, Atawodi SE. Effect of *Nauclea latifolia* leaves aqueous extracts on blood glucose levels of normal and alloxan-induced diabetic rats. *Afr J Biotechnol* 2005; 4: 91–93
- 48 Gidado A, Ameh DA, Atawodi SE, Ibrahim S. Hypoglycaemic activity of *Nauclea latifolia* SM. (Rubiaceae) in experimental animals. *Afr J Tradit Complement Alternat Med* 2008; 5: 201–208
- 49 Gidado A, Ameh DA, Atawodi SE, Ibrahim S. A preliminary study of the mechanism of hypoglycaemic activity of *Nauclea latifolia* leaf ethanolic extract. *J Complement Integr Med* 2012; DOI: 10.1515/1553–3840.1561
- 50 Yessoufou A, Gbenou J, Grissa O, Hitchami A, Simonin A, Tabka Z, Moudachirou M, Mountairou K, Khan N. Anti-hyperglycemic effects of three medicinal plants in diabetic pregnancy: modulation of T cell proliferation. *BMC Complement Alternat Med* 2013; 13: 77
- 51 Grace SE, Bob IAM, Godwin OI, Justin IA, Eyong UE, Patrick EE. Antioxidant enzymes activity and hormonal changes following administration of ethanolic leaves extracts of *Nauclea latifolia* and *Gongronema latifolium* in streptozotocin induced-diabetic rats. *Eur J Med Plants* 2013; 3: 297–309
- 52 Seung-Joo-Lee KU, Takayuki S, Kwang-Guen L. Identification of volatile components in basil (*Ocimum basilicum*) and thyme leaves (*Thymus vulgaris*) and their antioxidant properties. *Food Chem* 2004; 91: 131–137
- 53 Aguiyi JC, Obi CI, Gang SS, Igweh AC. Hypoglycaemic activity of *Ocimum gratissimum* in rats. *Fitoterapia* 2000; 71: 444–446
- 54 Onaolapo AY, Onaolapo OJ, Adewole SA. Ethanolic extract of *Ocimum gratissimum* leaves (Linn.) rapidly lowers blood glucose levels in diabetic Wistar rats. *Macedonia J Med Sci* 2011; 4: 351–357
- 55 Egesie UG, Adelaiye AB, Ibu JO, Egesie OJ. Safety and hypoglycaemic properties of aqueous leaf extract of *Ocimum gratissimum* in streptozotocin induced diabetic rats. *Niger J Physiol Sci* 2006; 21: 31–35
- 56 Mohammed A, Tanko Y, Okasha MA, Magaji RA, Yaro AH. Effects of aqueous leaves extract of *Ocimum gratissimum* on blood glucose levels of streptozotocin-induced diabetic Wistar rats. *Afr J Biotechnol* 2007; 6: 2087–2090
- 57 Asuquo OR, Edet AG, Mesembe O, Atanghwo J. Ethanolic extracts of *Vernonia amygdalina* and *Ocimum gratissimum* enhance testicular improvement in diabetic wistar rats. *Internet J Altern Med* 2010; DOI: 10.5580/89d
- 58 Joyce OO, Nworah DC, Ekwurugwu JN, Pughikumo DT, Joffa PPK. Comparative anti-diabetic effects of *Ocimum gratissimum*, *Vernonia amygdalina* and insulin on testicular architecture in STZ-induced diabetic rats. *Am J Pharm Tech Res* 2012; 2: 279–287
- 59 Oguanobi NI, Chijioko CP, Ghasi S. Anti-diabetic effect of crude leaf extracts of *Ocimum gratissimum* in neonatal streptozotocin-induced type-2 model diabetic rats. *Int J Pharm Pharm Sci* 2012; 5: 77
- 60 Okon UA, Ikpi DE, Ben EE. *Ocimum gratissimum* alleviates derangements in serum and biliary bilirubin, cholesterol and electrolytes in streptozotocin induced diabetic rats. *Int J Biochem Res Rev* 2013; 3: 171–189
- 61 Herbert OCM, Clement J, Idongesit J, Godwin E, Udemé E, Grace E. Evaluation of the hypoglycemic effect of aqueous extract of *Phyllanthus amarus* in alloxan-induced diabetic albino rats. *Int J Pharm Biomed Res* 2011; 2: 158–160
- 62 Owolabi OA, James DB, Anigo KM, Iormanger GW, Olaiya II. Combined effect of aqueous extracts of *Phyllanthus amarus* and *Vitex doniana* stem bark on blood glucose of streptozotocin (STZ) induced diabetes rats and some liver biochemical parameters. *Br J Pharmacol Toxicol* 2011; 2: 143–147
- 63 Povi L, Kwashie E, Amegnona A, Kodjo A, Edmond C, Messanvi G. Antidiabetic activity of *Phyllanthus amarus* Schum and Thonn (Euphorbiaceae) on alloxan induced diabetes in male Wistar rats. *J Appl Sci* 2011; 11: 2968–2973
- 64 Adeneye AA. The leaf and seed aqueous extract of *Phyllanthus amarus* improves insulin resistance diabetes in experimental animal studies. *J Ethnopharmacol* 2012; 144: 705–711
- 65 Aguwa CN, Ukwé CV, Inya-Agha SI, Okonta JM. Antidiabetic effect of *Picralima nitida* aqueous seed extract in experimental rabbit model. *J Nat Rem* 2001; 1: 135–139
- 66 Igboasoiyi AC, Essien EE, Eseyin OA, Ubam G. Screening of the seed of *Picralima nitida* for hypoglycaemic activity. *Pak J Biol Sci* 2007; 10: 828–830
- 67 Nimenibo-Uadia R. Effects of *Vernonia amygdalina* in alloxan-induced diabetic albino rats. *J Med Lab Sci* 2003; 12: 25–31
- 68 Chike CPR, Georgewill OA, Nnodi CU. Effect of aqueous leaf extract of *Vernonia amygdalina* (bitter leaf) on blood glucose concentration of alloxan induced diabetic albino Wistar rats. *Afr J Appl Zool Environ Biol* 2006; 8: 44–47
- 69 Osinubi AA. Effects of *Vernonia amygdalina* and chlorpropamide on blood glucose. *Med J Islamic World Acad Sci* 2007; 16: 115–119
- 70 Iwuji SC, Nwafor A, Adienbo OM, Ekwurugwu J, Iwuji NG, Oodo OM. Hypoglycaemic potential of aqueous leaf extract of *Vernonia amygdalina*: An Animal Model. *Afr J Med Physiol Biomed Eng Sci* 2010; 2: 9–13
- 71 Taiwo IA, Odeigah PGC, Ogunkanmi LA. The glycaemic effects of *Vernonia amygdalina* and *V. tenoreana* with tolbutamide in rats and the implications for the treatment of diabetes mellitus. *J Sci Res Dev* 2008; 11: 122–130
- 72 Fasola TR, Okeocha PC, Odetola A. Screening for the hypoglycaemic potentials of the extract of *Vernonia amygdalina*. *Ethnobot Leaflets* 2010; 14: 759–765
- 73 Owen OJ, Amakiri AO, Karibi-Botoye TA. Sugar-lowering effects of bitter leaf (*Vernonia amygdalina*) in experimental broiler finisher chickens. *Asian J Pharm Clin Res* 2011; 4: 19–21
- 74 Modu SA, Adeboye E, Maisaratu A, Mubi BM. Studies on the administration of *Vernonia amygdalina* Del. (Bitter leaf) and glucophage on blood glucose level of alloxan-induced diabetic rats. *Int J Med Plant Altern Med* 2013; 1: 13–19
- 75 Okolie UV, Okeke CE, Oli JM, Ehiemere JJ. Hypoglycemic indices of *Vernonia amygdalina* on postprandial blood glucose concentration of healthy humans. *Afr J Biotechnol* 2008; 7: 4581–4585
- 76 Atangwho IJ, Ebong PE, Eteng MU, Eyong EU, Obi AU. Effect of *Vernonia amygdalina* Del leaf on kidney function of diabetic rats. *Int J Pharmacol* 2007; 3: 143–148
- 77 Eteng MU, Bassey BJ, Atangwho IJ, Egbung GE, Eyoung EU, Ebong PE, Abolaji AO. Biochemical indices of macrovascular complication in diabetic rat model: compared effects of *Vernonia amygdalina*, *Cathartus roseus* and chlorpropamide. *Asian J Biochem* 2008; 3: 228–234
- 78 Akah PA, Alemji JA, Salawu OA, Okoye TC, Offiah NV. Effects of *Vernonia amygdalina* on biochemical and hematological parameters in diabetic rats. *Asian J Med Sci* 2009; 1: 108–113
- 79 Atangwho IJ, Eyong EU, Ebong PE, Egbung GE. Effects of combined administration of *Vernonia amygdalina* (Del.) and *Azadirachta indica* (A. Juss) extracts on compartmental glucose concentration and α -amylase activity of diabetic and non-diabetic rats. *Niger J Biochem Mol Biol* 2009; 24: 22–28
- 80 Atangwho IJ, Ebong PE, Eyong EU, Ashmawi MZ, Ahmad M. Synergistic antidiabetic activity of *Vernonia amygdalina* and *Azadirachta indica*: biochemical effects and possible mechanism. *J Ethnopharmacol* 2012; 141: 878–887
- 81 Akpaso MI, Atangwho IJ, Akpantah A, Fischer VA, Igiri AO, Ebong PE. Effect of combined leaf extracts of *Vernonia amygdalina* (Bitter leaf) and *Gongronema latifolium* (Utazi) on the pancreatic β -Cells of streptozotocin induced diabetic rats. *Br J Med Med Res* 2011; 1: 24–34
- 82 Okokon JE, Umoh UF, Ekpo BAJ, Etim EI. Antidiabetic study of combined extracts of *Vernonia amygdalina*, *Ocimum gratissimum*, and *Gongronema latifolium* on alloxan-induced diabetic rats. *J Nat Pharm* 2013; 4: 28–31
- 83 Kalejaiye OF, Iwalewa EO, Omobuwajo OR, Oyedapo OO. Hypoglycaemic effects of Nigerian *Zingiber officinale* rhizome on experimental diabetic rats. *Niger J Nat Prod Med* 2002; 6: 32–35
- 84 Ozougwu JC, Eyo JE. Evaluation of the activity of *Zingiber officinale* (ginger) aqueous extracts on alloxan-induced diabetic rats. *Pharmacology* 2011; 1: 258–269
- 85 Eyo JE, Ozougwu JC, Echi PC. Hypoglycaemic effects of *Allium cepa*, *Allium sativum* and *Zingiber officinale* aqueous extracts on alloxan-induced diabetic *Rattus norvegicus*. *Med J Islamic World Acad Sci* 2011; 19: 121–126
- 86 Iranloye BO, Arikawe AP, Rotimi G, Sogbade AO. Anti-diabetic and antioxidant effects of *Zingiber Officinale* on alloxan-induced and insulin-resistant diabetic male rats. *Niger J Physiol Sci* 2011; 26: 89–96
- 87 Morakinyo AO, Akindele AJ, Ahmed Z. Modulation of antioxidant enzymes and inflammatory cytokines: possible mechanism of anti-diabetic effect of ginger extracts. *Afr J Biomed Res* 2011; 14: 195–202
- 88 Arikawe AP, Daramola AO, Olatunji-Bello II, Obika LF. Insulin, pioglitazone and *Zingiber officinale* administrations improve proliferating cell nuclear antigen immunostaining effects on diabetic and insulin resistant rat testis. *J Exp Clin Med* 2013; 30: 49–55

- 89 Hilaly JE, Lyoussi B. Hypoglycaemic effect of the lyophilised aqueous extract of *Ajuga iva* in normal and streptozotocin diabetic rats. *J Ethnopharmacol* 2002; 80: 109–113
- 90 Hilaly JE, Tahraoui A, Israili ZH, Lyoussi B. Hypolipidemic effects of acute and sub-chronic administration of an aqueous extract of *Ajuga iva* L. whole plant in normal and diabetic rats. *J Ethnopharmacol* 2006; 105: 441–448
- 91 Hilaly JE, Tahraoui A, Israili ZH, Lyoussi B. Acute hypoglycemic, hypocholesterolemic and hypotriglyceridemic effects of continuous intravenous infusion of a lyophilised aqueous extract of *Ajuga iva* L. Schreber whole plant in streptozotocin-induced diabetic rats. *Pak J Pharm Sci* 2007; 20: 261–268
- 92 Taleb-Senouci D, Ghomari H, Krouf D, Bouderbala S, Prost J, Lacaille-Dubois MA, Bouchenak M. Antioxidant effect of *Ajuga iva* aqueous extract in streptozotocin-induced diabetic rats. *Phytomedicine* 2009; 16: 623–631
- 93 Hamden K, Ayadi F, Jamoussi K, Masuodi H, Elfeki A. Therapeutic effect of phytoecdysteroids rich extract from *Ajuga iva* on alloxan induced diabetic rats liver, kidney and pancreas. *Biofactors* 2008; 33: 165–175
- 94 Taj Eldin IM, Ahmed EM, Abd Elwahab HM. Preliminary study of the clinical hypoglycemic effects of *Allium cepa* (Red Onion) in type 1 and type 2 diabetic patients. *Env Health Insights* 2010; 4: 71–77
- 95 El-Demerdash FM, Yousef MI, Abou-El-Naga NI. Biochemical study on the hypoglycemic effects of onion and garlic in alloxan-induced diabetic rats. *Food Chem Toxicol* 2005; 43: 57–63
- 96 El-Soud NA, Khalil M. Antioxidative effects of *Allium cepa* essential oil in streptozotocin induced diabetic rats. *Macedonia J Med Sci* 2010; 3: 344–351
- 97 Samir AMZ, Somaia ZAR, Mattar AF. Anti-diabetic properties of water and ethanolic extracts of *Balanites aegyptiaca* fruits flesh in senile diabetic rats. *Egypt J Hosp Med* 2003; 10: 90–108
- 98 Gad MZ, El-Sawalhi MM, Ismail MF, El-Tanbouly ND. Biochemical study of the anti-diabetic action of the Egyptian plants fenugreek and balanites. *Mol Cell Biochem* 2006; 281: 173–183
- 99 Singh G, Kapoor IP, Pandey SK, Singh UK, Singh RK. Studies on essential oils. Part 10. Antibacterial activity of volatile oils of some spices. *Phytother Res* 2002; 16: 680–682
- 100 Eddouks M, Lemhadri A, Michel JB. Caraway and caper: potential anti-hyperglycaemic plants in diabetic rats. *J Ethnopharmacol* 2004; 94: 143–148
- 101 Lemhadri A, Eddouks M, Michel JB. Cholesterol and triglycerides lowering activities of caraway fruits in normal and streptozotocin diabetic rats. *J Ethnopharmacol* 2006; 106: 321–326
- 102 Dawidar AM, Abdel-Mogib M, Abou-Elzahab MM, Berghot MA, Mahfouz M, El-Ghorab AH, Hussien KH. Effect of photo-oxygenation on biological activities of some commercial Egyptian essential oils. *Rev Latin-am Quim* 2010; 38: 168–179
- 103 Lemhadri A, Eddouks M, Sulpice T, Burcelin R. Anti-hyperglycaemic and anti-obesity effects of *Capparis spinosa* and *Chamaemelum nobile* aqueous extracts in HFD mice. *Am J Pharmacol Toxicol* 2007; 2: 106–110
- 104 Lemhadri A, Burcelin L, Sulpice T, Eddouks M. *Chamaemelum nobile* L. aqueous extract represses endogenous glucose production and improves insulin sensitivity in streptozotocin-induced diabetic mice. *Am J Pharmacol Toxicol* 2007; 2: 116–122
- 105 Singab ANB, El-Beshbishy HA, Yonekawa M, Nomura T, Fukai T. Hypoglycemic effect of Egyptian *Morus alba* root bark extract: effect on diabetes and lipid peroxidation of streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2005; 100: 333–338
- 106 El-Sayyad HHH, El-Sherbiny MA, Sobh MA, Abou-El-Naga AM, Ibrahim MAN, Mousa SA. Protective effects of *Morus alba* leaves extract on ocular functions of pups from diabetic and hypercholesterolemic mother rats. *Int J Biol Sci* 2011; 7: 715–728
- 107 El-Beshbishy HA, Singab AN, Sinkkonen J, Pihlaja K. Hypolipidemic and antioxidant effects of *Morus alba* L. (Egyptian mulberry) root bark fractions supplementation in cholesterol-fed rats. *Life Sci* 2006; 78: 2724–2733
- 108 Mathur ML, Gaur J, Sharma R, Haldiya KR. Antidiabetic properties of a spice plant *Nigella sativa*. *J Endocrinol Metab* 2011; 1: 1–8
- 109 El-Dakhakhny M, Mady N, Lambert N, Ammon HP. The hypoglycemic effect of *Nigella sativa* oil is mediated by extrapancreatic actions. *Planta Med* 2002; 68: 465–466
- 110 Mohamed AM, EL-Sharkawy FZ, Ahmed SAA, Aziz WM, Badary OA. Glycemic control and therapeutic effect of *Nigella sativa* and *Curcuma longa* on rats with streptozotocin-induced diabetic hepatopathy. *J Pharmacol Toxicol* 2009; 4: 45–57
- 111 Benhaddou-Andaloussi A, Martineau L, Vuong T, Meddah B, Madiraju P, Settaf A, Haddad PS. The *in vivo* antidiabetic activity of *Nigella sativa* is mediated through activation of the AMPK pathway and increased muscle Glut4 content. *Evid Based Complement Alternat Med* 2011; DOI: 10.1155/2011/538671
- 112 Abdel-Zahera AO, Salima SY, Assaf MH, Abdel-Hady RH. Antidiabetic activity and toxicity of *Ziziphus spina-christi* leaves. *J Ethnopharmacol* 2005; 101: 129–138
- 113 Hala MH, Eman ME, Aataa AS. Antihyperglycemic, antihyperlipidemic and antioxidant effects of *Ziziphus spina-christi* and *Ziziphus jujuba* in alloxan induced diabetic rats. *Int J Pharmacol* 2006; 2: 563–570
- 114 Michel CG, Nesseem DI, Ismail MF. Anti-diabetic activity and stability study of the formulated leaf extract of *Ziziphus spina-christi* (L.) Willd with the influence of seasonal variation. *J Ethnopharmacol* 2011; 133: 53–62
- 115 Afolayan AJ, Sunmonu TO. *Artemisia afra* Jacq. ameliorates oxidative stress in the pancreas of streptozotocin-induced diabetic Wistar rats. *Biosci Biotechnol Biochem* 2011; 75: 2083–2086
- 116 Afolayan AJ, Sunmonu TO. Protective role of *Artemisia afra* aqueous extract on tissue antioxidant defense system in streptozotocin-induced diabetic rats. *Afr J Tradit Complement Alternat Med* 2013; 10: 15–20
- 117 Ojewale JAO. Antinociceptive, anti-inflammatory and antidiabetic properties of *Bryophyllum pinnatum* (Crassulaceae) leaf aqueous extract. *J Ethnopharmacol* 2005; 99: 13–19
- 118 Mpiana PT, Masunda TA, Longoma BF, Tshibangu DST, Ngbolua KN. Anti-hyperglycemic activity of *Raphia gentiliana* De Wild. (Arecaceae). *Eur J Med Plants* 2013; 3: 233–240
- 119 Ojewale JAO. Hypoglycemic effect of *Sclerocarya birrea* [(A. Rich.) Hochst.] [Anacardiaceae] stem-bark aqueous extract in rats. *Phytomedicine* 2003; 10: 675–681
- 120 Ojewale JAO. Evaluation of the analgesic, anti-inflammatory and antidiabetic properties of *Sclerocarya birrea* (A. Rich.) Hochst. stem-bark aqueous extract in mice and rats. *Phytother Res* 2004; 18: 601–608
- 121 Musabayane CT, Gondwe M, Kamadyaapa DR, Moodley K, Ojewale JAO. The effects of *Sclerocarya birrea* [(a. rich.) hochst.] [Anacardiaceae] stem-bark aqueous extract on blood glucose, kidney and cardiovascular function in rats. *Endocr Abstr* 2006; 12: 36
- 122 Gondwe M, Kamadyaapa DR, Tufts M, Chuturgoon AA, Musabayane CT. *Sclerocarya birrea* [(A. Rich.) Hochst.] [Anacardiaceae] stem-bark ethanolic extract (SBE) modulates blood glucose, glomerular filtration rate (GFR) and mean arterial blood pressure (MAP) of STZ-induced diabetic rats. *Phytomedicine* 2008; 15: 699–709
- 123 Ojewale JAO. Analgesic, anti-inflammatory and hypoglycemic effects of *Sutherlandia frutescens* R.Br. (variety *incana* E. MEY.) [Fabaceae] shoot aqueous extract. *Methods Find Exp Clin Pharmacol* 2004; 26: 409–416
- 124 Chadwick WA, Roux S, Van de Venter M, Louw J, Oelofsen W. Anti-diabetic effects of *Sutherlandia frutescens* in Wistar rats fed a diabetogenic diet. *J Ethnopharmacol* 2007; 109: 121–127
- 125 MacKenzie J, Koekemoer T, Van de Venter M, Dealtry G, Roux S. *Sutherlandia frutescens* limits the development of insulin resistance by decreasing plasma free fatty acid levels. *Phytother Res* 2009; 23: 1609–1614
- 126 Nyah NG, Watcho P, Nguetefack T, Kamanyi A. Hypoglycaemic activity of the leaves extracts of *Bersama engleriana* in rats. *Afr J Tradit Complement Alternat Med* 2007; 2: 215–221
- 127 Watcho P, Anchountsa JHG, Mbiakop CU, Nguetefack TB, Kamanyi A, Wankeu-Nya M. Hypoglycemic and hypolipidemic effects of *Bersama engleriana* leaves in nicotinamide/streptozotocin-induced type 2 diabetic rats. *BMC Complement Alternat Med* 2012; 12: 264
- 128 Kuete D, Etoundi BC, Ngondi JL, Oben E. Effects of *Dichrostachys glomerata* spice on cardiovascular diseases risk factors in normoglycemic and type 2 diabetic obese volunteers. *Food Res Int* 2011; 44: 1197–1202
- 129 Mallidis C, Agbaje I, McClure N, Kliesch S. The influence of diabetes mellitus on male reproductive function: a poorly investigated aspect of male infertility. *Urol Assoc* 2011; 50: 33–37
- 130 Wankeu-Nya M, Watcho P, Florea A, Balici S, Matei H, Kamanyi A. *Draena arborea* alleviates ultra-structural spermatogenic alterations in streptozotocin-induced diabetic rats. *BMC Complement Alternat Med* 2013; 13: 71

- 131 Kamgang R, Mboumi YR, Fondjo AF, Tagne MAF, Mengue N'dille GPR, Yonkeu JN. Antihyperglycaemic potential of the water-extract of *Kalanchoe crenata* (Crassulaceae). *J Nat Med* 2008; 62: 34–40
- 132 Fondjo FA, Kamgang R, Oyono JE, Yonkeu JN. Anti-dyslipidemic and antioxidant potentials of methanol extract of *Kalanchoe crenata* whole plant in streptozotocin-induced diabetic nephropathy in rats. *Trop J Pharm Res* 2012; 11: 767–775
- 133 Tamiru W, Engidawork E, Kaleab A. Evaluation of the effects of 80% methanolic leaf extract of *Caylusea abyssinica* (Fresen.) fisch. & Mey. on glucose handling in normal, glucose loaded and diabetic rodents. *BMC Complement Alternat Med* 2012; 12: 151
- 134 Piero NM, Murugi NJ, Kibiti MC, Ngeranwa JJ, Njue MW, Maina D, Gathumbi KP, Njagi NE. Hypoglycemic activity of some Kenyan plants traditionally used to manage diabetes mellitus in eastern province. *J Diabet Metab* 2011; 2: 1–6
- 135 Leung L, Birtwhistle R, Kotecha J. Anti-diabetic and hypoglycaemic effects of *Momordica charantia* (bitter melon): a mini review. *Br J Nutr* 2009; 102: 1703–1708
- 136 Matheka DM, Kiama TN, Alkizim FO, Bukachi F. Glucose-lowering effects of *Momordica charantia* in healthy rats. *Afr J Diabet Med* 2011; 19: 15–19
- 137 Nardos A, Makonnen E, Debella A. Effects of crude extracts and fractions of *Moringa stenopetala* (Baker f) Cufodontis leaves in normoglycemic and alloxan-induced diabetic mice. *Afr J Pharm Pharmacol* 2011; 5: 2220–2225
- 138 Toma A, Makonnen E, Debella A, Tesfaye B. Antihyperglycemic effect on chronic administration of butanol fraction of ethanol extract of *Moringa stenopetala* leaves in alloxan induced diabetic mice. *Asian Pac J Trop Biomed* 2012; 2: 606–610
- 139 Salisu Y, Agunu A, Abubakar MS, Ibrahim G. Hypoglycaemic effects of *Acacia albida* Del. (Mimosaceae) methanol root bark extract. *Niger J Pharm Sci* 2009; 8: 66–72
- 140 Ikewuchi J, Onyeike EN, Uwakwe AA, Ikewuchi CC. Effect of aqueous extract of the leaves of *Acalypha wilkesiana* 'Godseffiana' Muell. Arg. (Euphorbiaceae) on the hematology, plasma biochemistry and ocular indices of oxidative stress in alloxan induced diabetic rats. *J Ethnopharmacol* 2011; 137: 1415–1424
- 141 Tanko Y, Yerima M, Mahdi MA, Yaro AH, Musa KY, Mohammed A. Hypoglycemic activity of methanolic stem bark of *Adansonia digitata* extract on blood glucose levels of streptozotocin-induced diabetic Wistar rats. *Int J Appl Res Nat Prod* 2008; 1: 32–36
- 142 Odo RI, Asuzu IU, Aba PE. The antidiabetic activities of the methanolic root bark extract of *Afzelia africana* in alloxan-induced diabetic mice. *J Complement Integr Med* 2012; 9: 1–11
- 143 Egunyomi A, Gbadamosi IT, Animashahun MO. Hypoglycaemic activity of the ethanol extract of *Ageratum conyzoides* Linn. shoots on alloxan-induced diabetic rats. *J Med Plants Res* 2011; 5: 5347–5350
- 144 Agunbiade OS, Ojezele OM, Ojezele JO, Ajayi AY. Hypoglycaemic activity of *Commelina africana* and *Ageratum conyzoides* in relation to their mineral composition. *Afr Health Sci* 2012; 12: 198–203
- 145 Mohammed RK, Ibrahim S, Atawodi SE, Eze ED, Suleiman JB. Anti-diabetic and haematological effects of n-butanol fraction of *Alchornea cordifolia* leaf extract in streptozotocin-induced diabetic Wistar rats. *Glob J Med Plant Res* 2012; 1: 14–21
- 146 Ozougwu JC. Anti-diabetic effects of *Allium cepa* (onions) aqueous extracts on alloxan-induced diabetic *Rattus norvegicus*. *J Med Plants Res* 2011; 5: 1134–1139
- 147 Obi HI, Ilodigwe EE, Ajaghaku DE, Okonta JM. The antidiabetic activity of combine aqueous extracts of *Gongronema latifolium* (Benth) and *Allium cepa*. *J Pharm Biomed Sci* 2012; 19: 1–5
- 148 Yusuf UA, Adeeyo OA, Salawu EA, Enaibe BU, Omotoso OD. *Allium cepa* protects renal functions in diabetic rabbits. *World J Life Sci Med Res* 2012; 2: 86–90
- 149 Ogunmodede OS, Saalu LC, Ogunlade B, Akunna GG, Oyewopo AO. An evaluation of hypoglycemic, antioxidant and hepatoprotective potentials of Onion (*Allium cepa* L.) on alloxan-induced diabetic rabbits. *Int J Pharmacol* 2012; 8: 21–29
- 150 Ojo RJ, Memudu AE, Akintayo CO, Akpan IS. Preventive effect of *Allium sativum* on alloxan induced diabetic rat. *ARPN J Agric Biol Sci* 2012; 7: 609–612
- 151 Saidu AN, Mann A, Balogun S. The hypoglycemic effect of aqueous extract of the *Anacardium occidentale* Linn leaves grown in Nigeria on normoglycemic albino rats. *J Emerg Trends Eng Appl Sci* 2012; 3: 302–308
- 152 Sani D, Sanni S, Ngulde SI. Hypoglycaemic effect of aqueous stem extract of *Anisopus mannii* in normal rats. *Afr J Pharm Pharmacol* 2009; 3: 481–484
- 153 Manosroi J, Zaruwa MZ, Manosroi A. Potent hypoglycemic effect of Nigerian anti-diabetic medicinal plants. *J Complement Integr Med* 2011; DOI: 10.2202/1553–3840.1482
- 154 Olubomehin OO, Abo KA, Ajaiyeoba EO. Alpha-amylase inhibitory activity of two *Anthocleista* species and *in vivo* rat model anti-diabetic activities of *Anthocleista djalensis* extracts and fractions. *J Ethnopharmacol* 2013; 146: 811–814
- 155 Ibeh BO, Ezeaja M. Preliminary study of antidiabetic activity of the methanolic leaf extract of *Axonopus compressus* (P. Beauv) in alloxan-induced diabetic rats. *J Ethnopharmacol* 2011; 138: 713–716
- 156 Aguh BI, Nock IH, Ndams IS, Agunu A. Hypoglycaemic activity and Nephro-protective effect of *Bauhinia rufescens* in alloxan-induced diabetic rats. *Int J Appl Pharm Biol Chem* 2013; 2: 249–255
- 157 Aguh BI, Nock IH, Ndams IS, Agunu A, Ayeku PO. Hypolipidemic properties of *Bauhinia rufescens* in alloxan-induced diabetic rats. *Sci J Biol Sci* 2013; 2: 68–75
- 158 Kolawole OM, Oladoyinbo SO, Agbede OO, Adu FD. The effect of *Bridelia ferruginea* and *Senna alata* on plasma glucose concentration in normoglycemic and glucose induced hyperglycemic rats. *Ethnobot Leaflets* 2006; 10: 209–218
- 159 Adeneye AA, Olagunju AA. Preliminary hypoglycemic and hypolipidemic activities of the aqueous seed extract of *Carica papaya* Linn. in Wistar rats. *Biol Med* 2009; 1: 1–10
- 160 Osadolor HB, Ariyo II, Emokpae MA, Anukam KC. Hypoglycemic effect of unripe pawpaw on streptozotocin induced diabetic albino rats. *Res J Med Plants* 2011; 5: 90–94
- 161 Ene AC, Milala MA, Nwankwo EA. The effect of different doses of black caraway (*Carum carvi* L.) oil on the liver enzymes of alloxan-induced diabetic rats. *J Med Sci* 2006; 6: 994–998
- 162 Ene AC, Bukbuk DN, Ogunmola OO. The effect of different doses of black caraway (*Carum carvi* L.) oil on the serum creatinine of alloxan-induced diabetic rats. *J Med Sci* 2006; 6: 701–703
- 163 Ene AC, Nwankwo EA, Samdi LM. Alloxan-induced diabetic rats and the effects of black caraway (*Carum carvi* L.) oil on their body weight. *Res J Med Med Sci* 2007; 2: 48–52
- 164 Ene AC, Nwankwo EA, Samdi LM. Alloxan-induced diabetic rats and the effects of black caraway (*Carum carvi* L.) oil on their body weight. *J Pharmacol Toxicol* 2007; 3: 141–146
- 165 Nadro MS. Hypoglycaemic effects of *Cassia italica* leaf extracts on streptozotocin-induced diabetic albino rats. *J Res Biosci* 2009; 6: 38–41
- 166 Nadro MS, Onoagbe IO. Anti-hyperlipidaemic and antioxidant effect of aqueous and ethanolic extracts of *Cassia italica* leaves in streptozotocin-induced diabetes in rats. *J Med Plants Res* 2012; 6: 4675–4681
- 167 Emeke EJ, Clement UO. Comparative study of hypoglycemic and biochemical effects of *Catharantus roseus* (Linn.), apocyanaceae (Madagascar periwinkle) and chlorpropamide (diabenese) on alloxan induced diabetic rats. *Biokemistri* 2005; 7: 149–156
- 168 Ladeji O, Omekarah I, Solomon M. Hypoglycemic properties of aqueous bark extract of *Cela pentandra* in streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2002; 84: 139–142
- 169 N'guessan K, Amoikon KE, Tièbré MS, Kadja B, Zirihi GN. Effect of aqueous extract of *Chrysophyllum cainito* leaves on the glycaemia of diabetic rabbits. *Afr J Pharm Pharmacol* 2009; 3: 501–506
- 170 Ezekwesili CN, Ogbunugafor HA, Ezekwesili-Ofili JO. Anti-diabetic activity of aqueous extracts of *Vitex doniana* leaves and *Cinchona calisaya* bark in alloxan-induced diabetic rats. *Int J Trop Dis Health* 2012; 2: 290–300
- 171 Tanko Y, Yaro AH, Isa AI, Yerima M, Saleh MIA, Mohammed A. Toxicological and hypoglycemic studies on the leaves of *Cissampelos mucronata* (Menispermaceae) on blood glucose levels of streptozotocin-induced diabetic Wistar rats. *J Med Plants Res* 2007; 1: 113–116
- 172 Ekeanyanwu RC, Udeme AA, Onuigbo AO, Etienajirhevwe OF. Anti-diabetic effect of ethanol leaf extract of *Cissampelos owariensis* (lungwort) on alloxan induced diabetic rats. *Afr J Biotechnol* 2012; 11: 6758–6762
- 173 Opajobi AO, Esume CO, Campbell P, Onyesom I, Osasuyi A. Effects of aqueous extracts of *Rauvolfia vomitoria* and *Citrus aurantiensis* on liver enzymes of streptozotocin – induced diabetic and normal rabbits. *Cont J Med Res* 2011; 5: 1–5

- 174 Adeneye AA. Hypoglycemic and hypolipidemic effects of methanol seed extract of *Citrus paradisi* Macfad (Rutaceae) in alloxan-induced diabetic Wistar rats. *Niger Q J Hosp Med* 2008; 18: 211–215
- 175 Adebajo AC, Iwalewa EO, Obuotor EM, Ibikunle GF, Omisore NO, Adekunmi CO, Obaparusi OO, Klaes M, Adetogun GE, Schmidt TJ, Verspoh EJ. Pharmacological properties of the extract and some isolated compounds of *Clausena lansium* stem bark: anti-trichomonal, antidiabetic, anti-inflammatory, hepatoprotective and antioxidant effects. *J Ethnopharmacol* 2009; 122: 10–19
- 176 Adisa RA, Choudhary MI, Adewoye EO, Olorunsogo OO. Hypoglycaemic and biochemical properties of *Cnestis ferruginea*. *Afr J Tradit Complement Alternat Med* 2010; 7: 185–194
- 177 Chika A, Bello SO. Antihyperglycaemic activity of aqueous leaf extract of *Combretum micranthum* (Combretaceae) in normal and alloxan-induced diabetic rats. *J Ethnopharmacol* 2010; 129: 34–37
- 178 Sarah N, Oluwatosin A, Ajaiyeoba E. Oral administration of extract from *Curcumin longa* lowers blood glucose and attenuates alloxan-induced hyperlipidemia in diabetic rabbits. *Pak J Biol Sci* 2009; 8: 625–628
- 179 Iwueke AV, Nwodo OFC. Effect of combined root extract of *Sarcocephalus latifolius* and *Daniella oliveri* on blood glucose and serum lipids in alloxan-diabetic rats. *Niger J Biochem Mol Biol* 2008; 23: 37–41
- 180 Iwueke AV, Nwodo OFC. Antihyperglycaemic effect of aqueous extract of *Daniella oliveri* and *Sarcocephalus latifolius* roots on key carbohydrate metabolic enzymes and glycogen in experimental diabetes. *Biochemistri* 2008; 20: 63–70
- 181 Omoniwa BP, Luka CD. Antidiabetic and toxicity evaluation of aqueous stem extract of *Ficus asperifolia* in normal and alloxan-induced diabetic rats. *Asian J Exp Biol Sci* 2012; 3: 726–732
- 182 Adeyi AO, Idowu AB, Mafiana CF, Oluwalana SA, Ajayi OL. Effects of aqueous leave extract of *Ficus exasperata* on pathophysiology and histopathology of alloxan-induced diabetic albino rats. *J Med Plants Res* 2012; 6: 5730–5736
- 183 Adewole SO, Adenowo TK, Thajasvarie N, Ojewole AO. Hypoglycaemic and hypotensive effects of *Ficus exasperata* VAHL. (Moraceae) leaf aqueous extract in rats. *Afr J Tradit Complement Alternat Med* 2011; 8: 275–283
- 184 Mohammed A, Adelaiye AB, Bakari AG, Mabrouk MA. Anti-diabetic and some haematological effects of ethylacetate and n-butanol fractions of *Ganoderma lucidum* aqueous extract in alloxan-induced diabetic Wistar rats. *Int J Med Med Sci* 2009; 1: 530–535
- 185 Gnanoran BN, N'guessan BB, Amoateng P, Dosso K, Yapo AP, Ehile EE. Hypoglycaemic activity of ethanolic leaf extract and fractions of *Holarrhena floribunda* (Apocynaceae). *J Med Biomed Sci* 2012; 1: 46–54
- 186 Okokon JE, Bassey SA, Basil NI. Antidiabetic effects of *Homalium letestui* (Flacourtiaceae) in streptozotocin induced diabetic rats. *Res J Med Plant* 2007; 1: 134–138
- 187 Adeneye AA, Adeyemi OO. Further evaluation of antihyperglycaemic activity of *Hunteria umbellata* (K. Schum) Hallier f. seed extract in experimental diabetes. *J Ethnopharmacol* 2009; 126: 238–243
- 188 Ezeigbo II, Asuzu IU. Anti-diabetic activities of methanol leaf extract of *Hymenocardia acida* (TUL.) in alloxan-induced diabetic rats. *Afr J Tradit Complement Alternat Med* 2012; 9: 204–209
- 189 Danmalam UH, Abdullahi LM, Agunu A, Musa KY. Acute toxicity studies and hypoglycemic activity of the methanol extract of the leaves of *Hypptis suaveolens* Poit. (Lamiaceae). *Niger J Pharm Sci* 2009; 8: 87–92
- 190 Omonkhua AA, Onoagbe IO. Effects of long-term oral administration of aqueous extracts of *Irvingia gabonensis* bark on blood glucose and liver profile of normal rabbits. *J Med Plants Res* 2012; 6: 2581–2589
- 191 Momoh MA, Muhammed U. Toxicological and hypoglycemic studies on the oil extracted from seeds of *Khaya senegalensis* on blood glucose levels of alloxan-induced diabetic albino rats. *J Med Plants Res* 2011; 5: 4021–4024
- 192 Kolawole OT, Kolawole SO, Ayankunle AA, Olaniran OI. Anti-hyperglycemic effect of *Khaya senegalensis* stem bark aqueous extract in Wistar rats. *Eur J Med Plants* 2012; 2: 66–73
- 193 Bello A, Aliero AA, Saidu Y, Muhammad S. Hypoglycaemic and hypolipidaemic effects of *Leptadenia hastata* (Pers.) Decne in alloxan induced diabetic rats. *Niger J Basic Appl Sci* 2011; 19: 187–192
- 194 Osadebe PO, Omeje EO, Nworu SC, Esimone CO, Uzor PF, David EK, Uzoma JU. Antidiabetic principles of *Loranthus micranthus* Linn. parasitic on *Persea americana*. *Asian Pac J Trop Biomed* 2010; 3: 619–623
- 195 Osadebe PO, Omeje EO, Uzor PF, David EK, Obiorah DC. Seasonal variation for the antidiabetic activity of *Loranthus micranthus* methanol extract. *Asian Pac J Trop Biomed* 2010; 3: 196–199
- 196 Ugbenyen AM, Odetola AA. Hypoglycemic potential of the young leaves methanolic extract of *Magnifera indica* in alloxan induced diabetic rats. *Pak J Nutr* 2009; 8: 239–241
- 197 Okokon JE, Bassey SA, Osuji LC, Pius MU. Antidiabetic and hypolipidemic effect of *Mammea africana* (Guttiferae) in streptozotocin induced diabetic rats. *J Pharmacol Toxicol* 2007; 2: 278–283
- 198 Enomfo JA, Okokon JE, Offiong E. Antidiabetic and hypolipidemic activities of ethanolic leaf extract and fractions of *Melanthera scandens*. *Asian Pac J Trop Biomed* 2012; 2: 523–527
- 199 Ataman JE, Grillo DB, Omongbai EKI, Idu M, Amaechina F, Okonji V, Ayinde BA. Effect of methanolic extract of *Momordica charantia* leaves on alloxan treated Wistar rats. *J Med Sci* 2006; 6: 828–832
- 200 Tarkang PA, Ofogba CJ. Evaluation of the hypoglycemic activity and safety of *Momordica charantia* (Cucurbitaceae). *Afr J Pharm Sci Pharm* 2012; 3: 17–29
- 201 Odutuga AA, Dairo JO, Minari JB, Bamisaye FA. Ant-diabetic effect of *Morinda lucida* stem bark extracts on alloxan induced diabetic rats. *Res J Pharmacol* 2010; 4: 78–82
- 202 Adeneye AA, Agbaje EO. Pharmacological evaluation of oral hypoglycemic and antidiabetic effects of fresh leaves ethanol extract of *Morinda lucida* Benth. in normal and alloxan induced diabetic rats. *Afr J Biomed Res* 2008; 11: 65–71
- 203 Edoga CO, Njoku OO, Amadi EN, Okeke JJ. Blood sugar lowering effect of *Moringa oleifera* Lam in albino rats. *Int J Sci Technol* 2013; 3: 88–90
- 204 Salau BA, Ajani EO. Methanolic extract of *Musa sapientum* (L var. paradisiacal) sucker improves lipid profiles in alloxan induced diabetic rats. *Asian J Biol Sci* 2012; 5: 322–327
- 205 Adewoye EO, Ige AO, Latona CT. Effect of methanolic extract of *Musa sapientum* leaves on gastrointestinal transit time in normal and alloxan induced diabetic rats: possible mechanism of action. *Niger J Physiol Sci* 2011; 26: 083–088
- 206 Adeneye AA, Ajagbonna OP, Ayodele OW. Hypoglycemic and antidiabetic activities on the stem bark aqueous and ethanol extracts of *Musanga cecropioides* in normal and alloxan-induced diabetic rats. *Fito-terapia* 2007; 78: 502–505
- 207 Owolabi OJ, Amaechina FC, Okoro M. Effect of ethanol leaf extract of *Newbouldia laevis* on blood glucose levels of diabetic rats. *Trop J Pharm Res* 2011; 10: 249–254
- 208 Umar IA, Mohammed A, Dawud FA, Kyari H, Abdullahi M. Anti-diabetic action of the aqueous extract of *Ocimum suave* in alloxan-induced diabetic rats. *Afr J Biotechnol* 2012; 11: 9243–9247
- 209 Igbakin AP, Adanlawo IG. The effect of solvent extracts of *Parinari microphylla* on metabolites of alloxan induced diabetic rats. *Int J Biol Chem* 2009; 3: 49–55
- 210 Ndiaye M, Diatta W, Sy AN, Dieye AM, Faye B, Bassene E. Antidiabetic properties of aqueous barks extract of *Parinari excelsa* in alloxan-induced diabetic rats. *Fito-terapia* 2008; 79: 267–270
- 211 Odetola AA, Akinloye O, Egunjobi C, Adekunle WA, Ayoola AO. Possible antidiabetic and antihyperlipidaemic effect of fermented *Parkia biglobosa* (Jacq) extract in alloxan-induced diabetic rats. *Clin Exp Pharmacol Physiol* 2006; 33: 808–812
- 212 Saba AB, Oyagbemi AA, Azeze OI. Antidiabetic and haematinic effects of *Parquetina nigrescens* on alloxan induced type-1 diabetes and normocytic normochromic anaemia in Wistar rats. *Afr Health Sci* 2010; 10: 276–282
- 213 Edem DO. Hypoglycemic effects of ethanolic extracts of alligator pear seed (*Persea americana* Mill) in rats. *Eur J Sci Res* 2009; 33: 669–678
- 214 Nwanjo HU, Oze G, Okafor MC, Nwosu D, Nwankpa P. Protective role of *Phyllanthus niruri* extract on serum lipid profiles and oxidative stress in hepatocytes of diabetic rats. *Afr J Biotechnol* 2007; 6: 1744–1749
- 215 Mbaka GO, Ogbonnia SO, Oyeniran KJ, Awopetu PI. Effect of *Raphia hookeri* seed extract on blood glucose, glycosylated haemoglobin and lipid profile of alloxan induced diabetic rats. *Br J Med Med Res* 2012; 2: 621–635
- 216 Ikewuchi JC, Ikewuchi CC. Hypoglycemic, hypocholesterolemic and ocular-protective effects of an aqueous extract of the rhizomes of *Sansevieria senegambica* baker (Agavaceae) on alloxan-induced diabetic Wistar rats. *Am J Biochem Mol Biol* 2012; 2: 48–66
- 217 Onakpa MM, Ajagbonna OP. Antidiabetic potentials of *Cassia occidentalis* leaf extract on alloxan induced diabetic albino mice. *Int J PharmTech Res* 2012; 4: 1766–1769
- 218 Mohammed A, Atiku MK. Anti-hyperglycemic and anti-hyperlipidemic effect of leaves and stem bark methanol extracts of *Senna siamea* in alloxan induced diabetic rats. *Curr Res Cardiovasc Pharmacol* 2012; 1: 10–17

- 219 Odason EE, Kolawole JA. Anti-diabetic properties and brine shrimp toxicity of the aqueous extract of the root of *Cassia siamea* Lam (Cesalpiniaceae). Niger J Pharm Res 2007; 6: 66–69
- 220 Okokon JE, Bassey AL, Nwidi LL. Antidiabetic and hypolipidemic effects of ethanolic root extract of *Setaria megaphylla*. Int J Pharmacol 2007; 3: 91–95
- 221 Martins E, Adesina OO, Adewale GB, Kolawole SO, Tomilola EJ, Oye-mayowa VS. Comparative evaluation of the protective effect of ethanolic and methanolic leaf extracts of *Sida acuta* against hyperglycaemia and alterations of biochemical and haematological indices in alloxan induced diabetic rats. J Pharmacol Toxicol 2010; 5: 1–12
- 222 Kade JJ, Barbosa NBV, Ibukun EO, Igbakin AP, Nogueira CW, Rocha JBT. Aqueous extracts of *Sphenocentrum trilobata* attenuates streptozotocin-induced hyperglycaemia in rat models by modulating oxidative stress parameters. Biol Med 2010; 2: 1–13
- 223 Godwin M, Adeyemi O, Osinubi A, Noronha C, Okanlawon A. The effect of aqueous root extract of *Sphenocentrum jollyanum* on blood glucose level of rabbits. J Med Plants Res 2009; 3: 870–874
- 224 Isah AB, Ibrahim YKE, Abdulrahman EM, Ibrahim MA. The hypoglycaemic activity of the aqueous extract of *Stachytarpheta angustifolia* (Verbanaceae) in normoglycaemic and alloxan-induced diabetic rats. Pak J Biol Sci 2007; 10: 137–141
- 225 Olorunfemi AE, Patrick E, Arit E, Arnold I, Emmanuel O. Hypoglycemic effect of the seed extract of *Telfairia occidentalis* in rats. Pak J Biol Sci 2007; 10: 498–501
- 226 Teugwa CW, Boudjeko T, Tchinda BT, Mejiato PC, Zofou D. Anti-hyperglycaemic globulins from selected *Cucurbitaceae* seeds used as anti-diabetic medicinal plants in Africa. BMC Complement Alternat Med 2013; 13: 2–8
- 227 N'guessan K, Fofie NBY, Zirihi GN. Effect of aqueous extract of *Terminalia catappa* leaves on the glycaemia of rabbits. J Appl Pharm Sci 2011; 1: 59–64
- 228 Kingsley O, Iyere OO, Georgina EO, Esosa US, Kazeem AO, Frank AO. Effects of aqueous root extract of *Treculia africana* on blood glucose, lipid profile and body weight changes of streptozotocin-induced diabetic and normal rats. Int J Plant Physiol Biochem 2011; 3: 169–175
- 229 Prohp TP, Onoagbe IO. Anti-diabetic properties and toxicological studies of *Triplochiton scleroxylon* on the heart enzymes in normal and streptozotocin-induced diabetic rabbits. Pak J Nutr 2009; 8: 1025–1029
- 230 Sy GY, Cissé A, Nongonierma RB, Sarr M, Mbodj NA, Faye B. Hypoglycaemic and antidiabetic activity of acetonic extract of *Vernonia colorata* leaves in normoglycaemic and alloxan-induced diabetic rats. J Ethnopharmacol 2005; 98: 171–175
- 231 Adaramoye O, Amanlou M, Habibi-Razaei M, Pasalar P, Moosavi-Movahedi A. Methanolic extract of African mistletoe (*Viscum album*) improves carbohydrate metabolism and hyperlipidemia in streptozotocin-induced diabetic rats. Asian Pac J Trop Biomed 2012; 5: 427–433
- 232 Ngugi MP, Murugi NJ, Kibiti MC, Ngeranwa JJ, Njue MW, Maina D, Gathumbi KP, Njagi NE. Hypoglycemic activity of some Kenyan plants traditionally used to manage diabetes mellitus in Eastern province. J Diabetes Metab 2011; 2: 1–6
- 233 Njagi JM, Piero MN, Ngeranwa JN, Njagi ENM, Kibiti CM, Njue WM, Maina D, Gathumbi PK. Assessment of antidiabetic potential of *Ficus sycomorus* on alloxan-induced diabetic mice. Int J Diabet Res 2012; 1: 47–51
- 234 Karau GM, Njagi ENM, Machochi AK, Wangai LN, Kamau PN. Hypoglycemic activity of aqueous and ethylacetate leaf and stem bark extracts of *Pappea capensis* in alloxan-induced diabetic BALB/c mice. Br J Pharmacol Toxicol 2012; 3: 251–258
- 235 Dinku T, Tadesse S, Asres K. Antidiabetic activity of the leaf extracts of *Pentas schimperiana* subsp. *schimperiana* (A. Rich) Vatke on alloxan-induced diabetic mice. Ethiop Pharm J 2010; 28: 22–26
- 236 Kambouche N, Merah B, Derdour A, Bellahouel S, Bouayed J, Dicko A, Younos C, Soulimani R. Hypoglycemic and antihyperglycemic effects of *Anabasis articulata* (Forssk) Moq (Chenopodiaceae), an Algerian medicinal plant. Afr J Biotechnol 2009; 8: 5578–5583
- 237 Hamza N, Berkea B, Chezea C, Aglib A, Robinsona P, Ginc H, Moore N. Prevention of type 2 diabetes induced by high fat diet in the C57BL/6J mouse by two medicinal plants used in traditional treatment of diabetes in the east of Algeria. J Ethnopharmacol 2010; 128: 513–518
- 238 Eddouks M, Lemhadri A, Michel JB. Hypolipidemic activity of aqueous extract of *Capparis spinosa* L. in normal and diabetic rats. J Ethnopharmacol 2005; 98: 345–350
- 239 Sefi M, Fetoui H, Lachkar N, Tahraoui A, Lyoussi B, Zeghal N, Boudawara T. *Centaureum erythraea* (Gentianaceae) leaf extract alleviates streptozotocin-induced oxidative stress and β -cell damage in rat pancreas. J Ethnopharmacol 2011; 135: 243–250
- 240 Mohammed MS, Hossam FA, Samir AE, Osama MS. Biomedical effect of cinnamon extract on obesity and diabetes relevance in Wistar rats. Am J Biochem Mol Biol 2012; 2: 133–145
- 241 Nahla SE, Ismail MA. Hypoglycemic effect of *Cleome droserifolia* ethanolic leaf extract in experimental diabetes and on non-enzymatic antioxidant, glycogen, thyroid hormone and insulin levels. Diabetol Croat 2006; 35: 15–22
- 242 Mohamed A. Antidiabetic, antihyperlipidemic and antioxidant effects of aqueous of the roots of *Cynara cornigera* in alloxan induced experimental diabetic rats. Int J Pharmacol 2011; 7: 782–789
- 243 Houacine C, Elkhawad AO, Ayoub SMH. A comparative study on the anti-diabetic activity of extracts of some Algerian and Sudanese plants. J Diabet Endocrinol 2012; 3: 25–28
- 244 Jouad H, Maghrani M, Eddouks M. Hypoglycaemic effect of *Rubus fruticosus* L. and *Globularia alypum* L. in normal and streptozotocin-induced diabetic rats. J Ethnopharmacol 2002; 81: 351–356
- 245 Zeggwagh NA, Eddouks M, Lemhadri A. Study of hypoglycaemic and hypolipidemic effects of *Inula viscosa* L. aqueous extract in normal and diabetic rats. J Ethnopharmacol 2006; 108: 223–227
- 246 Eddouks M, Maghrani M, Michel JB, Zeggwagh NA. Study of the hypoglycaemic activity of *Lepidium sativum* L. aqueous extract in normal and diabetic rats. J Ethnopharmacol 2005; 97: 391–395
- 247 Sayed MR, Mourad IM, Sayed DA. Biochemical changes in experimental diabetes before and after treatment with *Mangifera indica* and *Psidium guajava* extracts. Int J Pharm Biomed Sci 2011; 2: 29–41
- 248 Zeggwagh NA, Sulpice T, Eddouks M. Anti-hyperglycaemic and hypolipidemic effects of *Ocimum basilicum* aqueous extract in diabetic rats. Am J Pharmacol Toxicol 2007; 2: 123–129
- 249 El-Khayat Z, Hussein J, Ramzy T, Ashour M. Antidiabetic antioxidant effect of *Panax ginseng*. J Med Plants Res 2011; 5: 4616–4620
- 250 Jouad H, Eddouks M, Lacaille-Dubois MA, Lyoussi B. Hypoglycaemic effect of *Spergularia purpurea* in normal and streptozotocin-induced diabetic rats. J Ethnopharmacol 2000; 71: 169–177
- 251 Jouad H, Lemhadri A, Maghrani M, Zeggwagh NA, Eddouks M. Cholesterol-lowering activity of the aqueous extract of *Spergularia purpurea* in normal and recent-onset diabetic rats. J Ethnopharmacol 2003; 87: 43–49
- 252 Benwahhoud M, Jouad H, Eddouks M, Lyoussia B. Hypoglycemic effect of *Suaeda fruticosa* in streptozotocin-induced diabetic rats. J Ethnopharmacol 2001; 76: 35–38
- 253 El Amrani F, Rhallab A, Alaoui T, El Badaoui E, Chakir S. Hypoglycaemic effect of *Thymelaea hirsuta* in normal and streptozotocin-induced diabetic rats. J Med Plants Res 2009; 3: 625–629
- 254 Eddouks M, Maghrani M, Michel JB. Hypoglycaemic effect of *Triticum repens* P. Beauv. in normal and diabetic rats. J Ethnopharmacol 2005; 102: 228–232
- 255 Abd-Elraheem AE, Muhammad MAS, Mahrour MAM. Effect of ginger extract consumption on levels of blood glucose, lipid profile and kidney functions in Alloxan induced-diabetic rats. Egypt Acad J Biol Sci 2009; 2: 153–162
- 256 Oyedemi SO, Adewusi EA, Aiyegoro OA, Akinpelu DA. Antidiabetic and haematological effect of aqueous extract of stem bark of *Azela africana* (Smith) on streptozotocin-induced diabetic Wistar rats. Asian Pac J Trop Biomed 2011; 1: 353–358
- 257 Islam MS, Choi H, Loots DT. Effects of dietary onion (*Allium cepa* L.) in a high-fat diet streptozotocin-induced diabetes rodent model. Ann Nutr Metab 2008; 53: 6–12
- 258 Gundidza M, Masuku S, Humphrey G, Magwa ML. Anti-diabetic activity of *Aloe excelsa*. Cent Afr J Med 2005; 51: 115–120
- 259 Loots DT, Pieters M, Islam MS, Botes L. Antidiabetic effects of *Aloe ferox* and *Aloe greatheadii* var. *davyana* leaf gel extracts in a low-dose streptozotocin diabetes rat model. S Afr J Sci 2011; 107: 1–6
- 260 Mellem JJ, Baijnath H, Odhav B. Effect of the methanolic extract of *Brachylaena discolor* in a streptozotocin-induced diabetic rat model. Afr J Pharm Pharmacol 2013; 7: 636–642
- 261 Islam MS. Effects of the aqueous extract of white tea (*Camellia sinensis*) in a streptozotocin-induced diabetes model of rats. Phytomedicine 2011; 19: 25–31
- 262 Ojewole JAO. Hypoglycaemic effect of *Clausena anisata* (Wild) Hook methanolic root extract in rats. J Ethnopharmacol 2002; 81: 231–237

- 263 Ojewole JAO, Adewunmi CO. Hypoglycaemic effects of methanolic leaf extract of *Catharanthus roseus* (Linn.) G. Don (Apocynaceae) in normal and diabetic mice. *Acta Med Biol* 2000; 48: 55–58
- 264 Deuschländer MS, Lall N, Van de Venter M, Hussein AA. Hypoglycemic evaluation of a new triterpene and other compounds isolated from *Euclea undulata* Thunb. var. *myrtina* (Ebenaceae) root bark. *J Ethnopharmacol* 2011; 133: 1091–1095
- 265 Ojewole JAO. Antinociceptive, anti-inflammatory and antidiabetic properties of *Hypoxis hemerocallidea* Fisch. & C.A. Mey. (Hypoxidaceae) corm ['African Potato'] aqueous extract in mice and rats. *J Ethnopharmacol* 2006; 103: 126–134
- 266 Mahomed IM, Ojewole JA. Hypoglycemic effect of *Hypoxis hemerocallidea* corm (African potato) aqueous extract in rats. *Methods Findings Exp Clin Pharmacol* 2003; 25: 617–623
- 267 Oyedemi SO, Yakubu MT, Afolayan AJ. Antidiabetic activities of aqueous leaves extract of *Leonotis leonurus* in streptozotocin induced diabetic rats. *J Med Plants Res* 2011; 5: 119–125
- 268 Dzeufiet PDD, Ohandja DY, Tédong L, Asongalem EA, Dimo T, Sokeng SD, Kamtchouing P. Antidiabetic effect of *Ceiba pentandra* extract on streptozotocin-induced non-insulin-dependent diabetic (NIDDM) rats. *Afr J Tradit Complement Alternat Med* 2007; 4: 47–54
- 269 George C, Lochner A, Huisamen B. The efficacy of *Prosopis glandulosa* as antidiabetic treatment in rat models of diabetes and insulin resistance. *J Ethnopharmacol* 2011; 137: 298–304
- 270 Ojewole JA. Hypoglycemic and hypotensive effects of *Psidium guajava* Linn. (Myrtaceae) leaf aqueous extract. *Methods Findings Exp Clin Pharmacol* 2005; 27: 689–695
- 271 Ojewole JAO. Analgesic, anti-inflammatory and hypoglycaemic effects of *Rhus chirindensis* (Baker F.) [Anacardiaceae] stem-bark aqueous extract in mice and rats. *J Ethnopharmacol* 2007; 113: 338–345
- 272 Oyedemi SO, Bradley G, Afolayan AJ. Beneficial effect of aqueous extracts of *Strychnos henningsii* Gilg in streptozotocin-nicotinamide induced type 2 diabetic Wistar rats. *Int J Pharmacol* 2011; 7: 773–781
- 273 Sokeng SD, Lontsi D, Moundipa PF, Jatsa HB, Watcho P, Kamtchouing P. Hypoglycemic effect of *Anacardium occidentale* L. methanol extract and fractions on streptozotocin-induced diabetic rats. *Glob J Pharmacol* 2007; 1: 1–5
- 274 Kamtchouing P, Kahpui SM, Dzeufiet D, Tédong L, Asongalem EA, Dimo T. Anti-diabetic activity of methanol/methylene chloride stem bark extracts of *Terminalia superba* and *Canarium schweinfurthii* on streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2006; 104: 306–309
- 275 Dimo T, Rakotonirina SV, Tan PV, Azay J, Dongo E, Kamtchouing P, Cros G. Effect of *Sclerocarya birrea* (Anacardiaceae) stem bark methylene chloride/methanol extract on streptozotocin-diabetic rats. *J Ethnopharmacol* 2007; 110: 434–438
- 276 Makom IGN, Frigerio F, Casimir M, Nguiguim FT, Dongo E, Kamtchouing P, Dimo T, Maechler P. *Sclerocarya birrea* (Anacardiaceae) stem-bark extract corrects glycaemia in diabetic rats and acts on beta-cells by enhancing glucose-stimulated insulin secretion. *J Endocrinol* 2010; 205: 79–86
- 277 Ojewole JAO, Adewole SO, Olayiwola G. Hypoglycaemic and hypotensive effects of *Momordica charantia* Linn (Cucurbitaceae) whole-plant aqueous extract in rats. *Cardiovasc J S Afr* 2006; 17: 227–232