Are endoscopic mucosal resection and endoscopic submucosal dissection risky for patients with cirrhosis?

Endoscopic mucosal resection (EMR) and endoscopic submucosal dissection (ESD) have become a standard treatment in selected cases of early gastric cancer [1,2]. The risk of complications may increase in patients with cirrhosis, due both to severe clotting impairment and to decreased platelet counts [3]. We present our experience in three patients with cirrhosis.

A 68-year-old woman with cryptogenic cirrhosis and esophageal varices, who had normal laboratory parameters and platelet count, was diagnosed with a high-grade gastric adenoma. ESD was decided on. Lesion margins were marked. A solution of mixed saline, adrenaline, and indigo-carmin was used for submucosal injection. Circumferential incision and submucosal dissection were carried out using a HookKnife and an ITknife2 (Olympus, Tokyo, Japan). A Coagrasper (Olympus) was used for preventive hemo-

stasis. The lesion was removed en bloc in 61 minutes (• Figs. 1, 2). Pathological examination revealed a high-grade gastric adenoma. A prophylactic antibiotic was administered. Endoscopy at 6-month follow-up showed no recurrence.

A 69-year-old man was diagnosed with decompensated cryptogenic cirrhosis (esophageal varices and ascites). Serum bilirubin and albumin levels were 1.9 mg/dL and 2.2 g/dL respectively. Platelet count was $59 \times 10^3/\mu\text{L}$ and INR was 1.42. Endoscopic evaluation showed a sessile polyp (width 10 mm) in the gastric body. The polyp was elevated by administering a mixture of saline and adrenaline, and was then removed en bloc using a polypectomy snare. Pathological examination revealed a grade 1 neuroendocrine tumor. Prophylactic antibiotic therapy was administered.

A 55-year-old woman was diagnosed as having hepatitis C virus-related cirrhosis. Blood chemistry was normal except for the platelet count ($91 \times 10^3/\mu\text{L}$). Five sessile polyps (the largest 15 mm, the others 6–8 mm) were seen in the gastric body during endoscopy. The largest polyp was elevated as described above and removed en bloc using a polypectomy snare. EMR with a ligation device was planned for the remaining polyps. The lesions were elevated, band ligation was applied, and EMR was performed below the band with a polypectomy snare (• Figs. 3, 4). Pathological examination revealed grade 1 neuroendocrine tumor for all lesions. A prophylactic antibiotic was administered. Patients with cirrhosis are at increased risk of developing gastric lesions [4]. Surgical treatment in these patients is associated with high mortality [5]. The risk of bleeding during ESD in patients with cirrhosis is reported to be 4%–33%. INR below 1.5 and a platelet count above $50 \times 10^3/\mu\text{L}$ are likely to be safe thresholds for endoscopic resection [3, 6–8]. There is a risk of sepsis in cirrhotic patients due to impairment of defensive mechanisms [9]. Prophylactic antibiotics can reduce the risk of bacteremia and sepsis.

Fig. 1 a Endoscopic appearance (i-scan) of gastric lesion. b Marking the lesion with the HookKnife. c Submucosal injection.

Fig. 2 a View of submucosal tissue through endoscope hood. b Mucosal defect after endoscopic submucosal dissection. c Resected specimen.

Fig. 3 a Endoscopic appearance of gastric lesion. b,c Application of endoscopic band ligation to lesions.

Fig. 4 a View after endoscopic resection with snare. b Mucosal defects after endoscopic mucosal resection. c Resected specimens

Endoscopy_UCTN_Code_TTT_1AO_2AG

Competing interests: None

**Fatih Aslan¹, Zehra Akpınar¹,
Ali Rıza Seren², Emrah Alper¹,
Cem Cekiç¹, Nese Ekinci³,
Sezgin Vatansever¹, Belkis Unsal¹**

¹ Department of Gastroenterology,
Katip Celebi University Atatürk Training
and Research Hospital, Izmir, Turkey

² Department of Anesthesiology and
Reanimation, Katip Celebi University
Atatürk Training and Research Hospital,
Izmir, Turkey

³ Department of Pathology, Katip Celebi
University Atatürk Training and Research
Hospital, Izmir, Turkey

References

- 1 Yamamoto H, Kita H. Endoscopic therapy of early gastric cancer. *Best Pract Res Clin Gastroenterol* 2005; 19: 909–926
- 2 Soetikno R, Kaltenbach T, Yeh R et al. Endoscopic mucosal resection for early cancers of the upper gastrointestinal tract. *J Clin Oncol* 2005; 23: 4490–4498
- 3 Ferro D, Angelico F, Caldwell SH et al. Bleeding and thrombosis in cirrhotic patients: what really matters? *Dig Liver Dis* 2012; 44: 275–279
- 4 Zullo A, Romiti A, Tomao S et al. Gastric cancer prevalence in patients with liver cirrhosis. *Eur J Cancer Prev* 2003; 12: 179–182
- 5 Douard R, Lentschener C, Ozier Y et al. Operative risks of digestive surgery in cirrhotic patients. *Gastroenterol Clin Biol* 2009; 33: 555–564
- 6 Kwon YL, Kim ES, Lee KI et al. Endoscopic treatments of gastric mucosal lesions are not riskier in patients with chronic renal failure or liver cirrhosis. *Surg Endosc* 2011; 25: 1994–1999
- 7 Repici A, Pagano N, Hassan C et al. Endoscopic submucosal dissection of gastric neoplastic lesions in patients with liver cirrhosis: a systematic review. *J Gastrointest Liver Dis* 2012; 21: 303–307
- 8 Zullo A, Hassan C, Bruzzese V. Comment to “Bleeding and thrombosis in cirrhotic patients: what really matters?” *Dig Liver Dis* 2012; 44: 1049
- 9 Garcia-Tsao G. Current management of the complications of cirrhosis and portal hypertension: variceal hemorrhage, ascites, and spontaneous bacterial peritonitis. *Gastroenterology* 2001; 120: 726–748

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1364946>
Endoscopy 2014; 46: E149–E150
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Fatih Aslan, MD
Yalı Mahallesi
I. Tarık Sarı Caddesi
Aktas Evleri-1, B Blok
No: 16, Daire: 5
35310, Güzelbahçe
İzmir
Turkey
drfatihhaslan@hotmail.com