Transgastric and percutaneous transhepatic rendezvous maneuver for the treatment of difficult intrahepatic stones

Interventional therapy of hepatic biliary stones is challenging if the patient has been subject to previous gastric or biliary surgery. Several variations have been reported for the endoscopic and endoscopic ultrasound (EUS)-guided transgastric approach to the biliary tract as salvage techniques after the failure of peroral endoscopic drainage [1]. A success rate of 89% has been reported, with complications seen in 12.5% of cases [2, 3].

In the present case, a 62-year-old patient with a Roux-en-Y biliodigestive anastomosis for an intraoperative dissection of the biliary tract that occurred during conventional cholecystectomy presented with progressive jaundice and recurrent colicky pain. Magnetic resonance cholangiopancreatography (MRCP) demonstrated intrahepatic gallstones.

The biliodigestive anastomosis could not be reached by enteroscopy. Therefore percutaneous transhepatic cholangiographic drainage (PTCD) with cholangioscopy was performed for the treatment of the intrahepatic stones. Despite this, persistently elevated levels of alkaline phosphatase and γ -glutamyltransferase were evident. Repeat MRCP showed a stone-filled, dilated biliary segment in the left hepatic lobe (► Fig. 1) that had not been possible to inspect by cholangioscopy during the PTCD (► Fig. 2) because of complete obstruction of the segmental orifice.

The dilated biliary segment was punctured transgastrically under EUS guidance (GF-UE160 linear echoendoscope; Olympus Deutschland GmbH, Hamburg, Germany) using a 19G needle (Cook Echo Tip, Cook Deutschland GmbH, Mönchengladbach, Germany) placed with contrast guidance (► Fig. 3). A JagWire (0.035 inch $\times$ 450 cm, straight tip; Boston Scientific GmbH, Ratingen, Germany) was inserted into the dilated segment and beyond the stenosis into the main biliary duct. The wire was grasped with a forceps via the percutaneous route by means of cholangioscopy. Balloon dilation (6 mm, MaxForce; Boston Scientific GmbH) of the segmental stenosis was carried out by the rendezvous procedure. This was followed by laser lithotripsy and complete stone extraction by

Fig. 1 Magnetic resonance cholangiopancreatography (MRCP) showing biliary stones in a single dilated biliary segment (arrow) of the left hepatic lobe.

Fig. 2 Percutaneous transhepatic cholangiographic drainage (PTCD) that failed to show the dilated segment later shown on magnetic resonance cholangiopancreatography (MRCP).

basket catheter (► Fig. 4), after which the wire was removed via the transcutaneous route.

In this procedure, not increasing the diameter of the transgastric route to more than the diameter of the needle minimises the risk of biliary leakage, therefore

making the transgastric approach to the biliary tract a possible alternative option, even in benign situations.

Endoscopy_UCTN_Code_TTT_1AR_2AH

Competing interests: None

Fig. 3 Endoscopic ultrasound (EUS) image showing the puncture of the dilated biliary segment using contrast guidance.

Fig. 4 PTCD and cholangioscopy showing the formerly obstructed segment now stone free after dilation of the orifice and extraction of the stone.

Erik-Sebastian Fuchs, Bettina Hartmann, Ralf Jakobs

Med. Klinik C, Klinikum der Stadt Ludwigshafen, Ludwigshafen, Germany

References

- 1 Yamao K, Hara K, Mizuno N et al. EUS-guided biliary drainage. *Gut Liver* 2010; 4: 67–75
- 2 Bories E, Pesenti C, Caillol F et al. Transgastric endoscopic ultrasonography-guided biliary drainage: results of a pilot study. *Endoscopy* 2007; 39: 287–291
- 3 Will U, Thieme A, Fuedner F et al. Treatment of biliary obstruction in selected patients by endoscopic ultrasonography (EUS)-guided transluminal biliary drainage. *Endoscopy* 2007; 39: 292–295

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1365113>
Endoscopy 2014; 46: E184–E185
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Erik-Sebastian Fuchs, MD
 Medizinische Klinik C
 Klinikum der Stadt Ludwigshafen gGmbH
 Bremsenstraße 79
 67063 Ludwigshafen
 Germany
 Fax: +49-621-5034114
 fuchse@klilu.de