Massive hemobilia following transpapillary bile duct biopsy treated by using a covered self-expandable metal stent

Fig. 1 **a** Cholangiogram showing luminal narrowing of the common bile duct with bile duct dilation. **b** Endoscopic transpapillary forceps biopsy was performed.

Fig. 2 After the transpapillary forceps biopsy, massive hemobilia developed.

Severe hemobilia after a forceps biopsy is very rare [1,2]. The management of uncontrolled hemobilia after endoscopic procedures includes percutaneous radiologic intervention or surgery [3,4]. Recently, successful endoscopic hemostasis using covered self-expandable metal stents (CSEMS) for uncontrolled bleeding following sphincterotomy, stent removal, or papillary balloon dilation has been reported [3–5]. We report a case of massive hemobilia after a transpapillary forceps biopsy of the bile duct and successful endoscopic hemostasis with placement of a CSEMS.

A 51-year-old woman was admitted because of abdominal pain and jaundice. Abdominal CT showed cancer of the pancreatic head with a stricture of the common bile duct (CBD). Cholangiogram showed dilatation of the proximal bile

duct with abrupt luminal narrowing of the CBD (▶ **Fig. 1a**). After endoscopic biliary sphincterotomy, transpapillary forceps biopsy was performed. A rat-tooth biopsy forceps (FB-39Q-1, Olympus, Tokyo, Japan) was used for tissue sampling at the constricted segment of the CBD (▶ **Fig. 1b**), but massive hemobilia occurred immediately afterwards (▶ **Fig. 2**, ▶ **Video 1**). To control the hemobilia, balloon tamponade with a controlled radial expansion balloon (Boston Scientific, Natick, MA, USA; 10 mm, 8 atm, 60 seconds) was attempted three times (▶ **Fig. 3a**). As the

Video 1

After a forceps biopsy was taken, sudden massive hemobilia occurred, flowing out of the duodenum.

uncontrolled hemobilia continued, a partially covered SEMS (10 mm diameter, 8 cm long; Bonastent, Standard Sci-Tech, Seoul, Korea) was inserted to achieve hemostasis through compression exerted by the expandable stent (▶ **Fig. 3b**). Once the CSEMS had been placed, the hemobilia decreased and then stopped. The patient's hemodynamic profile and vital signs were stable, and there was no further bleeding.

Complications related to transpapillary forceps biopsies are infrequent; usually, iatrogenic hemobilia is minor and can be controlled with conservative treatment [1]. Uncontrolled massive hemobilia after a transpapillary forceps biopsy has not previously been reported. Recently, placement of a CSEMS has been introduced as an effective means of endoscopic hemostasis for bleeding that could not be controlled with conventional methods [3–5]. If massive bleeding from the bile duct occurs during endoscopic retrograde cholangiopancreatography, hemostasis can be achieved promptly using a CSEMS, without radiological intervention or surgery. Endoscopic treatment using a CSEMS could be a useful way of providing effective hemostasis in selected patients with uncontrolled hemobilia of the extrahepatic bile duct.

Endoscopy_UCTN_Code_TTT_1AR_2AD

Competing interests: None

Fig. 3 **a** Fluoroscopy showing balloon tamponade performed in an attempt to control the hemobilia. **b** A partially covered self-expanding metallic stent was placed with nasobiliary drainage.

- 3 Itoi T, Yasuda I, Doi S et al. Endoscopic hemostasis using covered metallic stent placement for uncontrolled post-endoscopic sphincterotomy bleeding. *Endoscopy* 2011; 43: 369–372
- 4 Valats JC, Funakoshi N, Bauret P et al. Covered self-expandable biliary stents for the treatment of bleeding after ERCP. *Gastrointest Endosc* 2013; 78: 183–187
- 5 Aslinia F, Hawkins L, Darwin P et al. Temporary placement of a fully covered metal stent to tamponade bleeding from endoscopic papillary balloon dilation. *Gastrointest Endosc* 2012; 76: 911–913

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1365115>
Endoscopy 2014; 46: E161–E162
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Jong Ho Moon, MD, PhD
 SoonChunHyang University School of Medicine
 Digestive Disease Center
 SoonChunHyang University Bucheon Hospital
 170 Jomaru-ro, Wonmi-gu
 Bucheon 420-767
 Korea
 Fax: +82-32-6215080
jhmoon@schmc.ac.kr

**Jeong-Yeop Song, Jong Ho Moon,
 Hyun Jong Choi, Dong Choon Kim,
 Moon Han Choi, Tae Hoon Lee,
 Sang-Woo Cha, Young Deok Cho,
 Sang-Heum Park**

Digestive Disease Center and Research
 Institute, Department of Internal Medicine,
 SoonChunHyang University School of
 Medicine, Bucheon and Seoul, Korea

References

- 1 Jaiwal J, Fogel EL, Sherman S et al. Triple-tissue sampling at ERCP in malignant biliary obstruction. *Gastrointest Endosc* 2000; 51: 383–390
- 2 Ikeda M, Maetani I, Terada K et al. Usefulness of endoscopic retrograde biliary biopsy using large-capacity forceps for extrahepatic biliary strictures: a prospective randomized study. *Endoscopy* 2010; 42: 837–841