Rectal perforation caused by mesalazine enema in a patient with ulcerative colitis

Fig. 1 Colonoscopy in a 48-year-old woman with ulcerative colitis. **a** Colonoscopy revealed large depressions in the rectum below the peritoneal reflection. **b** Perforation was observed in one of the depressions.

Fig. 2 T2-weighted magnetic resonance imaging showed a continuous localized signal from the left to the dorsal aspect of the anus (arrow) and contiguity from the left side wall of the rectum to the external aspect of the wall (arrow head). Perianal abscess was suspected.

We report on a rare complication of rectal perforation caused by mesalazine enema in a patient with ulcerative colitis. A 48-year-old woman with ulcerative colitis presented with a relapse of the disease and was started on treatment with mesalazine enemas.

The patient returned to the hospital 14 days later complaining of perianal pain. Colonoscopy revealed two depressions in the rectum below the peritoneal reflection (Fig. 1 a); one of the depressions contained a perforation

(Fig. 1 b). T2-weighted magnetic resonance imaging showed a continuous localized signal from the left to the dorsal aspect of the anus and contiguity from the left side wall of the rectum to the external aspect of the wall. A perianal abscess was suspected (Fig. 2). Based on these findings, the patient was diagnosed as having a rectal perforation.

After bowel rest and treatment with a steroid and antibiotics, the pain disappeared and remission was achieved. The patient was discharged 10 days later. Colonoscopy after 3 months confirmed leveling out of the two depressions.

The cause of the perforation was thought to be the incorrect insertion method for the mesalazine enema [1]: the patient had held the mesalazine enema in her right hand, and inserted it while standing with legs apart. The correct insertion method requires the patient to lie down on their side with their top leg bent up toward their chest.

Endoscopy_UCTN_Code_CPL_1AJ_2AH

Competing interests: None

Keiichiro Kume, Tatsuyuki Watanabe, Jyunzi Oshima, Ichiro Yoshikawa, Masaru Harada

Third Department of Internal Medicine, University of Occupational and Environmental Health, School of Medicine, Kitakyusyu, Japan

References

- 1 Tanswell IJ, Irfan K, Koskowsky T et al. Rectal perforation in ulcerative colitis: complication of an enema tip. *Gastrointest Endosc* 2009; 69: 344

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1365148>
Endoscopy 2014; 46: E190
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Keiichiro Kume, MD, PhD
Third Department of Internal Medicine
University of Occupational and Environmental Health
School of Medicine
1-1, Iseigaoka, Yahatanishi-ku
Kitakyusyu 807-8555
Japan
Fax: +81-93-6920107
k-kume@med.uoeh-u.ac.jp