Novel use of a self-expanding metal stent for an esophageal stricture after radiofrequency ablation treatment of Barrett's esophagus

Fig. 1 Endoscopic images showing: **a** a tight stricture with circumferential ulceration at the proximal end of the area of Barrett's epithelium that had been treated by radiofrequency ablation (RFA); **b** a fully covered metal esophageal stent deployed in the esophagus.

Fig. 2 Radiographic image following injection of contrast showing a waist in the mid-portion of the stent, with free flow of contrast into the stomach.

Fig. 3 Endoscopic view of the esophagus following removal of the stent 2 months later.

Fig. 4 The fully covered, 23×105-mm, esophageal self-expanding metal stent (SEMS).

Fig. 5 Endoscopic appearance 6 months later showing a well-healed fibrotic stricture.

Radiofrequency ablation (RFA) is effective and safe in the treatment of Barrett's esophagus [1]. The incidence of esophageal stricture after RFA treatment is reported to be up to 8% [2]. Stricture rates may be increased with RFA of long-segment Barrett's esophagus. Strictures are treated endoscopically with balloons or Savary dilators; however, there is a risk of perforation with these treatments. We report on the successful treatment of a patient with a stricture following RFA using a self-expanding metal stent (SEMS). A 71-year old man with long-segment Barrett's esophagus (C7M7) and low-grade dysplasia underwent circumferential RFA. A month later he reported dysphagia and odynophagia, and endoscopy revealed a tight stricture with circumferential ulceration at the proximal end of the RFA-treated area of Barrett's epithelium (● **Fig. 1 a**).

A gastroscop with a 5.9-mm diameter was advanced to the proximal end of the stricture; however, the distal end of the stricture could not be traversed. A gastroscop with an 8.8-mm diameter was therefore inserted and a 9–12-mm extraction balloon (Extractor Pro RX; Boston Scientific, Natick, Massachusetts, USA) was introduced. Injection of contrast revealed a 4–5 cm long stricture in the mid-esophagus. A stent introducer was passed over a 450-cm, 0.035-inch guidewire (Dreamwire; Boston Scientific), which had been passed through the stricture under fluoroscopic guidance. A fully covered metal esophageal stent (23×105 mm,

WallFlex; Boston Scientific) was deployed (● Fig. 1b). A further attempt to pass the 5.9 mm gastroscope through the stricture was unsuccessful. The extraction balloon was reintroduced and injection of contrast showed a waist in the mid-portion of the stent, but with free flow of contrast into the stomach (● Fig. 2).

The stent was removed 2 months later (● Fig. 3 and ● Fig. 4) and after 6 months, the patient had no symptoms of dysphagia and was found to have a well-healed fibrotic stricture on endoscopy (● Fig. 5).

To our knowledge, this is the first case of an esophageal stricture occurring after RFA that was successfully treated by placement of a fully covered removable metal stent. Use of a self-expandable metal stent has also been reported for a stricture occurring after photodynamic therapy for Barrett's esophagus [3]. Treatment of tight strictures with metal stents may be a cost-effective treatment as it avoids the need for repeated dilations and the possible subsequent complications [4].

Endoscopy_UCTN_Code_CPL_1AH_2AJ

Competing interests: None

Traci Murakami, Bhaskar Banerjee, Nuri Ozden

Division of Gastroenterology, Hepatology and Nutrition, University of Arizona Medical Center, Tucson, Arizona, USA

References

- 1 Shaheen N, Sharma P, Overholt B et al. Radio-frequency ablation in Barrett's esophagus with dysplasia. *N Engl J Med* 2009; 360: 2277–2288
- 2 Garman K, Shaheen N. Ablative therapies for Barrett's esophagus. *Curr Gastroenterol Rep* 2011; 13: 226–239
- 3 Cheon YK. Metal stenting to resolve post-photodynamic therapy stricture in early esophageal cancer. *World J Gastroenterol* 2011; 17: 1379–1382
- 4 Van Halsema E, Wong Kee Song L, Baron T et al. Safety of endoscopic removal of self-expandable stents after treatment of benign esophageal diseases. *Gastrointest Endosc* 2013; 77: 18–28

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1365428>
 Endoscopy 2014; 46: E269–E270
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding authors

Nuri Ozden, MD
 1501 N. Campbell Avenue
 PO Box 245028
 Tucson
 AZ 85721
 USA
 Fax: +1-520-874-7133
 nozden1@gmail.com

Traci Murakami, MD
 1501 N. Campbell Avenue
 PO Box 245028
 Tucson
 AZ 85721
 USA
 traci.murakami@gmail.com