Impacted bone fragment in a small-bowel diverticulum: an extremely rare cause of obscure gastrointestinal bleeding

Fig. 1 Large-mouthed small-bowel diverticulum.

Fig. 2 Evidence of bleeding in a small-bowel diverticulum.

Fig. 3 a, b Small-bowel diverticulum with impacted bone fragment.

Obscure gastrointestinal tract bleeding (OGIB) is a common and challenging issue [1]. Here we present an extremely rare case of overt OGIB secondary to impacted bone fragment within a small-bowel diverticulum.

An 88-year-old man presented with hypotension and a 2-day history of maroon-colored stools. Initial hemoglobin concentration was 7.7 g/dL. Urgent upper endoscopy showed a large duodenal diverticulum with no signs of bleeding. Colonoscopy was also unremarkable. Capsule small-bowel study was attempted, but the capsule was retained in the duodenal diverticulum, requiring endoscopic retrieval. Mesenteric angiography was negative. Delayed images on Tc-99m-labeled red blood cell scan showed activity in the small bowel. Antegrade and retrograde

single-balloon enteroscopy to the distal jejunum and proximal ileum showed large-mouthed diverticula but no signs of active bleeding. Intraoperative enteroscopy showed a segment in the mid-small bowel with multiple large-mouthed diverticula, including one diverticulum with a blood clot suggestive of recent bleeding (Fig. 1, Fig. 2). Segmental resection of this small-bowel segment containing the bleeding diverticulum and surrounding large-mouthed diverticula (approx. 60 cm) was performed. Postoperatively, the patient's hemoglobin levels remained stable with no further episodes of gastrointestinal bleeding. Examination of the resected small-bowel specimen showed large diverticula with a single diverticulum having an impacted 3-cm bone fragment with hemorrhagic

changes in the surrounding mucosa (Fig. 3).

Most of the cases of OGIB are due to small-bowel pathology. Identifying and treating the etiology of OGIB can be challenging [2]. There are only few case reports of foreign body ingestion causing OGIB. Our case is the first report of a bone fragment causing OGIB [3,4].

Endoscopy_UCTN_Code_CCL_1AC_2AH

Competing interests: None

Hazem T. Hammad¹, Leen Al-Sayyed¹, Shoba Theivanayagam¹, Van Nguyen², Arthur Rawlings³, Matthew L. Bechtold¹

¹ Division of Gastroenterology, University of Missouri School of Medicine, Columbia, Missouri, USA

² Department of Pathology and Anatomical Sciences, University of Missouri School of Medicine, Columbia, Missouri, USA

³ Department of Surgery, University of Missouri School of Medicine, Columbia, Missouri, USA

References

- 1 Singh A, Baptista V, Stoicov C et al. Evaluation of small bowel bleeding. *Curr Opin Gastroenterol* 2013; 29: 119–124
- 2 Raju GS, Gerson L, Das A et al. American Gastroenterological Association (AGA) Institute technical review on obscure gastrointestinal bleeding. *Gastroenterology* 2007; 133: 1697–717
- 3 Samyn I, Reynaert H, Op de Beeck B et al. Recurrent gastrointestinal blood loss of obscure origin: report of an exceptional case. *Acta Gastroenterol Belg* 1998; 61: 382–384
- 4 Nanavati SA, Opekun AR, Hacken-Bitar JB et al. Migrating shell fragment as a cause of recurrent obscure gastrointestinal bleeding. *J Clin Gastroenterol* 2004; 38: 137–138

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1365430>
Endoscopy 2014; 46: E262
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Hazem Hammad, MD
Division of Gastroenterology
Five Hospital Drive
Columbia, MO 65212
United States of America
Fax: +1-573-884-4595
hammadh@health.missouri.edu