Clozapine-associated Development of Second-onset Obsessive Compulsive Symptoms in Schizophrenia: Impact of Clozapine Serum Levels and Fluvoxamine Add-on

M. Gahr, K. Rehbaum, B. J. Connemann

Department of Psychiatry and Psychotherapy III, University of Ulm, Ulm, Germany

Abstract

Among antiserotonergic second generation antipsychotics (SGA), particularly treatment with clozapine (CLZ) is associated with the development of second-onset obsessive compulsive symptoms (OCS) in schizophrenia. However, less is known regarding the factors that increase the individual susceptibility for the development of SGA-associated second-onset OCS in schizophrenia. Here we present the case of a 29-year-old female patient with disorganized schizophrenia who exhibited OCS due to fluvoxamine-induced elevation of CLZ serum levels via inhibition of CYP 1A2 und 2C19. The severity of the observed OCS featured an association with CLZ serum levels. The case illustrates the interaction between fluvoxamine add-on and CLZ serum levels on the development of OCS in schizophrenia and emphasizes the need of regular therapeutic drug monitoring.

Key words

adverse drug reactions · antipsychotic treatment · second generation antipsychotics · side effects · therapeutic drug monitoring

Introduction

Epidemiological studies indicate that patients diagnosed with schizophrenia feature a considerable lifetime risk for obsessive compulsive (OC) symptoms with prevalence rates ranging from 10–25% [1–4]; occasionally estimates even vary between 0.5 and 55% [5]. The neurobiological processes underlying this comorbidity have not yet been fully elucidated. Besides several heterogeneous pathogenetic concepts regarding OC symptoms in schizophrenia (e.g., independent disorders that coincide accidentally [6] or “schizo-obsessive subtype of schizophrenia” [7,8]), especially treatment with second-generation antipsychotics (SGA) is associated with a significant risk for the development of OC symptoms [9]. This particular adverse drug reaction (ADR) was first reported for treatment with clozapine (CLZ) [10,11]. In accordance with this observation, it was hypothesized that antiserotonergic SGA (such as CLZ or olanzapine [OLZ]) might be responsible for SGA-induced second-onset OC symptoms in schizophrenia [12–14]. Indeed, although there is evidence that OLZ may also facilitate therapeutic effects on second-onset OC symptoms in patients with schizophrenia [15,16], clinical studies showed that OC symptoms were significantly

more prevalent and severe in patients treated with CLZ or OLZ vs. amisulpride or aripiprazole [12,14]. Among the group of antiserotonergic SGA, CLZ was assessed to be particularly capable of inducing OC symptoms in patients with schizophrenia [6,9,14,17,18]. Here, selective serotonin reuptake inhibitors (such as fluvoxamine [FLV] as an add-on) were reported to be successful not only in the treatment of OC symptoms in schizophrenia [19], but also in CLZ-induced second-onset OC symptoms in schizophrenia [20].

In order to contribute to the available experience regarding the pharmacological conditions that may trigger the development of OC symptoms in patients with schizophrenia we report the following case of a patient who developed second-onset OC symptoms under a therapeutic regimen consisting of CLZ, OLZ, and add-on of FLV.

Case Report

A 29-year-old female patient was admitted to our psychiatric ward due to OC symptoms which had been increasing for 6 weeks and which induced significant psychological impairment in daily life. She was diagnosed with schizophrenia (originally classified as disorganized subtype according to DSM-IV-TR 295.10 criteria) 6 years ago (cranial MRI and cerebrospinal fluid without pathological findings); the family anamnesis was positive only for major depressive disorder (patient's brother); she did not use any psychotropic substances (alcohol, nicotine or any illicit drugs). Leading psychopathological findings on admission were reduced attention, slowed thinking and psychomotor function, blunted affect and OC symptoms {distressing intrusive thoughts and obsessive counting/arithmetic; the *Yale-Brown obsessive compulsive scale* (Y-BOCS) [21] was not performed} without unambiguous positive symptoms of schizophrenia. Notably, OC symptoms occurred for the first time in the patient's course of disease. Current medication was CLZ (250 mg per day; 50 mg-0–200 mg-0), FLV (50 mg O.D.), and OLZ (20 mg O.D.). During the last psychiatric in-patient treatment (2 years prior to the current hospital admission), a regimen of CLZ (250 mg per day) and FLV add-on (25 mg per day) was established. FLV was additionally given as an inhibitor of the cytochrome P450 monooxidase [CYP] 1A2 and 2C19 [22] in order to increase CLZ serum levels and to reduce side effects as well as negative symptoms [23,24]. At this time the patient already exhibited CLZ serum levels in the upper therapeutic range. Approximately 2 months prior to the current readmission, FLV was increased (from 25 to 50 mg per day) for unknown reasons and OLZ had been established as a new pharmacological strategy in the context of the ambulant treatment. After consultation of the psychiatrist, who was responsible for the ambulant treatment, it turned out that OLZ had been administered due to problems with sleep.

Apart from overweight (body mass index 31.9 kg/m²) and several anticholinergic side effects such as obstipation (defecation 2–3 times per week), dryness of the mouth and occasionally blurred vision, physical examination was without any pathology. The medication was taken in correspondence to the medically administered dose and regimen; CLZ serum levels on admission time were 834 ng/mL (reference range 350–600 ng/mL) and N-desmethyl-CLZ 288 ng/mL, respectively [CLZ serum levels were measured by high-performance liquid chromatogra-

phy (HPLC) at the Neurological Laboratory, Department of Psychiatry, University of Mainz].

Due to significant anticholinergic side effects, we tapered OLZ (the serum levels of which presumably have also been gradually increased by FLV-induced inhibition of CYP1A2 [25]; OLZ serum levels on day of admission 60 ng/mL) during the first week after admission; subsequently anticholinergic ADR subsided completely. As OC symptoms persisted thereafter and, based on corresponding evidence [5,17,26], the elevated CLZ serum level was suggested to be a possible trigger of the second-onset OC symptoms, the daily doses of FLV (25 mg per day) and CLZ (200 mg per day) were reduced 10 days after admission. Accompanied by a slight clinical improvement of OC symptoms, CLZ serum levels measured after one week under the above described regimen were still increased (CLZ 778 ng/mL; N-desmethyl-CLZ 435 ng/mL). Thus, CLZ was reduced to a daily dose of 150 mg and FLV was kept at 25 mg per day. After further 2 weeks, OC symptoms subsided completely and CLZ serum levels were within the reference range, albeit still comparatively high (CLZ 528 ng/mL; N-desmethyl-CLZ 304 ng/mL).

Discussion

Apart from the well-known potential of SGAs to induce second onset OC symptoms [6], this case also illustrates the influence of CLZ treatment dose and serum levels as well as treatment duration and FLV add-on on OC symptoms. It was already suggested that second-onset OC symptoms may be a dose-related side-effect of CLZ [6], corresponding with observations of second-onset OC symptoms in schizophrenia significantly improving after CLZ reduction [27–30]. Our case report supports this assumption. Furthermore, duration of treatment with antiserotonergic SGA was positively correlated with severity of second-onset OC symptoms in a recent clinical study [12]. In our case, the preceding 2-year treatment interval with CLZ might thus have contributed to the comparatively fast development of severe OC symptoms after increase of CLZ serum levels. In the present case, OC symptoms occurred after dose escalation of FLV and additional prescription of OLZ. This, firstly, suggests that the combined effects of increased CLZ serum levels and OLZ might have been responsible for the observed second-onset OC symptoms, considering that OLZ as an antiserotonergic SGA was described to feature the potential of inducing OC symptoms in schizophrenia [12,14]. However, OLZ was described to facilitate also therapeutic effects on second-onset OC symptoms in patients with schizophrenia [15,16], and thus, CLZ might have been the main culprit concerning the de novo-development of OC symptoms. Apart from that, the administration of two antiserotonergic antipsychotics (in the present case CLZ and OLZ) should be avoided and is against the tendency of recent guidelines. Secondly, our case illustrates the ambivalent role of FLV regarding OC symptoms in the treatment of schizophrenia, especially in patients treated with CLZ: on the one hand there is evidence regarding beneficial effects of FLV on second-onset OC symptoms in schizophrenia [19,21,31,32], on the other hand FLV add-on increases CLZ serum levels and thus may trigger the development of second-onset OC symptoms. Moreover, the extent of FLV-induced CYP inhibition also depends on FLV serum levels; these were not measured in the present case; however, information on FLV serum levels may have allowed a better

understanding of the presented clinical course and thus, parallel measurements of CLZ and FLV are suggested.

Finally, concerning the pharmacological treatment of patients with schizophrenia, it should be considered that the combination of CLZ and OLZ should be avoided and add-on of selective serotonin reuptake inhibitors to CLZ requires therapeutic drug monitoring.

Conflict of Interest

The authors declare no conflicts of interest.

References

- 1 Buckley P, Miller B, Lehrer D et al. Psychiatric comorbidities and schizophrenia. *Schizophr Bull* 2009; 35: 383–402
- 2 Lysaker P, Whitney K. Obsessive-compulsive symptoms in schizophrenia: prevalence, correlates and treatment. *Expert Rev Neurother* 2009; 9: 99–107
- 3 Achim A, Maziade M, Raymond E et al. How prevalent are anxiety disorders in schizophrenia? A meta-analysis and critical review on a significant association. *Schizophr Bull* 2011; 37: 811–821
- 4 Schirmbeck F, Zink M. Comorbid obsessive-compulsive symptoms in schizophrenia: contributions of pharmacological and genetic factors. *Front Pharmacol* 2013; 4: 99
- 5 Mukhopadhyaya K, Krishnaiah R, Taye T et al. Obsessive-compulsive disorder in UK clozapine-treated schizophrenia and schizoaffective disorder: a cause for clinical concern. *J Psychopharmacol* 2009; 23: 6–13
- 6 Schirmbeck F, Zink M. Clozapine-induced obsessive-compulsive symptoms in schizophrenia: a critical review. *Curr Neuropharmacol* 2012; 10: 88–95
- 7 Poyurovsky M, Koran L. Obsessive-compulsive disorder (OCD) with schizotypy vs. schizophrenia with OCD: diagnostic dilemmas and therapeutic implications. *J Psychiatric Res* 2005; 39: 399–408
- 8 Dominguez R, Backman K, Lugo S. Demographics, prevalence, and clinical features of the schizo-obsessive subtype of schizophrenia. *CNS Spectr* 1999; 4: 50–56
- 9 Scheltens Beduin A, Swets M, Machielsen M et al. GROUP. Obsessive-compulsive symptoms in patients with schizophrenia: a naturalistic cross-sectional study comparing treatment with clozapine, olanzapine, risperidone, and no antipsychotics in 543 patients. *J Clin Psychiatry* 2012; 73: 1395–1402
- 10 Patil V. Development of transient obsessive-compulsive symptoms during treatment with clozapine. *Am J Psychiatry* 1992; 149: 272
- 11 Baker R, Chengappa K, Baird J et al. Emergence of obsessive compulsive symptoms during treatment with clozapine. *J Clin Psychiatry* 1992; 53: 439–442
- 12 Schirmbeck F, Esslinger C, Rausch F et al. Antiserotonergic antipsychotics are associated with obsessive-compulsive symptoms in schizophrenia. *Psychol Med* 2011; 41: 2361–2373
- 13 Zink M, Englisch S, Dressing H. Neurobiology confirms psychopathology. On the antagonism of psychosis and obsessive-compulsive syndromes. *Psychopathology* 2008; 41: 279–285
- 14 Schirmbeck F, Rausch F, Englisch S et al. Differential effects of antipsychotic agents on obsessive-compulsive symptoms in schizophrenia: a longitudinal study. *J Psychopharmacol* 2013; 27: 349–357
- 15 Poyurovsky M, Dorfman-Etrog P, Hermesh H et al. Beneficial effect of olanzapine in schizophrenic patients with obsessive-compulsive symptoms. *Int Clin Psychopharmacol* 2000; 15: 169–173
- 16 Poyurovsky M, Kurs R, Weizman A. Olanzapine-sertraline combination in schizophrenia with obsessive-compulsive disorder. *J Clin Psychiatry* 2003; 64: 611
- 17 Reznik I, Yavin I, Stryker R et al. Clozapine in the treatment of obsessive-compulsive symptoms in schizophrenia patients: a case series study. *Pharmacopsychiatry* 2004; 37: 52–56
- 18 Lykouras L, Alevizos B, Michalopoulou P et al. Obsessive-compulsive symptoms induced by atypical antipsychotics. A review of the reported cases. *Prog Neuropsychopharmacol Biol Psychiatry* 2003; 27: 333–346
- 19 Reznik I, Sirota P. Obsessive and compulsive symptoms in schizophrenia: a randomized controlled trial with fluvoxamine and neuroleptics. *J Clin Psychopharmacol* 2000; 20: 410–416
- 20 Poyurovsky M, Hermesh H, Weizman A. Fluvoxamine treatment in clozapine-induced obsessive-compulsive symptoms in schizophrenic patients. *Clin Neuropharmacol* 1996; 19: 305–313

- 21 Goodman W, Price L, Rasmussen S *et al.* The Yale-Brown obsessive compulsive scale. I. Development, use, and reliability. *Arch Gen Psychiatry* 1989; 46: 1006–1011
- 22 Olesen O, Linnet K. Fluvoxamine-clozapine drug interaction: inhibition in vitro of five cytochrome P450 isoforms involved in clozapine metabolism. *J Clin Psychopharmacol* 2000; 20: 35–42
- 23 Lu M, Lane H, Lin S *et al.* Adjunctive fluvoxamine inhibits clozapine-related weight gain and metabolic disturbances. *J Clin Psychiatry* 2004; 65: 766–771
- 24 Kishi T, Hirota T, Iwata N. Add-on fluvoxamine treatment for schizophrenia: an updated meta-analysis of randomized controlled trials. *Eur Arch Psychiatry Clin Neurosci* 2013; 263: 633–641
- 25 Hiemke C, Peled A, Jabarin M *et al.* Fluvoxamine augmentation of olanzapine in chronic schizophrenia: pharmacokinetic interactions and clinical effects. *J Clin Psychopharmacol* 2002; 22: 502–506
- 26 Lin S, Su S, Pan C. Higher plasma drug concentration in clozapine-treated schizophrenic patients with side-effects of obsessive/compulsive symptoms. *Ther Drug Monit* 2006; 28: 303–307
- 27 Zink M, Knopf U, Kuwilsky A. Management of clozapine-induced obsessive-compulsive symptoms in a man with schizophrenia. *Aust N Z J Psychiatry* 2007; 41: 293–294
- 28 Englisch S, Esslinger C, Inta D *et al.* Clozapine-induced obsessive-compulsive syndromes improve in combination with aripiprazole. *Clin Neuropharmacol* 2009; 32: 227–229
- 29 Lin S, Su S, Pan C. Higher plasma drug concentration in clozapine-treated schizophrenic patients with side effects of obsessive/compulsive symptoms. *Ther Drug Monit* 2006; 28: 303–307
- 30 Schirmbeck F, Zink M. Obsessive-compulsive syndromes in schizophrenia: A case for polypharmacy? In: Ritsner M. (ed.). *Polypharmacy in Psychiatric Practice*. Luxembourg, Berlin: Springer; 2013
- 31 Poyurovsky M, Isakov V, Hromnikov S *et al.* Fluvoxamine treatment of obsessive-compulsive symptoms in schizophrenic patients: an add-on open study. *Int Clin Psychopharmacol* 1999; 14: 95–100
- 32 Faucher S, Dardennes R, Ghaëm O *et al.* Obsessive-compulsive symptoms treatment in schizophrenia. *Can J Psychiatry* 2005; 50: 423–428

received 20.02.2014
 revised 19.03.2014
 accepted 22.03.2014

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1374611>
Pharmacopsychiatry 2014; 47: 118–120
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0176-3679

Correspondence

M. Gahr, MD, MA
 Department of Psychiatry and Psychotherapy III
 University of Ulm
 Leimgrubenweg 12–14
 89075 Ulm
 Germany
 Tel.: +49/(0)731/500 61552
 Fax: +49/(0)731/500 21549
maximilian.gahr@uni-ulm.de