Migration of embolization coils placed to stop a severe iatrogenic bleed following endoscopic ultrasound-guided cystogastrostomy

Fig. 1 Transcatheter arterial embolization (TAE) with metal coils being performed by the radiology service in a patient who experienced severe bleeding after endoscopic ultrasound (EUS)-guided cystogastrostomy.

Fig. 2 Radiologic and endoscopic views of the coils 5 days after initial placement (red rings indicate coils on the border of the cyst; black rings indicate coils that have migrated into the pseudocyst cavity).

A 32-year-old man with a large symptomatic pseudocyst was referred for endoscopic ultrasound (EUS)-guided cystogastrostomy. The procedure was complicated by an iatrogenic bleed from the pseudocyst border, which was associated with hypovolemia and a significant drop in hemoglobin.

A fully covered self-expandable metal stent (SEMS) inserted between the stomach and the pseudocyst did not stop the bleeding. As a result, transcatheter arterial embolization (TAE) with metal coils was performed by our radiology service (► **Fig. 1**). Hemostasis was achieved immediately after TAE.

After 5 days the patient developed fever and pain. A gastroscope was used to inspect the interior of the pseudocyst cavity, and the metal stent was replaced by two pigtail stents. Large blood clots and necrotic material observed in the cyst were removed with a Roth net. During this procedure, two of the metal coils used during TAE were seen on the border of the cyst at the site of the bleeding; however, two other coils appeared to have migrated directly from the vessel into the pseudocyst cavity (► **Fig. 2**). After the procedure, no further evidence of hemorrhage was recorded. Endoscopic pseudocyst lavage was repeated 4 days later, and the patient was discharged home 2 weeks later.

Embolization of selected arteries with metal coils has been previously used to manage iatrogenic EUS-guided cystogastrostomy bleeds [1]. Migration of the coils from the embolized vessels into the gastrointestinal tract is extremely uncommon, with only one documented case of migration of a steel-wire coil into the stomach after arterial embolization [2]. No reports of the migration of metal coils into a pancreatic pseudocyst have been previously reported. TAE appears to be a minimally invasive, highly effective treatment in patients who hemorrhage after cystogastrostomy.

Endoscopy_UCTN_Code_CPL_1AH_2AC

Competing interests: None

**Antonino Granata¹, Gabriele Curcio¹,
Neville Azzopardi¹, Ilaria Tarantino¹,
Luca Barresi¹, Roberto Miraglia²,
Mario Traina¹**

¹ Endoscopy Service, Department of
Diagnostic and Therapeutic Services,
Mediterranean Institute for Transplanta-
tion and Advanced Specialized Therapies
(ISMETT), Palermo, Italy

² Radiology Service, Department of
Diagnostic and Therapeutic Services,
Mediterranean Institute for Transplanta-
tion and Advanced Specialized Therapies
(ISMETT), Palermo, Italy

References

- 1 *Brandon JL, Ruden NM, Turba UC et al.* Angio-
graphic embolization of arterial haemor-
rhage following endoscopic US-guided cyst-
gastrostomy for pancreatic pseudocyst
drainage. *Diagn Interv Radiol* 2008; 14:
57–60
- 2 *Takahashi T, Shimada K, Kobayashi N et al.*
Migration of steel-wire coils into the stom-
ach after transcatheter arterial embolization
for a bleeding splenic artery pseudoaneu-
rysm: report of a case. *Surg Today* 2001;
31: 458–462

Bibliography

DOI [http://dx.doi.org/
10.1055/s-0034-1377212](http://dx.doi.org/10.1055/s-0034-1377212)
Endoscopy 2014; 46: E306–E307
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Antonino Granata, MD
Endoscopy Service
ISMETT
Via Tricomi 5
90127 Palermo
Italy
Fax: +39-091-2192400 (specify Endoscopy Service)
agranata@ismett.edu