A novel method to obtain biopsy samples from proximal biliary strictures

Brush cytology and forceps biopsy are the most commonly used techniques to allow a diagnosis to be reached in patients found with suspected proximal biliary malignancies during endoscopic retrograde cholangiography. However, the sensitivity of brush cytology for diagnosis of malignant biliary strictures ranges only from 30% to 60% [1], while in the case of the biopsy forceps it may in some cases be impossible to advance the forceps to the level of the stricture. Recently, we had experience of three cases of proximal biliary stricture in which cholangiocarino-

ma was strongly suspected. In all three it proved possible to obtain sufficient biopsy samples for a diagnosis using a novel method.

After cannulation of the main bile duct and identification of the proximal stricture under fluoroscopic guidance, endoscopic sphincterotomy was performed over a guidewire (0.035 inch; Boston Scientific, Natick, MA, USA). The stricture was dilated using a 6- or 8-mm diameter balloon, followed by brushing of the stricture to obtain cytology specimens. Then, a stent pusher (Jet-Set stent introducer sys-

tem for endoprotheses, diameter 2.1 mm, length 165 cm; MTW Endoskopie, Wesel, Germany) was cut 8 cm from the distal end (► **Fig. 1**) and pushed forward over the guidewire until it reached the stricture. The guidewire was withdrawn and a biopsy forceps (diameter 1.8 mm, length 160 cm; Endo-Technik, Solingen, Germany) advanced through the pusher. Biopsy samples were obtained from the stricture level with the forceps (► **Fig. 2**), and a plastic stent was subsequently placed. Histopathological examinations of the biopsy specimens in all three cases reported adenocarcinoma, whereas only one of the three brush cytology samples was reported as malignant.

A similar method using a double-balloon enteroscopy forceps has been previously reported [2]. However, the double-balloon enteroscopy forceps is more expensive and impractical than the standard biopsy forceps, and may not be available

in many endoscopy centers. We believe that the method presented here using a biopsy forceps through a pusher can be performed advantageously to obtain adequate tissue samples from proximal biliary strictures for accurate diagnosis.

Endoscopy_UCTN_Code_TTT_1AR_2AD

Competing interests: None

Fatih Tekin, Ilker Turan, Galip Ersoz, Oktay Tekesin, Omer Ozutemiz

Department of Gastroenterology, Ege University Medical School, Izmir, Turkey

References

- 1 Weber A, Schmid RM, Prinz C. Diagnostic approaches for cholangiocarcinoma. *World J Gastroenterol* 2008; 14: 4131–4136
- 2 Kulaksiz H, Strnad P, Römpf A et al. A novel method of forceps biopsy improves the diagnosis of proximal biliary malignancies. *Dig Dis Sci* 2011; 56: 596–601

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377282>
Endoscopy 2014; 46: E363–E364
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Fatih Tekin, MD
Ege Üniversitesi Tıp Fakültesi
Gastroenteroloji Bilim Dalı
Bornova 35100
İzmir
Turkey
Fax: +90-232-3427764
drtekinfatih@gmail.com