Endoscopic ultrasound-guided Nd:YAG laser ablation of recurrent pancreatic neuroendocrine tumor: a promising revolution?

Fig. 1 Pre-treatment ^{68}Ga -DOTA-NOC positron emission tomography (PET) revealed a lesion in the residual pancreatic body with metabolic activity (arrow). ^{68}Ga -DOTANOC PET 1 year after laser ablation (on the right), showed no metabolic activity in the pancreatic region.

Fig. 2 Computed tomography scan.
a Recurrent pancreatic neuroendocrine tumor appeared as a 9-mm nodular area with early contrast enhancement on the pancreatic residual body. **b** At the same site after treatment, a 35-mm coagulative necrotic area was seen with the onset of a small fluid collection in the omental bursa.

tomography (PET) (● Fig. 1a). Two years earlier she had undergone curative distal pancreatectomy for previous PNET in the setting of multiple endocrine neoplasia type I. At computed tomography (CT) scan, the lesion appeared as a 9-mm nodular area with early contrast enhancement on the pancreatic residual body (● Fig. 2a).

The patient refused total pancreatectomy. Laser ablation under EUS guidance was therefore performed using a neodymium-doped yttrium aluminum garnet (Nd:YAG) laser, at 4.0W for 300 seconds (● Fig. 3). No complications occurred during the procedure. At CT immediately after laser ablation, the ablated lesion appeared as a well-defined 35-mm coagulative necrotic area (● Fig. 2b). Neither perilesional parenchymal alteration nor vascular damages were observed. The 2-month follow-up CT scan showed the ablated area to be 18 mm (● Fig. 4a); at 1 year the area was 9 mm (● Fig. 4b), with no metabolic activity on ^{68}Ga -DOTA-NOC PET (● Fig. 1b).

Laser ablation is a minimally invasive method for destroying tumors within solid organs, and works by directing low-power laser light energy into the tissue. It has been used for primary and secondary liver tumors [2–4]. The potential advantage of laser ablation over other laser-in-

Endoscopic ultrasound (EUS) has developed from a diagnostic tool into a therapeutic one [1]. A 46-year-old woman was diagnosed with recurrent pancreatic neu-

roendocrine tumor (PNET) by positive ^{68}Ga -[1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid]-1-NaI3-octreotide (^{68}Ga -DOTA-NOC) positron emission

Fig. 3 Endoscopic ultrasound-guided laser ablation. **a** A hyperechoic line identified the laser fiber protruding from the tip of the needle. **b** During laser ablation, a slowly enlarging and coalescing hyperechoic zone appeared around the distal tip of the laser fiber. **c** After laser ablation, the entire treated area was occupied by an irregular and poorly defined echogenic zone.

Fig. 4 Follow-up computed tomography scans. **a** At 2 months after laser ablation, the ablated area was further reduced (18 mm vs. 35 mm). **b** At 1 year after laser ablation, a 9-mm area of coagulative necrosis remained.

Research, Campus Bio-Medico University of Rome, Rome, Italy

⁴ Surgical Endoscopy Unit, Sacred Heart Catholic University of Rome, Rome, Italy

References

- 1 Carrara S, Petrone MC, Testoni PA et al. Tumors and new endoscopic ultrasound-guided therapies. *World J Gastrointest Endosc* 2013; 5: 141–147
- 2 Pacella CM, Bizarri G, Francica G et al. Analysis of factors predicting survival in patients with hepatocellular carcinoma treated with percutaneous laser ablation. *J Hepatol* 2006; 44: 902–909
- 3 Francica G, Petrolati A, Di Stasio E et al. Effectiveness, safety, and local progression after percutaneous laser ablation for hepatocellular carcinoma nodules up to 4 cm are not affected by tumor location. *AJR Am J Roentgenol* 2012; 199: 1393–1401
- 4 Vogl TJ, Freier V, Nour-Eldin NE et al. Magnetic resonance-guided laser-induced interstitial thermotherapy of breast cancer liver metastases and other noncolorectal cancer liver metastases: an analysis of prognostic factors for long-term survival and progression-free survival. *Invest Radiol* 2013; 48: 406–412
- 5 Di Matteo F, Martino M, Rea R et al. EUS-guided Nd:YAG laser ablation of normal pancreatic tissue: a pilot study in a pig model. *Gastrointest Endosc* 2010; 72: 358–363

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377376>
Endoscopy 2014; 46: E380–E381
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Francesco Maria Di Matteo, MD
 Digestive Endoscopy Unit
 Campus Bio-Medico University of Rome
 Via Alvaro del Portillo 200
 00128 Rome
 Italy
 Fax: +39-06-225411658
 f.dimatteo@unicampus.it

duced thermotherapies is the short application time and the well-defined ablation area. Moreover, the use of thinner laser fibers enables insertion into standard EUS-needles and their potential application in deep abdominal organs, such as the pancreas, which are untreatable using the percutaneous approach. To date, our group has demonstrated the efficacy and safety of laser ablation with Nd:YAG in an in vivo animal model [5]. The current case is the first time we have applied this minimally invasive laser treatment to the human pancreas.

These results must be further assessed in additional patients affected by PNETS who are not suitable for conventional treatments.

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests: None

**Francesco Di Matteo¹,
 Francesca Picconi¹,
 Margareth Martino¹, Monica Pandolfi¹,
 Claudio Maurizio Pacella²,
 Emiliano Schena³, Guido Costamagna⁴**

¹ Digestive Endoscopy Unit, Campus Bio-Medico University of Rome, Rome, Italy

² Radiology Unit, Regina Apostolorum Hospital, Albano, Rome, Italy

³ Unit of Measurements and Biomedical Instrumentation, Center for Integrated