A case of rectal bleeding caused by digestive endometriosis resembling colon cancer

Fig. 1 Endoscopic view showing polypoid endometriosis of the colonic mucosa.

Fig. 2 Oral and intra-venous contrast-enhanced computed tomography (CT) image (axial view) showing diffuse and irregular thickening of the wall of the sigmoid colon.

Fig. 3 Macroscopic view of the resected sigmoid colon showing the thickened, fibrotic wall, which was bright white in the areas where endometriosis was found.

Fig. 4 Microscopic view of the resected specimen stained immunohistochemically for CD10 showing positive staining of the endometrial stroma (magnification $\times 40$).

A 42-year-old unmarried, nulliparous woman presented at our hospital with rectal bleeding. The patient had had iron deficiency anemia in the past. She had normal, regular menses with mild dysmenorrhea and had not taken hormonal drugs. Her physical examination was unremarkable. Colonoscopy revealed widespread polypoid mucosal thickening and luminal narrowing, along with erythema and edema, in the sigmoid colon. It was not possible to pass the endoscope beyond the lesion (● Fig. 1). Multiple endoscopic biopsies were taken, but histopathologic examination was nondiagnostic. An abdominal computed tomography (CT) scan revealed eccentric thickening of the wall of the sigmoid colon (● Fig. 2). The patient proceeded to laparotomy, which revealed that the lesion had penetrated the sigmoid colon to the posterior

uterine wall. Segmental resection of the sigmoid colon with side-to-side anastomosis was performed. On pathological examination, beneath a 4-cm fleshy-grey polypoid mass, the colonic wall was thickened over a 5-cm length (● Fig. 3). Microscopy revealed polypoid endometriosis of the bowel. Native colonic mucosa was also observed around the areas of endometriosis (● Fig. 4). Intestinal endometriosis penetrating the bowel wall may lead to localized narrowing and obstruction [1]. Polypoid endometriosis is a rare variant of endometriosis that is characterized by multiple large polypoid masses that may resemble malignancies. Endoscopic ultrasound (EUS) can be used to determine the size, margins, and layer of origin, as well as additional morphologic features that can suggest the endome-

triosis. A hypoechoic lesion infiltrating the muscularis propria and the serosa of the rectal wall and extending outside the rectal wall may be seen during EUS [2]. The endoscopic appearance of the mucosa is not diagnostic, and endoscopic biopsies commonly yield specimens that are insufficient for pathologic diagnosis [3]. EUS-guided fine needle aspiration (FNA) can be useful in the evaluation of rectosigmoid endometriosis [4]. Surgery is the gold-standard diagnostic method for endometriosis [5].

Endoscopy_UCTN_Code_CCL_1AD_2AD

Competing interests: None

**Bulent Yilmaz¹, Selma Cukur²,
Rula Sahin³**

¹ Department of Gastroenterology, Bolu Izzet Baysal State Hospital, Bolu, Turkey

² Department of Pathology, Bolu Izzet Baysal State Hospital, Bolu, Turkey

³ Department of Radiology, Bolu Izzet Baysal State Hospital, Bolu, Turkey

References

- 1 Schwartz JL, Schwartz LB. Extrapelvic endometriosis. In: Diamond MP, Osteen KG, eds. Endometrium and Endometriosis. Maden,

Massachusetts: Blackwell Science; 1997: 247–254

- 2 Pishvaian AC, Ahlawat SK, Garvin D et al. Role of EUS and EUS-guided FNA in the diagnosis of symptomatic rectosigmoid endometriosis. *Gastrointest Endosc* 2006; 63: 331–335
- 3 Parker RL, Dadmanesh F, Young RH et al. Polypoid endometriosis: a clinicopathologic analysis of 24 cases and a review of the literature. *Am J Surg Pathol* 2004; 28: 285–297
- 4 Bhutani MS. Recent developments in the role of endoscopic ultrasonography in diseases of the colon and rectum. *Curr Opin Gastroenterol* 2007; 23: 67–73
- 5 Buchweitz O, Poel T, Diedrich K et al. The diagnostic dilemma of minimal and mild endometriosis under routine conditions. *J Am Assoc Gynecol Laparosc* 2003; 10: 85–89

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377378>
Endoscopy 2014; 46: E357–E358
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Bulent Yilmaz, MD
 Bolu Izzet Baysal State Hospital
 Department of Gastroenterology
 PK: 14280
 Bolu
 Turkey
 Fax: +90-374-2534615
yilmazb76@mynet.com