Endoscopic ultrasound-guided fine needle aspiration diagnosis of a focal fatty mass in the pancreas

Focal fatty masses of the pancreas, which include focal fatty infiltration, lipoma, and liposarcoma, are rare entities [1]. A well-differentiated lipogenic liposarcoma can mimic a benign lesion on radiologic examination and only cytohistology can resolve the doubt [2].

A 60-year-old woman was referred to our department because of an incidental ultrasound diagnosis of a lesion in the head of the pancreas. The patient was asymptomatic. An unenhanced computed tomography (CT) scan showed a 43-mm homogeneous, hypodense lesion at the pancreatic head (● Fig. 1a). Abdominal magnetic resonance imaging (MRI) showed the lesion to be hypointense on fast imaging employing steady-state acquisition (FIESTA) with fat saturation (● Fig. 1b) and hyper-

intense on T2-weighted fast-spin echo (FSE) acquisition without fat saturation (● Fig. 1c). No clear contact with abdominal fat was seen.

An endoscopic ultrasound (EUS) showed a lobulated, ill-defined, heterogeneous lesion, which was slightly hypoechoic compared with the surrounding pancreatic parenchyma but with internal hyperechoic strands (● Fig. 2a). The appearance was compatible with both a pancreatic lipoma and focal fatty infiltration, but even a liposarcoma could not be excluded. For this reason, we decided to perform an EUS-guided fine needle aspiration (FNA) with a 19-gauge needle (one passage) (● Fig. 2b), which revealed clearly visible fat globules on the smears (● Fig. 3a).

Cytohistologic examination showed mature adipocytes characterized by large fat droplets and eccentrically placed small nuclei without atypia. The adipocytes were interspersed with islands of normal pancreatic tissue (● Fig. 3b,c). Immunohistochemical staining for Ki-67 showed no mitotic activity (● Fig. 3d).

This focal fatty mass of the pancreatic head had some of the characteristics of a lipoma (no clear contact with abdominal fat on imaging), and some of focal fatty infiltration (no clear cytohistologic demarcation from pancreatic parenchyma) [1, 2]. This difficulty in distinguishing between these nosologic entities has been previously reported, but a clear distinction is not necessary because conservative treatment is indicated for both types of lesion. EUS-FNA, on the other hand, permits clear classification of such lesions where the differential diagnosis includes liposarcoma, which has a surgical indication. Only three other cases of EUS-FNA of a focal fatty mass of the pancreas have been reported [3–5], but considering the

Fig. 1 Imaging of the lesion in the pancreatic head. **a** Computed tomography (CT) scan showing a hypodense lesion (*) in the pancreatic head. **b** Magnetic resonance imaging (MRI) scan showing the lesion is hypointense on fast imaging employing steady-state acquisition (FIESTA) with fat saturation in axial view; and **c** hyperintense on T2-weighted fast-spin echo (FSE) acquisition without fat saturation.

Fig. 2 Endoscopic ultrasound (EUS) images showing: **a** a lobulated, ill-defined, heterogeneous lesion, which is slightly hypoechoic compared with the surrounding pancreatic parenchyma but with internal hyperechoic strands; **b** EUS-guided fine needle aspiration (FNA) being performed with a 19-gauge needle.

Fig. 3 Histocytological appearance showing: **a** on Hemacolor staining of the smear, several round empty spaces, both singly and in groups (most likely adipose tissue), with no atypical cells; **b** on a hematoxylin and eosin (H&E)-stained section, adipose tissue (white arrow) without atypia, interspersed with islands of normal pancreatic acini (black arrow); **c** on an S100-stained section, adipose tissue and islands of normal pancreatic acini; **d** on Ki-67 staining, the adipose tissue is negative, indicating no mitotic activity.

low probability of complications and the potential benefit for the patient, EUS-FNA should be considered for such pancreatic lesions.

Endoscopy_UCTN_Code_CCL_1AF_2AZ_3AD

Competing interests: None

Luca Barresi¹, Ilaria Tarantino¹, Dario Ligresti¹, Rosa Liotta², Gabriele Curcio¹, Antonino Granata¹, Caruso Settimo³, Mario Traina¹

¹ Endoscopy Service, Department of Diagnostic and Therapeutic Services, Mediterranean Institute for Transplantation and Advanced Specialized Therapies (ISMETT), Palermo, Italy

² Pathology Service, Department of Diagnostic and Therapeutic Services, Mediterranean Institute for Transplantation and Advanced Specialized Therapies (ISMETT), Palermo, Italy

tion and Advanced Specialized Therapies (ISMETT), Palermo, Italy

³ Radiology Service, Department of Diagnostic and Therapeutic Services, Mediterranean Institute for Transplantation and Advanced Specialized Therapies (ISMETT), Palermo, Italy

References

- 1 Secil M, Igci E, Goktay AY et al. Lipoma of the pancreas: MRI findings. *Comput Med Imaging Graph* 2001; 25: 507–509
- 2 Temizoz O, Gencellac H, Unlu E et al. Incidental pancreatic lipoma: computed tomography imaging findings with emphasis on diagnostic challenges. *Can Assoc Radio J* 2010; 61: 156–161
- 3 Di Matteo FM, Shimpi L, Pandolfi M et al. EUS diagnosis of pancreatic lipoma. *Gastrointest Endosc* 2006; 64: 146–148
- 4 Suzuki R, Irisawa A, Hikichi T et al. Pancreatic lipoma diagnosed using EUS-FNA. A case report. *JOP* 2009; 10: 200–203

- 5 Pausawasdi N, Apisarnthanarak P, Pongpailul A et al. Pancreatic lipoma diagnosed by EUS-FNA. *Gastrointest Endosc* 2012; 76: 668–669

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377391>
Endoscopy 2014; 46: E393–E394
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Luca Barresi, MD
Gastroenterology and Endoscopy Service
ISMETT
Via Tricomi 1
Palermo 90127
Italy
Fax: +39-091-2192400
lbarresi@ismett.edu