Histoplasmosis presenting as pancreatic head mass lesion and gastric outlet obstruction in an immunocompetent patient

Fig. 1 Computed tomography scan showing pancreatic head mass and dilated stomach.

Fig. 2 Endoscopic ultrasound image showing hypoechoic pancreatic mass with infiltration into duodenal walls.

Fig. 3 Fine-needle aspiration material showing fungal organisms (silver stain $\times 100$).

The natural habitat of histoplasma is fecal-enriched soil found in avian habitats. Clinical manifestations of histoplasmosis vary, from asymptomatic pulmonary infection (most common form) to disseminated disease [1,2]. Disseminated disease is rare in immunocompetent individuals [2]. Pancreatic histoplasmosis is an uncommon form of gastrointestinal histoplasmosis, and intestinal obstruction, which has been reported in immunocompromised patients, is a rare form of the disease [2,3]. This case illustrates the importance of performing FNA of pancreatic head lesions.

Endoscopy_UCTN_Code_CCL_1AF_2AZ_3AD

Competing interests: None

A 61-year-old woman presented with vomiting and progressive weight loss. Blood counts, and liver and kidney function tests were normal. Upper gastrointestinal endoscopy showed a grossly dilated stomach with food residue, thickened duodenal folds, and luminal obstruction in the second part of the duodenum. Abdominal computed tomography showed features suggestive of gastric outlet obstruction and head/uncinate process mass lesion of the pancreas causing duodenal obstruction (● **Fig. 1**). An endoscopic ultrasound (EUS) was done, and revealed a hypoechoic lesion in the head of the pancreas, with loss of planes between the pancreas and

the duodenal walls (● **Fig. 2**). EUS-guided fine-needle aspiration (FNA) was performed, and subsequent periodic acid-Schiff staining showed thin-walled, round yeast forms with histoplasma morphology, suggestive of pancreatic histoplasmosis (● **Fig. 3**). Test for human immunodeficiency virus was negative.

The patient was treated with itraconazole. Her symptoms improved gradually over the next month. However, sudden cardiac death occurred at 3 months due to arrhythmia.

Histoplasma are intracellular dimorphic fungi, which exist as mycelial (infective forms in soil) and budding yeast forms.

Narendra S. Choudhary¹, Rajesh Puri¹, Maneesh Paliwal¹, Mridula Guleria², Randhir Sud¹

¹ Institute of Digestive and Hepatobiliary Sciences, Medanta, The Medicity, Gurgaon, Haryana, India

² Institute of Cytopathology, Medanta, The Medicity, Gurgaon, Haryana, India

References

- 1 Jaiswal S, Vij M, Chand G et al. Diagnosis of adrenal histoplasmosis by fine needle aspiration cytology: an analysis based on five cases. *Cytopathology* 2011; 22: 323–328
- 2 Lamps LW, Molina CP, West AB et al. The pathologic spectrum of gastrointestinal and

hepatic histoplasmosis. Am J Clin Pathol 2000; 113: 64–72

- 3 Spivak H, Schlasinger MH, Tabanda-Lichauco R et al. Small bowel obstruction from gastrointestinal histoplasmosis in acquired immune deficiency syndrome. Am Surg 1996; 62: 369–372

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377401>
Endoscopy 2014; 46: E411–E412
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Rajesh Puri, MD

Institute of Digestive and Hepatobiliary Sciences
Medanta, The Medicity
Sector 38, Gurgaon 122001
NCR Delhi
India
Fax: +91-124-4834111
purirajesh69@gmail.com