Intraductal papillary mucinous neoplasia with malignant biliary stricture in pancreas divisum

Fig. 1 A positron emission tomography-computed tomography scan showing uptake by the pancreatic mass, lymph nodes, and multiple hepatic masses.

Fig. 2 Minor papilla showing a typical "fish-mouth" aspect, due to presence of large amounts of mucinous fluid.

A 72-year-old man was evaluated because of epigastric abdominal pain and jaundice. His medical history included partial gastrectomy performed for peptic ulcer disease. Physical examination revealed jaundice and an epigastric mass, with tender-

Fig. 3 After contrast injection through the minor papilla, a pancreatogram revealed dorsal duct dilation and several filling defects.

Fig. 4 Deep cannulation of the major papilla and contrast injection showed a tight distal biliary stricture due to malignant compression by the pancreatic neoplasia.

Fig. 5 Final radiological and endoscopic aspects showing a partially covered metallic stent (1 × 6 cm) inserted into the common bile duct, across the stricture.

ness of the upper abdomen. Magnetic resonance imaging revealed a large, mixed cystic and solid mass at the level of the pancreatic head measuring 6.8 × 5.4 cm. The mass resulted in superior mesenteric vein retraction and superior mesenteric artery distortion. There were also multiple pancreatic cysts, suggesting intraductal papillary mucinous neoplasia. In addition, various hepatic nodules were present, as well as enlarged peripancreatic lymph nodes. A positron emission tomography-computed tomography scan revealed uptake by the pancreatic mass, lymph nodes, and multiple hepatic masses (● Fig. 1). An abdominal ultrasound-

guided biopsy of a liver mass confirmed metastatic adenocarcinoma. Because of diffuse disease the patient was referred for palliative biliary endoscopic drainage. During endoscopic retrograde cholangiopancreatography, the minor papilla showed a typical "fish-mouth" aspect (● Fig. 2). A pancreatogram revealed dorsal duct dilation and several filling defects (● Fig. 3). Contrast injected into the major papilla showed a very small and thin pancreatic ventral duct, and a tight distal biliary stricture due to malignant compression by the pancreatic neoplasm (● Fig. 4). A partially covered metallic stent (1 × 6 cm) was inserted (● Fig. 5).

Jaundice improved and serum total bilirubin returned to normal (1.2 mg/dL) 10 days after the procedure. The patient then started chemotherapy treatment.

Endoscopy_UCTN_Code_CCL_1AF_2AZ_3AB

Competing interests: None

Angelo P. Ferrari, Erika P. Macedo, Fernanda P. Martins, Maris C. B. Souza, Gustavo A. de Paulo

Endoscopy, Hospital Israelita Albert Einstein, São Paulo, Brazil

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377431>
Endoscopy 2014; 46: E431–E432
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Angelo P. Ferrari, MD

Endoscopy
Hospital Israelita Albert Einstein
Rua Dr. Bacelar, 317 ap 231
São Paulo 04026-001
Brazil
Fax: +55-11-55716454
angelo.ferrari@outlook.com