Postgastrectomy protein-losing cytomegalovirus jejunitis in an immunocompromised patient

A 59-year-old man was admitted because of nausea and fatigue. He had been diagnosed with gastric adenocarcinoma 7 months earlier, and because of gastric outlet obstruction had undergone total gastrectomy with esophagojejunostomy and omentectomy (pT₃N_{3b}M₀R₁). Post-operatively, platinum-based chemotherapy was initiated, followed by radio-chemotherapy (5-fluorouracil + 45 Gy) approximately 2 weeks before admission. Initially, the nausea and fatigue were attributed to ongoing radiochemotherapy. During admission the patient became bed-bound, with pronounced anasarca edema secondary to severe hypoalbuminemia (lowest level 9.9 g/L, normal 35–52 g/L) despite maximal supportive care, suggestive of protein-losing enteropathy (proteinuria was only mild). From the esophagojejunostomy onwards, the mucosa was

severely inflamed with ulcerations, desquamation, and spontaneous submucosal hemorrhages for approximately 15 cm in the jejunum (► Fig. 1 a, b). Microscopic examination revealed extensive ulceration with granulation and dense infiltration of mononuclear cells, cytomegaly, and Cowdry inclusions (► Fig. 1 c). The diagnosis of cytomegalovirus (CMV) jejunitis was strengthened by positive serum CMV polymerase chain reaction (PCR; 3.84 log copies/mL). After 2 weeks of treatment with ganciclovir, the patient's general condition recovered and CMV PCR became negative. At 1 month after discharge, the patient was in good general condition with an albumin level of 31.2 g/L (► Fig. 1 d). Although CMV infection is mostly asymptomatic or gives rise to only mild mononucleosis-like symptoms, in immunocompromised or critically ill patients it can

cause life-threatening complications [1–3]. CMV enteritis occurs in only 4% of patients with CMV disease and is mostly restricted to a defined area rather than being panenteric [4]. As shown in inflammatory bowel diseases where CMV infects areas of already inflamed mucosa [5], it is likely that mucosal changes secondary to radiotherapy provide favorable conditions for CMV infection. The protein-losing enteropathy was most likely a combination of inflammatory exudation and villous atrophy with increased permeability associated with CMV infection [6]. This case illustrates that CMV should be included in the differential diagnosis of protein-losing enteropathy, especially following radiation therapy, and that biopsies should be taken to investigate CMV even when other potential explanations of mucositis are present.

Endoscopy_UCTN_Code_CCL_1AC_2AH

Competing interests: None

**Philip Roelandt¹, Helene Poels¹,
Ester Meeus², Eric Van Cutsem³,
Hans Prenen³**

¹ Gastroenterology Department, University
Hospitals Leuven, Leuven, Belgium

² Pathology Department, University
Hospitals Leuven, Leuven, Belgium

³ Digestive Oncology Department,
University Hospitals Leuven, Leuven,
Belgium

References

- 1 Whitley RJ, Jacobson MA, Friedberg DN et al. Guidelines for the treatment of cytomegalovirus diseases in patients with AIDS in the era of potent antiretroviral therapy: recommendations of an international panel. International AIDS Society – USA. Arch Intern Med 1998; 158: 957–969
- 2 Chamberlain RS, Atkins S, Saini N et al. Ileal perforation caused by cytomegalovirus infection in a critically ill adult. J Clin Gastroenterol 2000; 30: 432–435
- 3 Jain M, Duggal S, Chugh TD. Cytomegalovirus infection in non-immunosuppressed critically ill patients. J Infect Dev Ctries 2011; 5: 571–579
- 4 Baroco AL, Oldfield EC. Gastrointestinal cytomegalovirus disease in the immunocompromised patient. Curr Gastroenterol Rep 2008; 10: 409–416
- 5 Kaufman HS, Kahn AC, Iacobuzio-Donahue C et al. Cytomegaloviral enterocolitis: clinical associations and outcome. Dis Colon Rectum 1999; 42: 24–30
- 6 Nakase H, Itani T, Mimura J et al. Transient protein-losing enteropathy associated with cytomegalovirus infection in a noncompromised host: a case report. Am J Gastroenterol 1998; 93: 1005–1006

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377433>
Endoscopy 2014; 46: E427–E428
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Philip Roelandt, MD, PhD
Gastroenterology Department
University Hospitals Leuven
Herestraat 49
3000 Leuven
Belgium
Fax: +32-16-344387
Philip.roelandt@uzleuven.be