Perforation of benign peptic ulcer in hiatal hernia into the pericardium, resulting in pneumopericardium

Fig. 1 Chest radiograph showing pneumopericardium.

A 57-year-old woman with a medical history of hiatal hernia, peptic ulcer disease, alcoholism, and schizophrenia was admitted to the emergency department with epigastric pain and fever. The patient had not been taking her prescribed proton pump inhibitors.

A peptic ulcer perforation (1.5 cm) in the hiatal hernia sac was discovered during gastroscopy, and an attempt to close the perforation with metal clips was unsuccessful (● **Video 1**). The patient's general state became more severe following gastroscopy. Tachycardia with atrial fibrillation, dyspnea (saturation <80%), and hypotension (systolic blood pressure <60 mmHg) occurred. The patient was referred for surgery with suspicion of pneumopericardium (● **Fig. 1**).

A gastrotomy was made during laparotomy. A fistula to the pericardium was located and directly sutured. A tissue sample was taken from the ulceration. A cardiothoracic team performed pericardial drainage, which was complicated by the rupture of the fragile, inflamed wall of the right ventricle. Suturing of the right

ventricle through the sternotomy was performed. The patient died 36 hours later due to multiorgan failure.

Massive fibropurulent pericarditis was diagnosed in the autopsy. Specimens from the ulceration showed only fibrosis and inflammation with no trace of malignancy.

Peptic ulcer perforation located in the hiatal hernia sac with pneumopericardium is a very rare complication of benign stomach ulceration. Typically, this complication is related to malignant neoplasms of the stomach located in a hernia sac or in the fundus with invasion of the diaphragm and pericardium or as a complication after Nissen fundoplication [1]. Salling et al. described a case of pneumopericardium related to a gastric ulcer in a hiatal hernia, and cited seven similar cases from the literature [2]. The mortality rate is very high and it demands a multidisciplinary approach as well as a quick surgical response. Currently, there are no published guidelines or standards on how to manage such a complication [3,4]. We suggest that in the case of a gastropericardial fistula, a computed tomography scan of the chest and abdomen with administration of oral contrast should be performed instead of endoscopy, which could dramatically worsen circulatory insufficiency.

Endoscopy_UCTN_Code_CCL_1AB_2AD_3AC

Competing interests: None

Dariusz Łaski¹, Marian Łukiański¹, Michał Dubowik², Rafał Pawlaczyk³, Wojciech Żurek⁴, Tomasz Marjański⁴, Zbigniew Śledziński¹

¹ Department of General, Endocrine and Transplant Surgery, Medical University of Gdańsk, Gdańsk, Poland

² Department of Gastroenterology and Hepatology, Medical University of Gdańsk, Gdańsk, Poland

³ Department of Cardiosurgery, Medical University of Gdańsk, Gdańsk, Poland

⁴ Department of Thoracic Surgery, Medical University of Gdańsk, Gdańsk, Poland

References

- 1 Murthy S, Looney J, Jaklitsch MT. Gastropericardial fistula after laparoscopic surgery for reflux disease. *N Engl J Med* 2002; 346: 328–332
- 2 Salling N, Falenstein AM, Larsen LG. Non-traumatic perforation of gastric ulcer in a hiatal hernia to the pericardium. *Acta Med Scand* 1983; 213: 225–226
- 3 Letoquart JP, Fasquel JL, L'Huillier JP et al. Gastropericardial fistula. Review of the literature apropos of an original case. *J Chir* 1990; 127: 6–12
- 4 West AB, Nolan N, O'Brian DS. Benign peptic ulcers penetrating pericardium and heart: clinicopathological features and factors favoring survival. *Gastroenterology* 1988; 94: 1478–1487

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377437>
Endoscopy 2014; 46: E423
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Dariusz Łaski, MD
Invasive Medicine Center
Medical University of Gdańsk
ul. Smoluchowskiego 17
Gdańsk
Poland
Fax: +48-58-3493020
dlaski@gumed.edu.pl

Video 1

Gastroscopy with an unsuccessful attempt to close the perforation with metal clips.