Pocket-creation method of endoscopic submucosal dissection to achieve en bloc resection of giant colorectal subpedunculated neoplastic lesions

Fig. 1 Illustration showing the sequence of the pocket-creation method for endoscopic submucosal dissection of a giant, subpedunculated, neoplastic lesion. **a** A lesion with severe fibrosis at the center (lateral view). **b** A view of the same lesion from the top after submucosal injection. **c** Initial mucosal incision. **d** Creation of a submucosal pocket. **e** Extension of the submucosal pocket on both sides of the fibrosis. **f** Dissection of the fibrosis.

Fig. 2 Endoscopic images of the pocket-creation method of endoscopic submucosal dissection (ESD) for giant, subpedunculated, neoplastic lesions with severe fibrosis. **a** Endoscopic view of a subpedunculated, neoplastic lesion, 36 mm in diameter, in the ascending colon. **b** Severe fibrosis between the center of the lesion and the muscle layer after extension of the submucosal pocket on both sides. **c** The muscle layer was freed from the lesion after cutting the fibrotic area. **d** The muscle layer was flattened after ESD.

Endoscopic submucosal dissection (ESD) has become a standard minimally invasive therapy in Japan for large, superficial, colorectal neoplasms [1]. Although polypectomy or endoscopic mucosal resection is generally effective for polypoid lesions, ESD is an effective alternative to achieve en bloc resection in difficult cases, such as giant, subpedunculated, neoplastic lesions that are at least 3 cm in diameter [2]. However, ESD for giant, subpedunculated, neoplastic lesions is associated with technical difficulties because severe submucosal fibrosis occurs at the center of these lesions due to prolapse, resulting in the muscle layer being pulled upward. The pocket-creation method (PCM) is a new ESD strategy to overcome these difficulties (► Fig. 1; ► Video 1). The key feature of PCM is the creation of a large submucosal pocket under the lesion using a small-caliber tip transparent hood [3]. PCM has two advantages: maintenance of the thick submucosal layer with a minimal mucosal incision, which prevents the leakage of injection solution, and provision of good traction, which allows the submucosal tissue to be stretched and facilitates submucosal dissection.

PCM is very useful for ESD of giant, subpedunculated, neoplastic lesions with severe fibrosis (► Fig. 2). If there is severe fibrosis at the center of the lesion, submucosal pockets should be created on both sides of the fibrosis. By creating pockets on both sides of the fibrosis, the muscle layer can be clearly recognized. The muscle layer is pulled up at the fibrotic area, thus forming the shape of a mountain (► Fig. 2b). By using the small-caliber tip transparent hood and creating the pockets, the submucosal space is opened near the fibrotic area. Dissection of the fibrosis can then be easily achieved and the lesion is freed from the muscle layer (► Fig. 2c). After this, only dissection of the remaining mucosa is needed (► Fig. 2d; ► Fig. 3). PCM of ESD safely achieves an en bloc and complete resection of giant, subpedunculated, neoplastic lesions with severe fibrosis.

Video 1

The pocket-creation method of endoscopic submucosal dissection for a giant, subpedunculated, neoplastic lesion with severe fibrosis.

Endoscopy_UCTN_Code_TTT_1AQ_2AD

Fig. 3 Histology of the endoscopic submucosal dissection specimen showed a well-differentiated tubular adenocarcinoma in a high grade tubular adenoma (hematoxylin and eosin, Loupe image). The carcinoma was intramucosal cancer with no lymphovascular invasion and negative resection margins.

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377438>
 Endoscopy 2014; 46: E421–E422
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Hironori Yamamoto, MD, PhD
 Department of Medicine
 Division of Gastroenterology
 Jichi Medical University
 3311-1 Yakushiji
 Shimotsuke
 Tochigi, 329-0498
 Japan
 Fax: +81-285-448297
yamamoto@jichi.ac.jp

Competing interests: None

**Yoshikazu Hayashi¹, Keiichi Sunada¹,
 Haruo Takahashi¹, Hakuei Shinhata¹,
 Alan T. Lefor², Akira Tanaka³,
 Hironori Yamamoto¹**

¹ Department of Medicine, Division
 of Gastroenterology, Jichi Medical
 University, Shimotsuke, Japan

² Department of Surgery, Jichi Medical
 University, Shimotsuke, Japan

³ Department of Pathology, Jichi Medical
 University, Shimotsuke, Japan

References

- 1 Yamamoto H. Endoscopic submucosal dissection – current success and future directions. *Nat Rev Gastroenterol Hepatol* 2012; 9: 519–529
- 2 Choi YS, Lee JB, Lee EJ et al. Can endoscopic submucosal dissection technique be an alternative treatment option for a difficult giant (30 mm) pedunculated colorectal polyp? *Dis Colon Rectum* 2013; 56: 660–666
- 3 Yamamoto H, Kawata H, Sunada K et al. Successful en-bloc resection of large superficial tumors in the stomach and colon using sodium hyaluronate and small-caliber-tip transparent hood. *Endoscopy* 2003; 35: 690–694