Treating scar tissues during endoscopic submucosal dissection for gastrointestinal neoplasms

The extensive fibrosis in gastrointestinal neoplasms with severe scarring often makes endoscopic submucosal dissection (ESD) difficult and unsafe to perform. The submucosal fibrosis prevents the mucosa layer from lifting during injection. In addition, the vessels in scar tissues are sometimes extensive and barely visible. Moreover, the scar tissue is hard to cut with a standard endoscopic electrosurgical knife. Therefore, ESD for this type of lesion results in more complications, and is associated with a lower en bloc resection rate. Here, we describe a trick for dissection of these fibrotic tissues. A flat lesion was found near the cardia after previous endoscopic resection (► Fig. 1). The biopsy demonstrated high grade intraepithelial neoplasia. The ESD was started at the non-scarring site to create a wider submucosal layer for dissection before exposing the scar (► Fig. 2). A Coagrasper (FD-411UR; Olympus, Tokyo, Japan) was then used to grasp the fibrotic tissue. Using the electrosurgical generator (ESG-100; Olympus), a soft coagulation mode at 80 W was used briefly to coagulate the vessels in the fibrosis, and subsequently a pulse-cut-slow mode at 40 W was used to resect the tissue (► Video 1). The process was repeated to dissect the remaining fibrosis (► Fig. 3).

Video 1

Dissection of fibrotic tissues using a Coagrasper (Olympus, Tokyo, Japan) during endoscopic submucosal dissection.

A colonoscope type of Coagrasper was favored because its smaller jaws might easily reach the narrow scarring submucosal space and decrease the thermal damage to the surrounding tissues. As the scissors type knives are still unavailable outside of Japan, we suggest that the method described here would be useful for lesions with severe fibrosis.

Endoscopy_UCTN_Code_TTT_1AO_2AG

Competing interests: None

**Peng Jin, Jian-qiu Sheng,
Dong-liang Yu, Kuang-I Fu**

Department of Gastroenterology, Beijing
Military General Hospital, Beijing, China

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377439>
Endoscopy 2014; 46: E420
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Kuang-I Fu, MD
Department of Gastroenterology
Beijing Military General Hospital
Nanmenchang 5#
Dongcheng District
Beijing 100700
China
Fax: +86-10-66721299
fukuangi@hotmail.com

Fig. 1 A flat lesion near the cardia recurred after previous endoscopic resection (green arrows showed the margin).

Fig. 2 The submucosal scar was exposed during endoscopic submucosal dissection.

Fig. 3 Wound surface after completed endoscopic submucosal dissection.