Temporary placement of a fully covered self-expanding metal stent to allow therapeutic ERCP

A 64-year-old man with a previous history of laryngectomy followed by chemotherapy and radiotherapy was referred to us in January 2014 because of dysphagia. Endoscopy detected a 4-cm-long benign-appearing stricture in the hypopharynx (diameter 7 mm) and a 6-cm-long malignant circumferential lesion in the mid-esophagus (diameter 8 mm).

The patient was judged to be unfit for surgery, and therefore a fully covered self-expandable metal stent (FC-SEMS), 180×80 mm, was placed. Through-the-scope balloon dilation (CRE 10–11–12 mm; Microvasive, Boston Scientific Co., Natick, Massachusetts, United States) of the hypopharyngeal stricture was performed.

In February 2014 the patient developed a relapse of the hypopharyngeal stricture and jaundice due to compression of the common bile duct by the lymph nodes. Through-the-scope dilation of the stricture was performed, and a 14×16×100 mm FC-SEMS (Conio-Niti-S; Taewoong Medical, Seoul, Korea) was placed. The stent was removed 48 hours later immediately before therapeutic endoscopic retrograde cholangiopancreatography (ERCP). The duodenoscope could be easily inserted through the hypo-

pharyngeal stricture, and ERCP revealed a 3-cm stricture (diameter 1 mm) of the common bile duct. A 60×10-mm biliary SEMS was placed at the biliary stricture.

Refractory hypopharyngeal stenosis is the most frequent long-term complication after laryngectomy followed by radiotherapy [1,2]. A stricture in the proximal esophagus occurs in 10%–58% of these patients [3]. Introduction of a dedicated and effective hypopharyngeal removable SEMS has changed the endoscopic therapy of these patients [4]. In this case, it enabled the endoscopic palliation of a biliary stricture, by allowing for the passage of the duodenoscope through a previously severely stenosed esophagus (▶ **Video 1**).

Video 1

Temporary placement of a fully covered, self-expandable, metal stent at an esophageal stricture enabled passage of the duodenoscope for therapeutic endoscopic retrograde cholangiopancreatography.

Endoscopy_UCTN_Code_TTT_1AR_2AK

Competing interests: None

**Benedetto Mangiavillano¹,
Deborah A. Fisher², Massimo Conio¹**

¹ Gastroenterology and Gastrointestinal Endoscopy, General Hospital, Sanremo, Italy

² Division of Gastroenterology, Durham Veterans Affairs Medical Center, Duke University, Durham, North Carolina, United States

References

- 1 *Jalisi M, Jalisi S.* Advanced laryngeal carcinoma: surgical and non-surgical management options. *Otolaryngol Clin North Am* 2005; 38: 47–57
- 2 *Nguyen NP, Moltz CC, Frank C et al.* Severity and duration of chronic dysphagia following treatment for head and neck cancer. *Anticancer Res* 2005; 25: 2929–2934
- 3 *Kronenberg MB, Meyers AD.* Dysphagia following head and neck cancer surgery. *Dysphagia* 1994; 9: 236–244
- 4 *Conio M, Bianchi S, Filiberti R et al.* A modified self-expanding Niti-S stent for the management of benign hypopharyngeal strictures. *Gastrointest Endosc* 2007; 65: 714–720

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377440>
Endoscopy 2014; 46: E419
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Benedetto Mangiavillano, MD
Gastroenterology and Gastrointestinal Endoscopy
Borea Hospital
Via Giovanni Borea n°56
18038 – Sanremo (IM)
Italy
b_mangiavillano@hotmail.com