Squamous cell carcinoma of the sigmoid colon arising on base of a nonpolypoid tubulovillous adenoma

Fig. 1 A 14-mm polyp located in the sigmoid colon, showing a combined Kudo type IV and type V pattern.

Fig. 2 Hematoxylin and eosin routine stain of complete sigmoid polypectomy, with invasive carcinoma at the 11 o'clock position (×12.5).

Colonoscopy represents the gold standard for screening and surveillance of colorectal polyps [1]. Of note, retrieval of the resected specimen is of paramount importance, as histopathological analysis gives additional information regarding the polyp's entity and the resection status [2]. A 56-year-old woman was referred for screening colonoscopy, which revealed two nonpolypoid lesions in the sigmoid colon, measuring 25 mm and 14 mm, respectively. The smaller polyp had a 5-mm area at the top that showed a Kudo type V pattern (● Fig. 1). Both polyps were removed using snare polypectomy.

Histology of the smaller polyp showed a tubulovillous adenoma with apical high grade dysplasia and, at the base, an abrupt transition to solid growing nests of squamous epithelium with invasive nests reaching the middle third of the submucosal level (depth of invasion 1.3 mm) (● Fig. 2 and ● Fig. 3). Neither lymphatic nor blood vessel permeation was observed but a mild budding of tumor cells was apparent (pT1 [sm2], LO, VO, RO, G3). No pre-existing non-neoplastic squamous cell metaplasia was detected. Immunohistochemically, the squamous cell carcinoma was positive for CK5/6, CDX-2, MLH1, MSH2, MSH6, MSH3, PMS2, and MGMT

(● Fig. 4). The tumor was negative for CK7 and CK20. The adjacent tubulovillous adenoma was positive for CK20 and CDX-2, positive for MLH1, MSH2, MSH6, MSH3, PMS2, and MGMT, but negative for CK7 and CK5/6.

After interdisciplinary evaluation, elective resection of the sigmoid was performed but neither lymph node metastasis nor distant metastases were found.

Squamous cell metaplasia and carcinoma of the colon are extremely rare, reported in only 0.025% of colorectal neoplasms [3]. Interestingly, despite its low frequency, the entity of squamous cell cancer of the colon has been long known in the literature, and it has been reported to have a poor prognosis, with a mean survival time of 8.5 months at Stage IV [4,5]. The present case highlights the potential of endoscopic resection techniques to treat early colon cancer [6]. Endoscopists have to be aware that every polyp may hide a specific pathology, which may alter patient therapy and can therefore only be truly diagnosed by subsequent histopathological analysis.

Endoscopy_UCTN_Code_CCL_1AD_2AB

Competing interests: None

Fig. 3 Higher magnification of ● Fig. 2, showing invasive squamous cell carcinoma with mild tumor budding and abrupt changes to overlying tubulovillous adenoma (hematoxylin and eosin, ×200).

Fig. 4 Immunohistochemical cytokeratin 5/6 reaction showing the abrupt transition to squamous cell carcinoma and tubulovillous adenoma (CK5/6 ×100).

Helmut Neumann¹, Michael Vieth²

¹ Department of Medicine I, University of Erlangen-Nuremberg, Erlangen, Germany

² Institute of Pathology, Klinikum Bayreuth, Bayreuth, Germany

References

- 1 Mönkemüller K, Neumann H, Malfertheiner P et al. Advanced colon polypectomy. Clin Gastroenterol Hepatol 2009; 7: 641–652
- 2 Vieth M, Quirke P, Lambert R et al. European guidelines for quality assurance in colorectal cancer screening and diagnosis. First edition – annotations of colorectal lesions. Endoscopy 2012; 44: 131–139

- 3 Ambe P, Shadoun S, Granetzny S et al. Squamous cell carcinoma of the colon. A rare histological entity. Chirurg 2011; 82: 1116–1119
- 4 Frizelle FA, Hobday KS, Batts KP et al. Adenosquamous and squamous carcinoma of the colon and upper rectum: a clinical and histopathologic study. Dis Colon Rectum 2001; 44: 341–346
- 5 Herxheimer G. Über heterologe Cancroide. Ziegler's Beitr Path Anat u allgem Pathol 1907; 41: 412
- 6 Bergmann U, Beger HG. Endoscopic mucosal resection for advanced non-polypoid colorectal adenoma and early stage carcinoma. Surg Endosc 2003; 17: 475–479

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377493>
Endoscopy 2014; 46: E455–E456
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Michael Vieth, MD
Institute of Pathology
Klinikum Bayreuth
Preuschwitzer Str. 101
95445 Bayreuth
Germany
vieth.lkpathol@uni-bayreuth.de