Endoscopy_UCTN_Code_TTT_1AQ_2AG

Recurrent rectovaginal fistula: treatment with self-expanding metal stents

Competing interests: None

**Antonietta Lamazza¹, Enrico Fiori¹,
Alberto Schillaci¹, Antonio V.
Sterpetti¹, Emanuele Lezoche²**

¹ University of Rome La Sapienza, Department of Surgery, Istituto Pietro Valdoni, Rome, Italy

² University of Rome, Department of General Surgery, Surgical Specialties and Organ Transplantation Paride Stefanini, Rome, Italy

Postoperative rectovaginal fistula is a rare complication after colorectal resection for cancer. This adverse event results in a scenario of fear and misunderstanding for the patient [1–6]. Often, local inflammation, widespread infection, friability of the vaginal and rectal tissue, and the inevitably reduced perfusion of blood to the lower end of the rectum make any form of treatment hazardous and likely to cause complications. Even in cases in which the fistula heals, altered colorectal motility is common.

Self-expandable metal stents can be used to treat patients with rectovaginal fistula after colorectal resection for cancer. Of 10 patients who had rectovaginal fistula after colorectal resection for cancer and were treated with endoscopic placement of a self-expandable metal stent, three had been referred after multiple failed operations. All three patients had a diverting

proximal stoma. We used fully covered colonic stents 10 cm in length and 28 mm in diameter (Tae Woong Medical, Gimpo Si, Gyeonggi-do, South Korea) in all of them. There were no complications after the procedures. In one patient, the rectovaginal fistula healed without evidence of major fecal incontinence 8 months after stent insertion (► Fig. 1a–f). In the remaining two patients, the fistula decreased significantly in size (from 4 × 4 cm to 1 × 1 cm) without evidence of local tissue inflammation; both of these patients underwent a successful flap transposition 8 months after stent placement. In all three patients, the stent dislodged 3 months after placement, and a new stent was placed.

Self-expandable metal stents are a valid adjunct to the treatment of patients with complex, recurrent rectovaginal fistula after colorectal resection for cancer.

References

- 1 Ommer A, Herold A, Berg E et al. German S3-guideline: rectovaginal fistula. *Ger Med Sci* 2013; 10: 15
- 2 Schloricke E, Zimmermann M, Hoffmann M et al. Surgical treatment and prognosis of rectovaginal fistulae according to their origin. *Zentralbl Chir* 2012; 137: 390–395
- 3 Auyang ED, Santos BF, Enter DH et al. Natural orifice tranluminal endoscopic surgery: a technical review. *Surg Endosc* 2011; 25: 3135–3148
- 4 Jarrar A, Church J. Advancement flap repair: a good option for complex anorectal fistulas. *Dis Colon Rectum* 2011; 54: 1537–1541

Fig. 1 Recurrent rectovaginal fistula. **a** View from the rectal lumen. **b** Same patient; view from the vaginal lumen. **c** Insertion of a self-expanding metal stent. View from the rectal lumen. **d** Insertion of a self-expanding metal stent. View from the vaginal wall. **e** Complete healing of the fistula. There is still some inflammation immediately after removal of the stent. **f** Resolution of the inflammation 2 weeks after stent removal.

- 5 Zimmermann MS, Hoffmann M, Hildebrand P et al. Surgical repair of rectovaginal fistulas: a challenge. *Int J Colorectal Dis* 2011; 26: 817–819
- 6 Machado GR, Bojalian MO, Reeves ME. Transanal endoscopic repair of rectal anastomotic defects. *Arch Surg* 2005; 140: 1219–1222

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377497>
Endoscopy 2015; 47: E149–E150
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Antonio V. Sterpetti, MD
Istituto Pietro Valdoni
Viale del Policlinico
00167 Rome
Italy
Fax: +39-6-49972245
antonio.sterpetti@uniroma1.it