Esophageal stricture caused by a mediastinal hamartoma invading the esophageal wall

Fig. 1 An esophageal stricture located in his upper esophagus, and smooth mucosa at the narrow segment of the esophagus.

Fig. 3 Computed tomography showed thickening of the mucosa at the upper esophagus.

Fig. 2 Narrow-band imaging did not show any abnormality of the mucosal surface.

Fig. 4 The esophageal stricture interfered with the insertion of the endoscope.

A 60-year-old man presented with dysphagia, which had persisted for 2 months. A barium meal test revealed a stricture in the upper esophagus, and smooth mucosa at the narrow segment of the esophagus (● Fig. 1). He subsequently underwent endoscopy. Endoscopic examination revealed a mucosal bulge in the esophagus at a distance of 18–24 cm from the incisors, extending to about half of the esophageal wall, and with smooth mucosa at the ridge surface. Narrow-band imaging did not show any abnormality of the mucosal surface (● Fig. 2). Although the esophagus was narrow, it allowed insertion of the endoscope. Gross endoscopic appearance indicated that the lesion would be firm when touched with biopsy forceps. Biopsy revealed inflammatory granulation tissue, and chest computed tomography showed thickening of the mucosa at the upper esophagus (● Fig. 3).

After 4 weeks, the patient underwent a follow-up endoscopic ultrasound examination. On this occasion, an esophageal stricture was observed, which interfered with the insertion of the endoscope. In addition, the mucosa appeared normal at the narrow segment of the esophagus (● Fig. 4). Ultrasound examination revealed an obvious thickening of the mucosal layer of the lesion, which appeared asymmetrically annular. There was an unclear boundary between the local and surrounding tissues (● Fig. 5). Open surgery was performed because the patient experienced a gradual worsening of dysphagia symptoms. During surgery, a tumor was observed between the esophagus and the trachea, which invaded the esophageal wall and had unclear boundaries with the trachea. Because the possibility of a malignant tumor could not be ruled out, esophagus detachment and

subtotal resection were performed. Post-operative histopathological examination of the upper esophagus tissues showed the presence of chronic inflammation in the mucosa, squamous epithelial hyperplasia, and chronic suppurative inflammation of the subepithelial focal lesion. Examination of the paraesophageal tumor showed proliferation of tissues including fibers, blood vessels, lymphatic and striated muscles, and nerves with hamartoma-like changes, associated with local inflammatory cell infiltration, degeneration, and necrosis (● Fig. 6). Hamartomas rarely occur in the esophagus. In this case, the tumor was externally located but had invaded the esophageal wall, which manifested as an esophageal stricture, thereby precluding a diagnosis before surgical operation.

Fig. 5 Endoscopic ultrasound revealed an obvious thickening of the mucosal layer of the lesion.

Fig. 6 Proliferation of tissues including fibers, blood vessels, lymphatic and striated muscles, and nerves with hamartoma-like changes.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AB

Competing interests: None

Xiao-jun Zhao, Hui Xie, Dong-liang Yu, Jian-qiu Sheng

Department of Gastroenterology, Beijing Military General Hospital, Beijing, China

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377533>
 Endoscopy 2015; 47: E151–E152
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Jian-qiu Sheng, MS
 Department of Gastroenterology
 Beijing Military General Hospital
 Nanmencang 5#
 Dongcheng District
 Beijing 100700
 China
 Fax: +86-10-66721299
jianqiu@263.net