Intraductal ultrasonography-directed endoscopic retrograde biliary drainage without fluoroscopy

Fig. 1 Intraductal ultrasonography (IDUS)-directed endoscopic retrograde biliary drainage (ERBD) without fluoroscopy. **a** Bile aspiration after biliary cannulation. **b** IDUS views of a stone in the common bile duct. **c** Placement of stent for ERBD using IDUS without fluoroscopy. **d** Plain abdominal radiograph after ERBD insertion.

Common bile duct (CBD) stones may cause septic shock and lead to life-threatening conditions [1]. Endoscopic sphincterotomy (EST) and stone removal is an accepted method for the treatment of CBD stones [2]. EST has the potential to cause bleeding in patients with a bleeding risk. However, endoscopic retrograde biliary drainage (ERBD) can be performed effectively and safely without EST in these patients. ERBD has been performed under fluoroscopy, resulting in a relatively long duration of radiation exposure [3,4]. Intraductal ultrasound (IDUS) provides real-time cross-sectional imaging within the bile duct. IDUS can be performed without fluoroscopy. Therefore, IDUS-directed ERBD can be used to reduce the time of radiation exposure for both examiners and patients. This report describes IDUS-directed ERBD with a plastic stent without fluoroscopy in patients with a risk of bleeding.

A total of nine IDUS-directed ERBD procedures without fluoroscopy were performed. All procedures were performed using a standard side-viewing duodenoscope (TJF-160F; Olympus, Tokyo, Japan). After cannulation with a 0.035-inch guidewire (Jagwire; Boston Scientific, Natick, Massachusetts, United States), the “bile aspiration” technique was used to indicate bile duct cannulation (● Fig. 1 a) [5]. A 2.0-mm IDUS probe with a frequency of 20-MHz (UM-G20-29R; Olympus) was advanced over the guidewire into the bile duct. After images of ductal anatomy and lesions had been obtained (● Fig. 1 b), the insertion length of the IDUS probe, from the papilla of Vater to the lesions, was used to determine the length of plastic stent (Percuflex DUODENAL BEND Biliary Stent; Boston Scientific) that was required. The plastic stent for ERBD was inserted over the guidewire and placed in the correct position after withdrawal of the IDUS

probe (● Fig. 1 c). The location of the ERBD stent was confirmed by plain radiography (● Fig. 1 d).

Successful intubation of the papilla, selective CBD cannulation, and appropriate ERBD were achieved in all patients (9/9, 100%). No complications were observed in any patient.

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests: None

Ji-Myoung Lee¹, Chang-Hwan Park¹, Soo-Jung Rew², Ha-Na Kim¹, Ho-Jun Lee¹, In-Hyung Park¹, Seon-Young Park¹

¹ Department of Internal Medicine, Chonnam National University Medical School, Gwangju, Korea

² Department of Internal Medicine, Gwangju Christian Hospital, Gwangju, Korea

References

- 1 Lu J, Guo CY, Xu XF et al. Efficacy of intraductal ultrasonography in the diagnosis of non-opaque choledocholith. *World J Gastroenterol* 2012; 18: 275–278
- 2 Jiang XW, Tang SH, Yang JQ et al. Ultrasound-guided endoscopic biliary drainage: a useful drainage method for biliary obstructions. *Dig Dis Sci* 2014; 59: 161–167
- 3 Tsapaki V, Paraskeva KD, Mathou N et al. Patient and endoscopist radiation doses during ERCP procedures. *Radiat Prot Dosimetry* 2011; 147: 111–113
- 4 Lorenzo-Zuniga V, Alvarez MA, Moreno de Vega V et al. Predictive factors of radiation dose in ERCP: a prospective study in 2 tertiary centers. *Surg Laparosc Endosc Percutan Tech* 2013; 23: 266–270
- 5 Siegel JH, Rodriguez R, Cohen SA et al. Endoscopic management of cholangitis: critical review of an alternative technique and report of a large series. *Am J Gastroenterol* 1994; 89: 1142–1146

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377587>
Endoscopy 2014; 46: E495
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Chang-Hwan Park, MD, PhD

Department of Internal Medicine
Chonnam National University Medical School
42 Jebongro, Dong-gu
Gwangju, 501-757
Korea
Fax: +82-62-2258578
p1052ccy@hanmail.net