Endoscopic intraductal radiofrequency ablation of remnant intrapapillary mucinous neoplasm with acute hemorrhage after incomplete surgical resection

Fig. 1 Pancreatic duct system in the head with visible stenosis caused by the bleeding tumor.

Fig. 2 Radiofrequency ablation probe positioned in the stenosis during ablation.

Fig. 3 Computed tomography scan 2 days after radiofrequency ablation, showing a hypo-dense lesion in the pancreatic head, corresponding to the ablation area.

An 83-year-old woman presented with recurrent hemosuccus pancreaticus. The patient had previously undergone a distal pancreatic resection due to intrapapillary mucinous neoplasm, with main duct involvement of the tail. Upper endoscopy revealed active bleeding through the ampulla of Vater. During a previous episode

of pancreatic bleeding, intraductal injection of fibrin glue had been performed. Endoscopic retrograde pancreatography demonstrated dilated pancreatic ducts of the remnant pancreatic head. Approximately 3 cm above the papilla, the main pancreatic duct showed a 10-mm-long irregular stricture (▶ **Fig. 1**). As the pa-

tient refused surgical treatment, radiofrequency ablation (RFA) was performed using a bipolar RFA catheter (EndoHPB; EMCision, Montreal, Canada) designed for biliary RFA. The RFA catheter was advanced over a 0.035-inch guidewire (Jagwire; Boston Scientific Corp., Natick, Massachusetts, USA) (▶ **Fig. 2**). RFA was applied for a total of 90 seconds using 8 W soft coagulation mode, effect 1 (ERBE VIO 300 D; ERBE Elektromedizin GmbH, Tübingen, Germany). The patient developed mild pancreatitis following RFA, with a maximum serum lipase of 9.6 $\mu\text{mol/L}$ after 12 hours, which returned to normal within 24 hours. Computed tomography scan 2 days after ablation showed a 20-mm cystic ablation area in the pancreatic head (▶ **Fig. 3**). The patient developed no further bleeding during 10 weeks of follow-up.

Whereas endoscopic ultrasound-guided RFA is under evaluation for the ablation of pancreatic lesions, we are not aware of any previous case of direct intraductal application of RFA [1]. Intraductal RFA has the potential to treat complications of intraductal tumor growth such as bleeding. Bleeding from the pancreatic duct often requires radiological or surgical intervention [2]. In the present case, RFA was used to treat bleeding. However, RFA may also be a treatment option for the treatment of small intraductal neoplasms. At present, intraductal RFA is approved for the treatment of malignant biliary strictures [3] and is an alternative to photodynamic therapy [4].

Endoscopy_UCTN_Code_TTT_1AR_2AK

Competing interests: None

**Jochen Weigt, Ane Kandulski,
Peter Malfertheiner**

Department of Gastroenterology,
Hepatology, and Infectious Diseases,
Otto-v.-Guericke University, Magdeburg,
Germany

References

- 1 Kim HJ, Seo DW, Hassanuddin A et al. EUS-guided radiofrequency ablation of the porcine pancreas. *Gastrointest Endosc* 2012; 76: 1039–1043
- 2 Han B, Song ZF, Sun B. Hemosuccus pancreaticus: a rare cause of gastrointestinal bleeding. *Hepatobiliary Pancreat Dis Int* 2012; 11: 479–488
- 3 Tal AO, Vermehren J, Friedrich-Rust M et al. Intraductal endoscopic radiofrequency ablation for the treatment of hilar non-resectable malignant bile duct obstruction. *World J Gastrointest Endosc* 2014; 6: 13–19
- 4 Strand DS, Cosgrove ND, Patrie JT et al. ERCP-directed radiofrequency ablation and photodynamic therapy are associated with comparable survival in the treatment of unresectable cholangiocarcinoma. *Gastrointest Endosc*. In press 2014. doi: 10.1016/j.gie.2014.02.1030

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377590>
Endoscopy 2014; 46: E489–E490
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Jochen Weigt, MD

Department of Gastroenterology, Hepatology,
and Infectious Diseases
Otto-v.-Guericke University
Leipziger Str 44
Magdeburg 39120
Germany
Fax: +49-391-6713105
jochen.weigt@med.ovgu.de