Mantle cell lymphoma masked by posttraumatic pancreatic hematoma

Fig. 1 Evolving peripancreatic lesion. **a** Abdominal computed tomography immediately after a car accident shows an enhancing peripancreatic mass. **b** After 6 months, the lesion has shrunk. **c** A huge peripancreatic mass is incidentally identified following appendectomy 18 months after the injury.

Pelvic computed tomography revealed a 15-cm peripancreatic mass in a 73-year-old man undergoing an operation for appendicitis associated with a periappendiceal abscess. The patient had been hospitalized 18 months previously for blunt abdominal trauma sustained during a motor vehicle collision. At that time, hemoperitoneum and a peripancreatic hematoma were diagnosed (● Fig. 1 a). The hematoma decreased in size within 1 week, and there was no evidence of major pancreatic duct injury; thus, he was managed conservatively. The lesion continued to shrink for 6 months after the injury (● Fig. 1 b), but the volume had increased at the time of the appendectomy, as shown by computed tomography (● Fig. 1 c). Endoscopic ultrasound-guided biopsy with a 19-gauge needle (ProCore; Cook Medical, Bloomington, Indiana, USA) was used to identify the lesion. Pathologic examination and immunohistochemical staining of the specimen yielded a diagnosis of mantle cell lymphoma (● Fig. 2 a–c).

Pancreatic injury, which occurs in approximately 5% of patients who sustain blunt abdominal trauma, is associated with a morbidity rate of up to 45% and a mortality rate of up to 30% [1]. Low grade blunt injuries without main pancreatic ductal damage are initially managed non-operatively rather than through operative exploration and repair [2]. The differentiation between an organizing hematoma and a mass like a lymphoma is not always straightforward [3,4]. Notably, the mass in this case became smaller during the first 6 months. The mantle cell lymphoma, which presented as a slowly growing mass, was stationary in the early stage, and absorption of the hematoma occurred more quickly than growth of the lymphoma [5].

This case highlights the fact that clinicians should continue observation until a hematoma that has arisen at an unusual site completely disappears. If the behavior of a lesion does not correspond to the patient's clinical presentation or exhibits

an unusual course, the patient must be evaluated to establish the correct diagnosis.

Endoscopy_UCTN_Code_CCL_1AZ_2AO

Competing interests: None

**Seungho Lee¹, Jung-Ho Han¹,
Ki Bae Kim¹, Seon Mee Park¹,
Min Ho Kang², Hye-Suk Han³,
Song-Yi Choi⁴**

¹ Department of Gastroenterology, Chungbuk National University College of Medicine, Cheongju, Republic of Korea

² Department of Radiology, Chungbuk National University College of Medicine, Cheongju, Republic of Korea

³ Department of Oncology, Chungbuk National University College of Medicine, Cheongju, Republic of Korea

⁴ Departments of Pathology and Oncology, Chungnam National University College of Medicine, Daejeon, Republic of Korea

Fig. 2 Pathologic examination and immunohistochemical staining of mantle cell lymphoma. **a** The specimen contains small lymphocytes with irregular nuclei. **b, c** Immunohistochemical staining reveals positivity for cyclin D1 and CD5.

Acknowledgment

▼
This study was supported by a grant from the Korea Healthcare Technology R&D Project, Ministry of Health & Welfare, Republic of Korea (A100054). The source of the grant had no role in the design, practice, or analysis of the study.

References

- 1 Subramanian A, Dente CJ, Feliciano DV. The management of pancreatic trauma in the modern era. *Surg Clin North Am* 2007; 87: 1515–1532
- 2 Lee PH, Lee SK, Kim GU et al. Outcomes of hemodynamically stable patients with pancreatic injury after blunt abdominal trauma. *Pancreatology* 2012; 12: 487–492
- 3 Iaccarino C, Schiavi P, Crafa P et al. Primary dural lymphoma mimicking a chronic epidural hematoma. Differential diagnosis of two rare conditions. *Clin Neurol Neurosurg* 2013; 115: 1510–1513
- 4 Dedekam E, Graham J, Streng K et al. Primary renal lymphoma mimicking a subcapsular hematoma: a case report. *J Radiol Case Rep* 2013; 7: 18–26
- 5 Argatoff LH, Connors JM, Klasa RJ et al. Mantle cell lymphoma: a clinicopathologic study of 80 cases. *Blood* 1997; 89: 2067–2078

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377597>
Endoscopy 2014; 46: E520–E521
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Joung-Ho Han, MD, PhD
Department of Gastroenterology
Chungbuk National University College of Medicine
410 SungBong-Ro Heungdeok-Gu
Cheongju-Si ChungBuk, 361-711
Republic of Korea
Fax: +82-43-273-3252
joungho@cbnu.ac.kr